

ESOMEPRAZOLE MAGNESIUM- esomeprazole magnesium capsule, delayed release

Dr. Reddy's Laboratories Inc

Drug Facts

Active ingredient(s) in each capsule

* Esomeprazole 20 mg

(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate, USP)

Purpose

Acid reducer

Use(s)

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- Do not use if you are allergic to esomeprazole
- Esomeprazole may cause severe skin reactions. Symptoms may include:
- skin reddening
- blisters
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadness
- frequent **chest pain.**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss

- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read directions and warnings before use
- keep the carton. It contains important information.
- store at 20°C to 25°C (68°F to 77°F)
- this product meets USP Dissolution Test 3

Inactive ingredients

FD&C Blue #1, ferrous ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, iron oxide red, magnesium stearate, methacrylic acid copolymer, polysorbate 80, potassium hydroxide, propylene glycol, shellac, simethicone, sugar spheres, talc, titanium dioxide and triethyl citrate.

Questions or comments

call toll-free weekdays 8 AM to 8 PM EST at **1-888-375-3784**

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Distributor:

Dr. Reddy's Laboratories Inc.,

Princeton, NJ 08540

Made in India

Issued: 04/2024

Principal Display Panel

Container Labeling

Mini Capsules

Treats Frequent Heartburn

Esomeprazole Magnesium

Delayed-Release Capsules
USP, 20 mg* | ACID REDUCER
 May take 1 to 4 days for full effect

14 Capsules
 One 14-Day Course of Treatment



TAMPER EVIDENT-FEATURES: Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or blue band around the center of each capsule is broken or missing.

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Active ingredient Purpose (in each capsule)

Esomeprazole 20 mg... Acid reducer (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate, USP)

Uses ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: ■ Do not use if you are allergic to esomeprazole
 ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
 If an allergic reaction occurs, stop use and seek medical help right away.

Distributed by:
 Dr. Reddy's Laboratories, Inc.
 Princeton, NJ 08540

Made in India

Revised: 11/24

LOT/EXP 15 0085 106

PEEL HERE 

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if

- you have ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 - frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain
- Ask a doctor or pharmacist before use if you are** ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens ■ you need to take this product for more than 14 days
 - you need to take more than 1 course of treatment every 4 months ■ you get diarrhea
 - you develop a rash or joint pain
- if pregnant or breast-feeding, ask a health professional before use.**
Keep out of reach of children.
 In case of overdose, get medical help.

help or contact a Poison Control Center right away (1-800-222-1222)

Directions

- adults 18 years of age and older
 - this product is to be used once a day (every 24 hours), every day for 14 days
 - may take 1 to 4 days for full effect
- 14-Day Course of Treatment**
- swallow 1 capsule with a glass of water before eating in the morning
 - take every day for 14 days
 - do not take more than 1 capsule a day
 - swallow whole. Do not crush or chew capsules.

■ do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information. ■ store at 20°C to 25°C (68°F to 77°F) ■ this product meets USP Dissolution Test 3

Inactive ingredients

FD&C Blue #1, ferrous ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, iron oxide red, magnesium stearate, methacrylic acid copolymer, polysorbate 80, potassium hydroxide, propylene glycol, shellac, simethicone, sugar spheres, talc, titanium dioxide and triethyl citrate.

Questions or comments?

call toll-free weekdays 8 AM to 8 PM EST at 1-888-375-3784

Carton

7 0 1 5 8 0 0 5 1



Drug Facts
Active ingredient (in each capsule)
Esomeprazole, 20 mg
Acid reducer
Esomeprazole magnesium USP

Uses
● treats frequent heartburn (occurs 2 or more days a week)
● not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert
● Do not use if you are allergic to esomeprazole
● Esomeprazole may cause severe skin reactions. Symptoms may include: ● skin redness ● blisters ● rash
● If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:
● trouble or pain swallowing food, vomiting with blood, or bloody or black stools
● heartburn with **lightheadedness, swelling or dizziness**
● chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
● frequent chest pain
● had heartburn over 3 months. This may be a sign of a more serious condition.
● frequent wheezing, particularly with heartburn

Ask a doctor or pharmacist before use if you are:
● taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:
● your heartburn continues or worsens
● you need to take this product for more than 14 days

1 5 0 0 8 5 1 0 7



Drug Facts (continued)
● you need to take more than 1 course of treatment every 4 months you get diarrhea ● you develop a rash or joint pain
● if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
● adults 18 years of age and older
● this product is to be used once a day (every 24 hours), every day for 14 days
● most take 1 to 4 days for full effect

14-Day Course of Treatment
● swallow 1 capsule with a glass of water before eating in the morning
● take every day for 14 days
● do not take more than 1 capsule a day
● swallow whole. Do not crush or chew capsules.
● do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
● you may repeat a 14-day course every 4 months
● do not take for more than 14 days or more often than every 4 months unless directed by a doctor

Other information
● read the directions and warnings before use
● keep the carton. It contains important information.
● store at 20° to 25°C (68° to 77°F).
● this product meets USP Dissolution Test 3

Inactive ingredients
FD-C-Blue #1, Microfines oxide, gelatin, glycerol monostearate, hydroxypropyl cellulose, hypromellose, iron oxide red, magnesium stearate, methacrylic acid copolymer, polyacrylate 80, potassium hydroxide, propylene glycol, stearic, ammonium, sugar spheres, talc, titanium dioxide and hydroxy citrate.

Questions or comments?
Call toll-free weekdays 8 AM to 5 PM EST at 1-888-375-3784

3 BOTTLES INSIDE



Actual Size

HealthCare  Aisle

NDC 43598-067-42

Mini Capsules

Esomeprazole Magnesium
Delayed-Release Capsules USP, 20 mg*
ACID REDUCER

HealthCare  Aisle

Mini Capsules

Treats Frequent Heartburn

Esomeprazole Magnesium
Delayed-Release Capsules USP, 20 mg*
ACID REDUCER

May take 1 to 4 days for full effect
See new warning information

42 Capsules
Three 14-Day Courses of Treatment

LOT
EXP



Compare to the active ingredient in Nexium® 24HR Clear Minis™

Compare to the active ingredient in Nexium® 24HR Clear Minis™

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

KEEP CAUTION FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION
TAMPER EVIDENT FEATURES: Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" or blue band around the center of each capsule is broken or missing.

****This product is not manufactured or distributed by GSK Consumer Healthcare, distributor of Nexium® 24HR. Nexium® is a registered trademark of AstraZeneca AB. Clinically Proven to Treat Frequent Heartburn. Distributed by: Dr. Reddy's Laboratories, Inc. Princeton, NJ 08540**

Revised: 11/24

ESOMEPRAZOLE MAGNESIUM

esomeprazole magnesium capsule, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:43598-067
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ESOMEPRAZOLE MAGNESIUM (UNII: R6DXU4WAY9) (ESOMEPRAZOLE - UNII:N3PA6559FT)	ESOMEPRAZOLE	20 mg

Inactive Ingredients

Ingredient Name	Strength
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:2) (UNII: XRK36F13ZZ)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B710)	
TALC (UNII: 7SEV7J4R1U)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
GELATIN (UNII: 2G86QN327L)	

FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
SUCROSE (UNII: C151H8M554)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
DIMETHICONE 350 (UNII: 2Y53S6ATLU)	

Product Characteristics

Color	BLUE (light blue cap and dark blue body)	Score	no score
Shape	CAPSULE	Size	11mm
Flavor		Imprint Code	926
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43598-067-42	3 in 1 CARTON	12/01/2024	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA207198	12/01/2024	

Labeler - Dr. Reddy's Laboratories Inc (802315887)