


GASTROGRAFIN- diatrizoate meglumine and diatrizoate sodium liquid
Bracco Diagnostics Inc

GASTROGRAFIN®
Diatrizoate Meglumine
and Diatrizoate Sodium
Solution USP


DESCRIPTION

Gastrografin (Diatrizoate Meglumine and Diatrizoate Sodium Solution) is a palatable lemon-flavored water-soluble iodinated radiopaque contrast medium for oral or rectal administration only. Each mL contains 660 mg diatrizoate meglumine and 100 mg diatrizoate sodium; pH has been adjusted to 6.0 to 7.6 with sodium hydroxide. *Each mL contains approximately 4.8 mg (0.21 mEq) sodium and 367 mg organically bound iodine.* Inactive ingredients: edetate disodium, flavor, polysorbate 80, purified water, saccharin sodium, simethicone, and sodium citrate.

Diatrizoate meglumine is designated chemically as 1-deoxy-1-(methylamino)-D-glucitol 3,5-diacetamido-2,4,6-triiodo-benzoate (salt); diatrizoate sodium is monosodium 3, 5-diacetamido-2,4,6-triiodobenzoate. Structural formulas:



diatrizoate meglumine
 $C_{11}H_9I_3N_2O_4 \cdot C_7H_{17}NO_5$
MW 809.13
Organically Bound Iodine: 47.1%
CAS-131-49-7



diatrizoate sodium
 $C_{11}H_8I_3N_2NaO_4$
MW 635.90
Organically Bound Iodine: 59.9%
CAS-737-31-5

CLINICAL PHARMACOLOGY

The most important characteristic of contrast media is the iodine content. The relatively high atomic weight of iodine contributes sufficient radiodensity for radiographic contrast with surrounding tissues.

Diagnostic enteral radiopaque agents have few known pharmacological effects. Diatrizoate meglumine and diatrizoate sodium exert a mild laxative effect attributable to their high osmolarity.

Diatrizoate meglumine and diatrizoate sodium are sparingly absorbed from the intact gastrointestinal tract, and therefore permit gastrointestinal opacification and delineation after oral or rectal administration. Oral administration is used for radiographic evaluation of the esophagus, stomach and proximal small intestine. Rectal administration is used for examination of the colon; however, visualization of the distal small bowel is generally unsatisfactory, since the hypertonicity of the medium causes intraluminal diffusion of water with subsequent dilution of the medium. Enough absorption from the gastrointestinal tract to permit incidental visualization of the urinary tract has been reported; this should also be considered when thyroid testing is being contemplated, since iodine-mediated thyrotropic effects may occur (see **PRECAUTIONS**).

INDICATIONS AND USAGE

Gastrografin (Diatrizoate Meglumine and Diatrizoate Sodium Solution) is indicated for radiographic examination of segments of the gastrointestinal tract (esophagus, stomach, proximal small intestine, and colon). The preparation is particularly indicated when a more viscous agent such as barium sulfate, which is not water-soluble, is not feasible or is potentially dangerous.

Gastrografin may also be used as an adjunct to contrast enhancement in computed tomography of the torso (body imaging); the preparation is indicated, in conjunction with intravenous administration of a radiopaque contrast agent, when unenhanced imaging may not provide sufficient definition in distinguishing normal loops of bowel from adjacent organs or areas of suspected pathology.

CONTRAINDICATIONS

Do not administer to patients with a known hypersensitivity to Gastrografin or any of its components.

WARNINGS

Dehydration: Administration of hypertonic Gastrografin solutions may lead to hypovolemia and hypotension due to fluid loss from the intestine. A 1 in 4.6 (1:4.6) dilution of Gastrografin yields an approximately isotonic 16.5 percent diatrizoate salts solution; less dilute solutions are hypertonic and may lead to intraluminal movement of fluid with resulting hypovolemia. In young or debilitated children and in elderly cachectic persons, the loss of plasma fluid may be sufficient to cause a shock-like state. If Gastrografin is used in infants and children (under 10 kg) or in dehydrated or debilitated patients, the solution must be prepared using the specific dilutions described in **DOSAGE AND ADMINISTRATION**. In debilitated patients and in patients with electrolyte imbalances, postprocedural monitoring of hydration, serum osmolarity, electrolytes and clinical status is essential. In pediatric or severely debilitated patients, the maintenance of an open intravenous fluid line for rehydration may be advisable should hypotension or shock supervene. Electrolyte disturbances must be corrected prior to the administration of any hypertonic Gastrografin solutions.

Aspiration: Aspiration of Gastrografin into the trachea and airways may result in serious pulmonary complications including, pulmonary edema, pneumonitis or death. Bronchial entry of any orally administered contrast medium causes a copious osmotic effusion. Therefore, avoid use of Gastrografin in patients with esophagotracheal fistula and minimize risks for pulmonary aspiration in all patients. If Gastrografin is given by nasogastric tube, the position of the tube in the stomach must be verified before administration.

Anaphylactic reactions: Anaphylactic reactions, including fatalities, have been reported with the use of Gastrografin. Patients at increased risk include those with a history of a previous reaction to a contrast medium, patients with a known sensitivity to iodine, and patients with a known clinical hypersensitivity (bronchial asthma, hay fever, and food allergies). Medical personnel trained in the treatment of anaphylactic reactions and the necessary drugs and medical equipment should always be readily available when Gastrografin is used.

PRECAUTIONS

General

Diagnostic procedures which involve the use of radiopaque contrast agents should be carried out under the direction of personnel with the prerequisite training and with a thorough knowledge of the particular procedure to be performed. Appropriate facilities should be available for coping with any complication of administration, as well as for treatment of reaction to the contrast medium (see **ADVERSE REACTIONS**, and **PRECAUTIONS, Information for the Patient**).

Rectal administration of undiluted Gastrografin (Diatrizoate Meglumine and Diatrizoate Sodium Solution) in any patient, particularly with large doses and/or in those with overdistention, has been reported to be associated with mucosal irritation.

Cases of hyperthyroidism have been reported with the use of oral contrast media. Some of these patients reportedly had multinodular goiters which may have been responsible for the increased hormone synthesis in response to excess iodine. Administration of an intravascular iodinated radiopaque diagnostic agent to a hyperthyroid patient precipitated thyroid storm; a similar situation could follow administration of oral preparations of iodides. Therefore, caution should be exercised when administering enteral gastrointestinal radiopaque agents to hyperthyroid and euthyroid goiterous patients.

Consideration should be given to the potential for precipitation of water-soluble contrast agents under conditions that may promote hyperacidity (i.e., fasting, emotional upset, or stress). Harmful effects directly attributable to precipitate formation have not been reported. However, the possibility of interpreting the precipitate radiologically as an anatomical abnormality (i.e., ulceration of the stomach or small intestine) or injury, should be kept in mind.

Information for the Patient

Patients should receive the following information and instructions:

1. This drug has been prescribed to perform an x-ray of the gastrointestinal tract.
2. Inform the physician if pregnant or if allergic to iodine, any foods, or x-ray materials.
3. The iodine in diatrizoate salts may interfere with some thyroid tests if these are needed in the future. Inform the attending physician at that time about this gastrointestinal study.
4. This drug may cause abdominal cramping, nausea, vomiting, diarrhea, skin rashes, itching, heartburn, dizziness, or headache in some patients, but most reactions are mild and pass quickly.

Drug/Laboratory Test Interactions

Thyroid Function Tests

The results of protein bound iodine (PBI) and radioactive iodine uptake studies, which depend on iodine estimations, will not accurately reflect thyroid function for six months, and possibly as long as one year, following the administration of diagnostic enteral radiopaque media.

Thyroid function tests, if indicated, generally should be performed prior to the administration of any iodinated agent. However, thyroid function can be evaluated after use of these agents by using T₃ resin uptake and total or free thyroxine (T₄) assays which are not dependent on iodine estimations.

Pancreatic Tests

Small quantities of contrast medium in the intestinal tract may cause false low trypsin values when determined spectrophotometrically. Therefore, duodenal instillation should not precede pancreatic function tests involving spectrophotometric trypsin assays.

Any test which might be affected by contrast media should be performed prior to administration of the

contrast medium.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have not been performed to evaluate carcinogenic or mutagenic potential, or possible impairment of fertility in males or females.

Pregnancy

When administered intravenously, diatrizoate salts cross the placenta and are evenly distributed in fetal tissues.

No teratogenic effects attributable to diatrizoate meglumine or diatrizoate sodium have been observed in teratology studies performed in animals. There are, however, no adequate and well-controlled studies in pregnant women. Because small amounts of these agents may be absorbed, and animal teratology studies are not always predictive of human response, these agents should be used during pregnancy only when clearly needed.

Procedures including radiation involve a certain risk related to the exposure of the fetus.

Nursing Mothers

Diatrizoate meglumine is excreted in breast milk following intravascular administration.

Because small amounts of enteral gastrointestinal radiopaque agents may be absorbed following oral or rectal administration, caution should be exercised when they are administered to a nursing woman.

Pediatric Use

See **WARNINGS**, and **PRECAUTIONS, General**.

Local injury to the colonic mucosa, particularly in the presence of underlying disease which interferes with intestinal viability, has been reported in cases where recommended doses and dilutions (see **DOSAGE AND ADMINISTRATION**) were not used; when extemporaneous dosage is elected, the polysorbate 80 level in the dose may be a contributing factor to injury.

ADVERSE REACTIONS

Most adverse reactions to enteral diagnostic radiopaque agents are mild and transitory. Nausea, vomiting and/or diarrhea, urticaria with erythema, hypoxia, acute dyspnea, tachyarrhythmia, and anaphylaxis have occurred following ingestion of the contrast medium, particularly when high concentrations of large volumes of solution are administered. Severe changes in serum osmolarity and electrolyte concentrations may produce shock-like states (see **WARNINGS**). It should be kept in mind that serious or anaphylactoid reactions that may occur with intravascular administration of radiopaque contrast agents are theoretically possible following administration by other routes.

OVERDOSAGE

See **WARNINGS** regarding potential hypovolemia, hypotension, or shock. The maintenance of an open intravenous fluid line for rehydration may be advisable. See **DOSAGE AND ADMINISTRATION** for appropriate doses and dilutions. Treatment of an overdose should be directed toward the support of all vital functions, and prompt institution of symptomatic therapy.

DOSAGE AND ADMINISTRATION

General

This medium is not to be used for the preparation of solutions for parenteral administration. Oral

or rectal administration only. Discard any unused portion after procedure.

The routine preparatory measures employed for barium studies are also appropriate for this agent.

For pediatric and severely cachectic patients the maintenance of an intravenous fluid line may be advisable.

Radiographic Examination of Segments of the Gastrointestinal Tract

Oral Administration: Adult oral dosage may range from 30 to 90 mL (11 to 33 g iodine), depending on the nature of the examination and the size of the patient. For infants and children less than 5 years of age, 30 mL (11 g iodine) are usually adequate; for children 5 to 10 years of age, the suggested dose is 60 mL (22 g iodine). These pediatric doses may be diluted 1:1, if desired, with water, carbonated beverage, milk, or mineral oil. When used in infants, the solution may be given in a nursing bottle. Pediatric doses may also be used in dehydrated and/or debilitated adult patients. A 1:1 dilution is also recommended when the contrast medium is used in elderly cachectic individuals.

For very young (under 10 kg) and debilitated children the dose should be diluted: 1 part Gastrografin (Diatrizoate Meglumine and Diatrizoate Sodium Solution) in 3 parts water is recommended.

For Enemas or Enterostomy Instillations: Gastrografin should be diluted when it is used for enemas and enterostomy instillations. When used as an enema, the suggested dilution for adults is 240 mL (88 g iodine) in 1,000 mL of tap water. For children under 5 years of age, a 1:5 dilution in tap water is suggested; for children over 5 years of age, 90 mL (33 g iodine) in 500 mL of tap water is a suitable dilution.

Tomography (Body Imaging)

A usual adult dose is 240 mL of a dilute Gastrografin solution prepared by diluting 25 mL (9.17 g iodine) to one liter with tap water. Less dilute solutions [up to 77 mL (28.26 g iodine) diluted to one liter with tap water] may be used when indicated. The dose is administered orally about 15 to 30 minutes prior to imaging in order to permit the contrast medium to reach the pelvic loops.

HOW SUPPLIED

Gastrografin (Diatrizoate Meglumine and Diatrizoate Sodium Solution USP) is available in packages of:

Twenty-four 30 mL single dose bottles (NDC 0270-0445-35).

Twelve 120 mL single dose bottles (NDC 0270-0445-40).

Storage

Protect from light. Store at 20-25°C (68-77°F) [See USP]; avoid excessive heat.

Manufactured by E-Z-EM Canada Inc,
Anjou, Quebec H1J2Z4, Canada
For Bracco Diagnostics Inc.
Monroe Township, NJ 08831

Rev. August 2019
CLE7903

Gastrografin® 120 mL Label
NDC 0270-0445-40



Bracco Diagnostics

120 mL

NDC 0270-0445-40

37% Organically Bound Iodine

GASTROGRAFIN®

Diatrizoate Meglumine
and Diatrizoate
Sodium Solution USP

Rx only

DO NOT INJECT

Single Dose Bottle
Discard Unused Portion
For gastrointestinal radiography
For oral or rectal use only.
Not for parenteral use

Usual Dosage: See package insert.
Each mL of aqueous, flavored
solution provides 660 mg
diatrizoate meglumine and
100 mg diatrizoate sodium; pH
adjusted to 6.0 – 7.6 with sodium
hydroxide. Each mL contains
approximately 4.8 mg (0.21 mEq)
sodium and 367 mg organically
bound iodine.

Protect from light • Store at
20-25°C (68-77°F) [See USP];
avoid excessive heat. Dispense
in tight, light-resistant containers

Manufactured by
E-Z-EM Canada Inc.
Anjou, Quebec H1J2Z4, Canada
For Bracco Diagnostics Inc.
Monroe Twp., NJ 08831

REV. 08/19



N 3 0270-0445-40 3



LOT

EXP

C1E7B02

Gastrografin® 12 x 120 mL Bottles Label
NDC 0270-0445-40

NDC 0270-0445-40

12 x 120 mL bottles



Bracco Diagnostics

37% Organically Bound Iodine

GASTROGRAFIN®

Diatrizoate Meglumine and Diatrizoate Sodium Solution USP

Rx only

Bulk Container – Not for Household Dispensing

Single Dose Bottles • Discard Unused Portion • For gastrointestinal radiography

• For oral or rectal use only • Not for parenteral use

DO NOT INJECT

Usual dosage: See package insert.

Each mL of aqueous, flavored solution provides 660 mg diatrizoate meglumine and 100 mg diatrizoate sodium; pH adjusted to 6.0-7.6 with sodium hydroxide. Each mL contains approximately 4.8 mg (0.21 mEq) sodium and 367 mg organically bound iodine.

Protect from light • Store at 20-25°C (68-77°F)

[See USP]; avoid excessive heat.

Dispense in tight, light-resistant containers

LOT

Lot and expiry encoding area

EXP

Manufactured by

E-Z-EM Canada Inc.

Anjou, Quebec H1J2Z4, Canada

For Bracco Diagnostics Inc.

Monroe Twp., NJ 08831



(01)30302700445404

REV. 08/19

CEE7A02



GASTROGRAFIN

diatrizoate meglumine and diatrizoate sodium liquid

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0270-0445
Route of Administration	ORAL, RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
diatrizoate meglumine (UNII: 3X9MR4N98U) (DIATRIZOIC ACID - UNII:5UVC90J1LK)	diatrizoate meglumine	660 mg in 1 mL
diatrizoate sodium (UNII: V5403H8VG7) (DIATRIZOIC ACID - UNII:5UVC90J1LK)	diatrizoate sodium	100 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CITRATE (UNII: 1Q73Q2JULR)	3.22 mg in 1 mL
EDETATE DISODIUM (UNII: 7FLD91C86K)	0.403 mg in 1 mL
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	2.77 mg in 1 mL

Product Characteristics

Color		Score	
Shape		Size	
Flavor	LEMON	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0270-0445-35	24 in 1 BOX	04/01/2005	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0270-0445-40	12 in 1 BOX	02/26/1958	
2		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA011245	02/26/1958	

Labeler - Bracco Diagnostics Inc (849234661)

Registrant - Bracco Diagnostics Inc (849234661)

Establishment

Name	Address	ID/FEI	Business Operations
E-Z-EM Canada Inc		204211163	MANUFACTURE(0270-0445) , ANALYSIS(0270-0445)

Establishment

Name	Address	ID/FEI	Business Operations
Justesa Imagen, S.A.U		477020325	LABEL(0270-0445) , PACK(0270-0445) , API MANUFACTURE(0270-0445) , ANALYSIS(0270-0445)

Establishment

Name	Address	ID/FEI	Business Operations
Merck SLU		460082644	LABEL(0270-0445) , PACK(0270-0445) , ANALYSIS(0270-0445) , API MANUFACTURE(0270-0445)

