

MILK OF MAGNESIA- magnesium hydroxide suspension

Major Pharmaceuticals

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Major Pharmaceuticals Milk of Magnesia Drug Facts

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not exceed the maximum recommended daily dose in a 24 hour period

- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- **each 15 mL contains:** calcium 20 mg, magnesium 595 mg, and sodium 2 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- close cap tightly after use
- avoid freezing

Inactive ingredients

light mineral oil, peppermint oil, purified water, saccharin sodium

Questions or comments?

1-800-616-2471

Principal Display Panel

Milk of Magnesia Saline Laxative

MINT FLAVOR

Effective overnight relief

Cramp free

Stimulant free

Sugar free

Compare to the Active Ingredient of Phillips'® Milk of Magnesia

12 FL OZ (355 mL)

Magnesium Hydroxide, 1200 mg per 15 mL



NDC 0904-6757-14

Milk of Magnesia

Saline Laxative MINT FLAVOR

- Effective overnight relief
- Cramp free
- Stimulant free
- Sugar free

Compare to the Active Ingredient of Phillips[®] Milk of Magnesia*

12 FL OZ (355 mL)

Magnesium Hydroxide,
1200 mg per 15 mL

: 33240 5C F1

Drug Facts**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.****Active ingredient (in each 15 mL)**

Magnesium hydroxide 1200 mg.....

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug.

Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).**Directions**

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older 30 mL to 60 mL

children 6 to 11 years 15 mL to 30 mL

children under 6 years ask a doctor

Other information

- each 15 mL contains: calcium 20 mg, magnesium 595 mg, and sodium 2 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- close cap tightly after use
- avoid freezing

Inactive ingredients

light mineral oil, peppermint oil, purified water, saccharin sodium

Questions or comments? 1-800-616-2471

T8 05 0H2EE :

**Gluten Free
Distributed by
MAJOR[®]**

PHARMACEUTICALS
17177 N Laurel Park Drive
Suite 233
Livonia, MI 48152
M-05 REV. 04/18
Re-Order No. 700956

*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips[®] Milk of Magnesia.**MILK OF MAGNESIA**

magnesium hydroxide suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-6757
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
WATER (UNII: 059QF0K00R)	

SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
PEPPERMINT OIL (UNII: AV092KU4JH)	

Product Characteristics

Color	WHITE (opaque)	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-6757-14	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/15/1991	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	04/15/1991	

Labeler - Major Pharmaceuticals (191427277)

Revised: 10/2018

Major Pharmaceuticals