

IBUPROFEN- ibuprofen tablet
BETTER LIVING BRANDS LLC.

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)*

SEE NEW WARNINGS

Active ingredient

(in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit

of aspirin

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

■ **do not take more than directed**

■ **the smallest effective dose should be used**

adults and children 12 years and older

■ take 1 tablet every 4 to 6 hours while symptoms persist

■ if pain or fever does not respond to 1 tablet, 2 tablets may be used

■ do not exceed 6 tablets in 24 hours, unless directed by a doctor

Children under 12 years

■ ask a doctor

Other information

■ store between 20-25°C (68-77°F). [See USP Controlled Temperature]

■ avoid excessive heat above 40°C (104°F)

Inactive ingredients

colloidal silicon dioxide, iron oxide red, maize starch, polyethylene glycol, povidone k-30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide.

Questions or comments?

call **1-877-770-3183** weekdays 8:00 AM EST to 5:00PM PST.

Principal Display Panel



Compare to Advil[®] Tablets active ingredient!
NDC 21130-396-25

Quality Guaranteed

Ibuprofen
Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)*
Contains no ingredient made from a gluten-containing grain (wheat, barley, or rye) Actual Size
250 Coated Tablets

Important: Read all product information before using. Do not use if foil liner seal is broken or missing.

Purposes
Pain reliever/fever reducer

Drug Facts
Active ingredient (in each tablet) Ibuprofen 200 mg (NSAID)*
Nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache
■ backache ■ the common cold ■ menstrual cramps
■ temporarily reduces fever

Warnings
Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
■ skin redness ■ rash ■ ulcers
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinner (anticoagulant) or aspirin drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
■ If you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
■ right before or after heart surgery
■ Ask a doctor before use if:
■ you have problems or serious side effects from taking pain relievers or fever reducers

Other information:
BETTER LIFE BRANDS, LLC
P.O. BOX 99
BIRMINGHAM, GA 30606-0099
1-888-752-3863

LOT 2113077744
EXP. 2025020403
700000005476

Scan here for more information

LOVE IT OR IT'S ON US!
This product is not manufactured or distributed by Mylan Consumer Healthcare, distributor of Advil[®] Tablets.

3 2113077744 1
MADE IN INDIA

nted label,
led line.

Inside (adhesive side)

Drug Facts (continued)

■ the stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are:
■ taking aspirin or heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
■ under a doctor's care for any serious condition
■ taking any other drug

When using this product:
■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if:
■ you experience any of the following signs of stomach bleeding:
■ red bark ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke: ■ chest pain
■ trouble breathing ■ weakness in one part or side of body
■ dazed speech ■ big swelling

■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
■ do not take more than directed
■ the smallest effective dose should be used

adults and children ■ take 1 tablet every 4 to 6 hours while symptoms last.
■ If pain or fever does not respond to 1 tablet, 2 tablets may be used.
■ do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years ■ ask a doctor

Other information
■ store between 20 to 25°C (68 to 77°F), (See USP Controlled Room Temperature).
■ avoid excessive heat above 40°C (104°F)

Inactive ingredients
colloidal silicon dioxide, iron oxide red, maize starch, polyethylene glycol, polydioxane k-50, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?
call 1-877-776-8125 weekdays 8:00 AM EST to 5:00 PM PST

IBUPROFEN

ibuprofen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-396
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFI6D95)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	brown (Reddish Brown color Tablets)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-396-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	07/12/2019	
2	NDC:21130-396-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	08/18/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	07/12/2019	

Labeler - BETTER LIVING BRANDS LLC. (009137209)

Revised: 12/2025

BETTER LIVING BRANDS LLC.