

**ACETAMINOPHEN DIPHENHYDRAMINE HCL- acetaminophen diphenhydramine  
hcl tablet  
Granules USA, Inc.**

-----  
***EXTRA STRENGTH***

**Pain Relief PM**

***Acetaminophen, USP 500 mg/ Diphenhydramine HCl, USP 25 mg  
Pain Reliever/ Nighttime Sleep-Aid***

***Non-Habit Forming***

**Active ingredients**

(in each caplet)

Acetaminophen, USP 500mg

Diphenhydramine HCl. 25mg

**Purposes**

Pain reliever

Nighttime sleep-aid

**Uses**

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness.

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

### **Ask a doctor or pharmacist before use if you have**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

### **Stop use and ask a doctor if**

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
- These could be signs of a serious condition.

### **If pregnant or breast-feeding**

ask a health professional before use.

### **Keep out of the reach of children.**

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

■ do not take more than directed (see overdose warning)

adults and children 12 years and over

■ take 2 caplets at bedtime do not use

■ do not take more than 2 caplets of this product in 24 hours

children under 12 years

■ do not use

## **Other information**

■ store between 20°-25°C (68°-77°F). See USP Controlled Room Temperature.

■ see end panel for expiration date and lot number


## **Inactive ingredients**

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, hypromellose, microcrystalline cellulose, polyethylene glycol, polysorbate, povidone, pregelatinized starch, stearic acid, titanium dioxide

## **Questions or comments?**

item	:CNS APAP RR 50C1		
code #	:		
size	:1+3/4 X 1+3/4 X 3+3/8	view	:PRIM
ref #	:PPI000045	date	:06/01
material	:.016 888		






**Dion modifications to side 2 incoming artwork:**

1) Adjusted size of bounding box so text does not extend into it

**IGS**

DE 1





# ACETAMINOPHEN DIPHENHYDRAMINE HCL

acetaminophen diphenhydramine hcl tablet

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69848-014
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)		DIPHENHYDRAMINE HYDROCHLORIDE	25 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	500 mg

## Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	OVAL (CAPLET SHAPED TABLET)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	G;651
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69848-014-02	20 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2019	
2	NDC:69848-014-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2019	
3	NDC:69848-014-15	150 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2019	

**Labeler** - Granules USA, Inc. (137098864)

Revised: 12/2024

Granules USA, Inc.