

**CONRX NIGHTTIME- acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate tablet**

**Eagle Distributors, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**ConRx® NightTime**

**Drug Facts**

<b>Active ingredients (in each Caplet)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

**Ask a doctor before use if you have**

- liver disease
- glaucoma

- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers.
- taking the blood thinning drug warfarin

**When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed - see Overdose warning
- do not exceed 4 doses per 24 hours

adult and children 12 years and over, 2 Caplets with water every 6 hours

children 4 to under 12 years, ask a doctor

children under 4 years, do not use

**Other information**

- store at room temperature

**Inactive ingredients**

cross carmellose sodium, D&C Yellow No.10, FD&C Blue No.1, cellulose microcrystalline, polysorbate 80, silicone dioxide, talc, titanium dioxide, corn starch, gelatin, magnesium stearate, sodium starch glycolate, sodium benzoate, hydroxy propyl methyl

## Questions?

1-855-619-7900

### **PRINCIPAL DISPLAY PANEL - 50 x 2 Caplet Pouch Box**

**Compare to the Active Ingredients in  
NyQuil®**

*ConRx® NightTime*

**COLD & FLU**

**NightTime Relief**

*Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr*

- ▣ **Aches, Fever & Sore Throat**
- ▣ **Sneezing, Runny Nose**
- ▣ **Cough**

*2 Caplets Each Pouch*

**TO OPEN**

**PUSH IN TAB AND PULL OUT**

**Compare to the Active Ingredients in  
NyQuil®**

*2 Caplets Each Pouch*

Compare to the Active Ingredients in

# NyQuil<sup>®</sup>

## ConRx<sup>®</sup> NightTime

**COLD & FLU**

NightTime Relief

Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

- ★ Aches, Fever & Sore Throat
- ★ Sneezing, Runny Nose
- ★ Cough

2 Caplets Each Pouch

TO OPEN  
PUSH IN TAB AND PULL OUT

Compare to the Active Ingredients in

# NyQuil<sup>®</sup>

2 Caplets Each Pouch

Compare to the Active Ingredients in

# NyQuil<sup>®</sup>

## ConRx<sup>®</sup> NightTime

**COLD & FLU**

NightTime Relief

Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

- ★ Aches, Fever & Sore Throat
- ★ Sneezing, Runny Nose
- ★ Cough

2 Caplets Each Pouch

★ Cough  
★ Sneezing, Runny Nose  
★ Aches, Fever & Sore Throat  
Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr  
NightTime Relief

**COLD & FLU**

# NyQuil<sup>®</sup>

## ConRx<sup>®</sup> NightTime

Compare to the Active Ingredients in

Failure to follow these warnings could result in serious consequences.

### Drug Facts

#### Active ingredients (in each Caplet)

Acetaminophen 325 mg ..... Pain reliever/fever reducer

Dextromethorphan HBr 15 mg ..... Cough suppressant

#### Purpose

Compare to the Active Ingredients in

# NyQuil<sup>®</sup>

Dextromethorphan HBr 10 mg Cough suppressant  
 Doxylamine succinate 6.25 mg Antihistamine

**Uses:** temporarily relieves common cold/flu symptoms:  
 • cough due to minor throat and bronchial irritation • sore throat • headache • minor aches and pains  
 • fever • runny nose and sneezing

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
 • more than 4 doses in 24 hrs, which is the maximum daily amount for this product  
 • with other drugs containing acetaminophen  
 • 3 or more alcoholic drinks every day while using this product  
**Sore throat warning:**  
 If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use:**  
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 • if you are now taking prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 • to make a child sleep

**Ask a doctor before use if you have** • liver disease • glaucoma • cough that occurs with too much phlegm (mucus)  
 • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** • taking sedatives or tranquilizers. • taking the blood thinning drug warfarin

**When using this product**  
 • do not use more than directed  
 • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast feeding, ask a health professional before use.**  
**Keep out of reach of children.**  
**Overdose warning:** Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs of symptoms.

**Directions** • take only as directed - see Overdose warning  
 • do not exceed 4 doses per 24 hours

adult and children 12 years and over	2 Caplets with water every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

**Other information** • store at room temperature

**Inactive ingredients** cross carmellose sodium, D&C Yellow No.10, FD&C Blue No.1, cellulose microcrystalline, polysorbate 80, silicone dioxide, talc, titanium dioxide, corn starch, gelatin, magnesium stearate, sodium starch glycolate, sodium benzoate, hydroxy propyl methyl

**Questions?** 1-855-619-7900

**NyxQuil**  
**ConRx® NightTime**  
**COLD & FLU**  
**NightTime Relief**

*Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr*

- ★ Aches, Fever & Sore Throat
- ★ Sneezing, Runny Nose
- ★ Cough

\*This product is not manufactured or distributed by: Proctor & Gamble. Eagle Distributors, Inc./ConRx® does not own the NyQuil® trademark.

  
 Product manufactured for:  
**Eagle Distributors, Inc.**  
 Los Angeles, CA 90011

## CONRX NIGHTTIME

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68737-239
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Acetaminophen</b> (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	325 mg
<b>Dextromethorphan Hydrobromide</b> (UNII: 9D2RTI9KYH) (Dextromethorphan - UNII:7355X3ROTS)	Dextromethorphan Hydrobromide	15 mg
<b>Doxylamine Succinate</b> (UNII: V9B19B5YI2) (Doxylamine - UNII:95QB77JKPL)	Doxylamine Succinate	6.25 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D6IU)	

<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)
<b>GELATIN</b> (UNII: 2G86QN327L)
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6B30)
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)
<b>ETHYLCELLULOSES</b> (UNII: 7Z8S9VYZ4B)

### Product Characteristics

<b>Color</b>	GREEN	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	CRX
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68737-239-23	50 in 1 BOX		
1		2 in 1 POUCH		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	02/10/2014	

**Labeler** - Eagle Distributors, Inc. (929837425)

Revised: 2/2014

Eagle Distributors, Inc.