

**NAPROXEN SODIUM- naproxen sodium tablet
Granules USA, Inc.**

Naproxen Sodium Tablets, USP 220 mg (NSAID)
Pain reliever/fever reducer
*STRENGTH TO LAST 12 HOURS***

Active ingredient (For Tablet)

(in each tablet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)**

**nonsteroidal anti-inflammatory drug

Active ingredient (For Caplet)

(in each caplet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)**

**nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Allergy alert:

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
 - it feels like the pill is stuck in your throat
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions (For Tablets)

- **do not take more than directed**
- **the smallest effective dose should be used**

■ drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12- hour period
- do not exceed 3 tablets in a 24- hour period

Children under 12 years

- ask a doctor

Directions (For Caplets)

- **do not take more than directed**
- **the smallest effective dose should be used**

■ drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12- hour period
- do not exceed 3 caplets in a 24- hour period

Children under 12 years

- ask a doctor

Other information (For Tablet)

- **each tablet contains:** sodium 20 mg
- store at 20 - 25°C (68 - 77°F). Avoid high humidity and excessive heat above 40°C (104°F)

Other information (For Caplet)

■ **each caplet contains:** sodium 20 mg

■ store at 20 - 25°C (68 - 77°F). Avoid high humidity and excessive heat above 40°C (104°F)

Inactive ingredients

FD&C blue #2 aluminum lake, hypromellose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, povidone k-30, sodium starch glycolate, stearic acid, titanium dioxide.

Questions or comments?

1-877-770-3183 Mon - Fri 9:00 AM to 4:00 PM EST.

Principal Display Panel

size :1+3/4 X 1+3/4 X 3+3/8
 ref # :PPI80604B
 material :.016 SBS

Drug Facts (continued)

Other Information

- each tablet contains sodium 20 mg
- store at 20° - 25°C (68° - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients FDAC, PABA #2, aluminum lactate, hydroxyacetone 2910, mezastron, microcrystalline cellulose, polyethylene glycol, potassium hydroxide, sodium starch glycolate, stearic acid, titanium dioxide.

Questions or comments?
 1-877-776-3183 Mon-Fri 9:00 AM to 4:00 PM EST.

This product is not manufactured or distributed by Granules Health, LLC, distributor of Aleve. Aleve is a registered trademark of Bayer HealthCare, LLC.

COATING FREE AREA

700000002396
 200000003203

KEEP CARTON FOR REFERENCE
 DO NOT USE IF CARTON IS OPEN OR IF FOIL SEAL ON BOTTLE OPENING IS MISSING OR BROKEN

Drug Facts (continued)

naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have had high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools

Drug Facts (continued)

- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"> take 1 tablet every 8 to 12 hours while symptoms last for the first dose you may take 2 tablets within the first hour do not exceed 2 tablets in any 5- to 12-hour period do not exceed 3 tablets in a 24-hour period
Children under 12 years	ask a doctor

50 Tablets

STRENGTH TO LAST 12 HOURS

Contains no xerodent made from a gluten-containing grain (wheat, barley, or rye)

Naproxen Sodium Tablets USP
220 mg (NSAID)**
 Pain reliever/fever reducer

GRANULES
 CONSUMER HEALTH

***Compare to the Active Ingredient of Aleve® Tablets**



Drug Facts

Active ingredient (in each tablet)
 Naproxen sodium 220 mg. Pain reliever/fever reducer (naproxen 200 mg) (NSAID)**

Purposes
 *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, ...)

COATING FREE AREA

200000003203
 700000002396

Distributed by:
 Granules Consumer Health
 A Division of Granules USA, Inc.
 35 Waterway Blvd., 2nd Floor
 Parsippany, NJ 07054

MADE IN INDIA

8 13874 02053 7

Lot
 Exp.

COATING FREE AREA

Principal Display Panel

size :1+3/4 X 1+3/4 X 3+3/8
 ref # :PP180604B
 material :.016 SBS

COATING
FREE AREA

200000003205
700000002397

Drug Facts (continued)

Other Information

- each tablet contains: sodium 20 mg
- store at 20 - 25°C (68 - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive Ingredients FDCC Blue #2 aluminum lake, hypromellose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, povidone K-30, sodium starch glycolate, stearic acid, titanium dioxide.

Questions or comments?
 1-877-276-3183 Mon-Fri 9:00 AM to 4:00 PM EST.

This product is not recommended or endorsed by Bayer Healthcare, U.S.A., distributor of Aleve. Aleve is a registered trademark of Bayer Healthcare, U.S.A.

KEEP CARTON FOR REFERENCE

DO NOT USE IF CARTON IS OPEN OR IF FOIL SEAL ON BOTTLE OPENING IS MISSING OR BROKEN

Drug Facts (continued)

naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have had high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools

Drug Facts (continued)

- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12-hour period
- do not exceed 3 tablets in a 24-hour period

Children under 12 years

- ask a doctor

100 Tablets

Contains no preservatives from a gluten-containing grain (wheat, barley, or rye)

STRENGTH TO LAST 12 HOURS

Granules Consumer Health

Naproxen Sodium Tablets USP

220 mg (NSAID) **

Pain reliever/fever reducer

Compare to the Active Ingredient of Aleve® Tablets



Drug Facts

Active ingredient (in each tablet)	Purposes
Naproxen sodium 220 mg. Pain reliever/fever reducer (naproxen 200 mg) (NSAID)**	muscle aches menstrual cramps toothache

**nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscle aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen,

COATING
FREE AREA

200000003205
700000002397

Distributed by:
 Granules Consumer Health
 (A division of Granules USA, Inc.)
 35 Waterway Blvd., 3rd Floor
 Parsippany, NJ 07054
 MADE IN INDIA

8 13874 02054 4

Lot
Exp.

COATING
FREE AREA

Principal Display Panel

size :1+3/4 X 1+3/4 X 3+5/8
 ref # :PP180604B
 material :.016 SBS



NAPROXEN SODIUM

naproxen sodium tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69848-010
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:5Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	220
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69848-010-01	10 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2019	
2	NDC:69848-010-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2023	
3	NDC:69848-010-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091353	07/01/2019	

NAPROXEN SODIUM

naproxen sodium tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69848-009
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	220
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69848-009-01	10 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091353	07/01/2019	12/31/2023

Labeler - Granules USA, Inc. (137098864)

Revised: 12/2025

Granules USA, Inc.