

IBUPROFEN ORAL- ibuprofen oral suspension
Preferred Pharmaceuticals Inc.

Ibuprofen Oral Suspension USP, 100 mg/5 mL (OTC)

ACTIVE INGREDIENT(S)

(in each 5 mL)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

PURPOSE

Pain reliever/fever reducer

USE(S)

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

WARNINGS

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:☐

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed
- **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

DO NOT USE

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

ASK A DOCTOR BEFORE USE IF

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

WHEN USING THIS PRODUCT

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

DIRECTIONS

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information

□

- **each 5 mL contains :** sodium 2 mg

- store between 20-25°C (68-77°F)
- **do not use if carton is opened or printed bottle neckband is broken or missing**
- see bottom panel for lot number and expiration date

INACTIVE INGREDIENT SECTION

Berry flavor (with dye)

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Yellow #10, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

Berry flavor (dye free)

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

Grape flavor

-

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Red #33, FD&C Blue #1, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

Bubble Gum flavor

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Red #33, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

-

-

QUESTIONS SECTION

Call 1-888-588-1418 from 9 AM to 5 PM EST,
Monday-Friday.



Distributed by:
Camber Consumer Care, Inc.

Piscataway, NJ 08854, USA

Relabeled By: Preferred Pharmaceuticals Inc.

PRINCIPAL DISPLAY PANEL

Ibuprofen oral suspension berry flavor with dye container carton

**Childrens
Ibuprofen Oral
Suspension**

Generic for Childrens Motrin

Active ingredient (in each 5mL): Ibuprofen, USP 100mg (NSAID)...pain reliever/fever reducer

Pkg Size: Exp Date: #####/###/####

Ins:
Mfg: Camber Consumer Care
Prod#:
Warning

Do not use if printed neckband is broken or missing. Store between 20°- 25° C (68°- 77°F). Sore throat warning: severe or persistent sore throat or sore throat accompanied by a high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. Read box for additional warnings, drug facts, directions, and other information. Original Berry Flavor. Alcohol free.



CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed.



Directions English
Take ___ teaspoonful(s)
) every ___ hours.



GTIN

SN #####
EXP #####/###/####

Instrucciones Espanol:
Toma ___ cucharadita(s)
) cada ___ horas

Childrens Ibuprofen Oral Suspension
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Childrens Ibuprofen Oral Suspension
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Childrens Ibuprofen Oral Suspension
Qty:
Insurance NDC:
Lot: Bat:

Childrens Ibuprofen Oral Suspension
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Log

Chart

Billing

Patient

IBUPROFEN ORAL			
ibuprofen oral suspension			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68788-7984(NDC:69230-308)
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL
Inactive Ingredients			
	Ingredient Name		Strength
	ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)		
	ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
	CARBOXYMETHYLCELLULOSE SODIUM (1.3 CARBOXYMETHYL SUBSTITUTION PER SACCHARIDE; 600 MPA.S AT 2%) (UNII: 72QQR5RYU4)		
	D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
	GLYCERIN (UNII: PDC6A3C0OX)		
	FD&C RED NO. 40 (UNII: WZB9127XOA)		
	MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
	POLYSORBATE 80 (UNII: 6OZP39ZG8H)		
	PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		

WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	ORANGE	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-7984-1	1 in 1 CARTON	08/05/2021	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210602	08/05/2021	

Labeler - Preferred Pharmaceuticals Inc. (791119022)

Registrant - Preferred Pharmaceuticals Inc. (791119022)

Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals Inc.		791119022	RELABEL(68788-7984)

Revised: 10/2025

Preferred Pharmaceuticals Inc.