GOOD NEIGHBOR PHARMACY INFANTS PAIN AND FEVERacetaminophen suspension Amerisource Bergen

Amerisource Bergen Infant's Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- · skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Insert syringe tip into bottle opening.
- flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.
- pull yellow part of syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
 replace cap tightly to maintain child resistance

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

* or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other information

- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Infants' Tylenol® active ingredient

FOR AGES 2 TO 3 YEARS

infants' Pain & Fever

acetaminophen 160 mg per 5 mL suspension liquid

Fever Reducer - Pain Reliever

Alcohol Free

Aspirin Free

Ibuprofen Free

Use only with enclosed syringe.

See side panel for more information.

Grape Flavor

2 fl oz (60 mL)

160 mg per 5 mL



Pain & Fever





Infants' Tylenol

FOR AGES infants'

acetaminophen 160 mg per 5 mL suspension liquid

Fever Reducer – Pain Reliever

Alcohol Free Aspirin Free Ibuprofen Free

Use only with enclosed syringe. See side panel for more information.

Grape Flavor

2 fl oz (60 mL) 160 mg per 5 mL

IMPORTANT:

Please see dosing chart or ask a doctor for your child's correct dose.

- STEP 1. See *Directions* for the correct dose for your child.
- STEP 2. Shake well before using.

STEP 3. Push air out of syringe. Insert syringe tip into bottle opening



- STEP 4. Flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle
- STEP 5. Pull yellow part of syringe until it reaches and stays at the correct dose.



STEP 6. Dispense liquid slowly into

Drug Facts

Active ingredient (in each 5 mL) Purpose

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to:

- flu headache sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe

- liver damage may occur if your child takes

 more than 5 doses in 24 hours, which is the maximum daily amount

 with other drugs containing acetaminophen
- With other order containing acceleration principle.

 Allergy alert: Acetaminophen may cause severe skin reactions.

 Symptoms may include: skin reddening blisters rash

 If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription), If you are not sure whether a

drug contains acetaminophen, ask a doctor or pharmacist.
■ if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver disease Ask a doctor or pharmacist before use if your child is taking

the blood thinning drug warfarin When using this product do not exceed recommended dose

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days fever gets worse or lasts more than 3 days
- new symptoms occur redness or swelling is present.

 These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms

Directions

- this product does not contain directions or complete warnings for adult use
- wainings for adult use

 do not give more than directed (see overdose warning)

 shake well before using mL = milliliter

 find right dose on chart. If possible, use weight to dose;
- otherwise, use age.

 push air out of syringe. Insert syringe tip into bottle opening.

 flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.

Drug Facts (continued)

- pull yellow part of syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child's
- mouth, toward inner cheek
 repeat dose every 4 hours while
- symptoms last ■ do not give more than 5 times in
- replace cap tightly to maintain
- child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*	
under 24	under 2 years	ask a doctor	
24-35	2-3 years	5 mL	

or as directed by a doctor Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing

Other information

- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing


Inactive ingredients anhydrous citric acid, butylparaben, calcium sulfat carrageenan, D&C red #33, FD&C blue #1 flavor, glycerin, high fructose com syrup hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments? 1-800-719-9260

**Good Neighbor Pharmacy® infants' Pain & Fever is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol®.

Distributed By AmerisourceBergen 1300 Morris Drive, Chesterbrook, PA 19087 Questions or Concerns? www.mygnp.co

Strenath





94616 29 C2

GOOD NEIGHBOR PHARMACY INFANTS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:46122-552

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
BUTYLPARABEN (UNII: 3QPI1U3FV8)		
CARRAGEENAN (UNII: 5C69YCD2YJ)		
D&C RED NO. 33 (UNII: 9DBA0SBB0L)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
GLYCERIN (UNII: PDC6A3C0OX)		
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SORBITOL (UNII: 506T60A25R)		
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)		

Product Characteristics			
Color	PURPLE	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-552- 46	1 in 1 CARTON	07/30/2018	08/31/2025
1		60 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/30/2018	08/31/2025

Labeler - Amerisource Bergen (007914906)

Revised: 11/2024 Amerisource Bergen