


**BENZONATATE- benzonatate capsule**  
**REMEDYREPACK INC.**

-----  
**Benzonatate Capsules USP, 100 mg and 200 mg**

**DESCRIPTION**

Benzonatate, a non-narcotic oral antitussive agent, is 2, 5, 8, 11, 14, 17, 20, 23, 26-nonaooctacosan-28-yl p-(butylamino) benzoate; with a molecular weight of 603.7.



Each Benzonatate Capsules USP contains: Benzonatate, USP 100 mg or 200 mg.

Benzonatate Capsules USP also contain: D&C Yellow 10, gelatin, glycerin, methylparaben, propylparaben and titanium dioxide.

**CLINICAL PHARMACOLOGY**

Benzonatate Capsule acts peripherally by anesthetizing the stretch receptors located in the respiratory passages, lungs, and pleura by dampening their activity and thereby reducing the cough reflex at its source. It begins to act within 15 to 20 minutes and its effect lasts for 3 to 8 hours. Benzonatate Capsule has no inhibitory effect on the respiratory center in recommended dosage.

**INDICATIONS AND USAGE**

Benzonatate Capsule is indicated for the symptomatic relief of cough.

**CONTRAINDICATIONS**

Hypersensitivity to benzonatate or related compounds.

**WARNINGS**

**Hypersensitivity**

Severe hypersensitivity reactions (including bronchospasm, laryngospasm and cardiovascular collapse) have been reported which are possibly related to local anesthesia from sucking or chewing the capsule instead of swallowing it. Severe reactions have required intervention with vasopressor agents and supportive measures.

### **Psychiatric Effects**

Isolated instances of bizarre behavior, including mental confusion and visual hallucinations, have also been reported in patients taking Benzonatate Capsule in combination with other prescribed drugs.

### **Accidental Ingestion and Death in Children**

Keep Benzonatate Capsules out of reach of children. Accidental ingestion of Benzonatate Capsules resulting in death has been reported in children below age 10. Signs and symptoms of overdose have been reported within 15-20 minutes and death has been reported within one hour of ingestion. If accidental ingestion occurs, seek medical attention immediately (see OVERDOSAGE).

### **PRECAUTIONS**

Benzonatate is chemically related to anesthetic agents of the para-amino-benzoic acid class (e.g. procaine; tetracaine) and has been associated with adverse CNS effects possibly related to a prior sensitivity to related agents or interaction with concomitant medication.

### **Information for Patients**

Swallow Benzonatate Capsules whole. Do not break, chew, dissolve, cut, or crush Benzonatate Capsules. Release of Benzonatate from the capsule in the mouth can produce a temporary local anesthesia of the oral mucosa and choking could occur. If numbness or tingling of the tongue, mouth, throat, or face occurs, refrain from oral ingestion of food or liquids until the numbness has resolved. If the symptoms worsen or persist, seek medical attention.

Keep Benzonatate Capsules out of reach of children. Accidental ingestion resulting in death has been reported in children. Signs and symptoms of overdose have been reported within 15-20 minutes and death has been reported within one hour of ingestion. Signs and symptoms may include restlessness, tremors, convulsions, coma and cardiac arrest. If accidental ingestion occurs, seek medical attention immediately.

Overdosage resulting in death may occur in adults.

Do not exceed a single dose of 200 mg and a total daily dosage of 600 mg. If you miss a dose of Benzonatate Capsules, skip that dose and take the next dose at the next scheduled time. Do not take 2 doses of Benzonatate Capsules at one time.

Repackaged By / Distributed By: RemedyRepack Inc.

625 Kolter Drive, Indiana, PA 15701

(724) 465-8762

Usage in Pregnancy

Pregnancy Category C.

Animal reproduction studies have not been conducted with Benzonatate Capsules. It is also not known whether Benzonatate Capsules can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Benzonatate Capsules should be given to a pregnant woman only if clearly needed.

### **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk caution should be exercised when Benzonatate Capsule is administered to a nursing woman.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Carcinogenicity, mutagenicity, and reproduction studies have not been conducted with Benzonatate Capsules.

### **Pediatric Use**

Safety and effectiveness in children below the age of 10 have not been established. Accidental ingestion resulting in death has been reported in children below age 10. Keep out of reach of children.

## **ADVERSE REACTIONS**

Potential Adverse Reactions to Benzonatate Capsules may include:

**Hypersensitivity reactions** including bronchospasm, laryngospasm, cardiovascular collapse possibly related to local anesthesia from chewing or sucking the capsule.

**CNS:** sedation; headache; dizziness; mental confusion; visual hallucinations.

**GI:** constipation; nausea; GI upset.

**Dermatologic:** pruritus; skin eruptions.

**Other:** nasal congestion; sensation of burning in the eyes; vague "chilly" sensation; numbness of the chest; hypersensitivity. Deliberate or accidental overdose has resulted in death, particularly in children.

## **OVERDOSAGE**

Intentional and unintentional overdose may result in death, particularly in children.

The drug is chemically related to tetracaine and other topical anesthetics and shares various aspects of their pharmacology and toxicology. Drugs of this type are generally well absorbed after ingestion.

### **Signs and Symptoms:**

The signs and symptoms of overdose of benzonatate have been reported within 15-20

minutes. If capsules are chewed or dissolved in the mouth, oropharyngeal anesthesia will develop rapidly, which may cause choking and airway compromise.

CNS stimulation may cause restlessness and tremors which may proceed to clonic convulsions followed by profound CNS depression. Convulsions, coma, cerebral edema and cardiac arrest leading to death have been reported within 1 hour of ingestion.

### **Treatment:**

In case of overdose, seek medical attention immediately. Evacuate gastric contents and administer copious amounts of activated charcoal slurry. Even in the conscious patient, cough and gag reflexes may be so depressed as to necessitate special attention to protection against aspiration of gastric contents and orally administered materials. Convulsions should be treated with a short-acting barbiturate given intravenously and carefully titrated for the smallest effective dosage. Intensive support of respiration and cardiovascular-renal function is an essential feature of the treatment of severe intoxication from overdosage. Do not use CNS stimulants.

## **DOSAGE AND ADMINISTRATION**

Adults and Children over 10 years of age: Usual dose is one 100 mg or 200 mg capsule three times a day as needed for cough. If necessary to control cough, up to 600 mg daily in three divided doses may be given. **Benzonatate Capsules should be swallowed whole.** Benzonatate Capsules are not to be broken, chewed, dissolved, cut or crushed.

## **HOW SUPPLIED**

Benzonatate Capsules USP , 200 mg are available as yellow, oval soft gelatin capsules with '2' imprinted in white ink.

NDC: 70518-3542-00

NDC: 70518-3542-01

NDC: 70518-3542-02

NDC: 70518-3542-03

NDC: 70518-3542-04

PACKAGING: 30 in 1 BLISTER PACK

PACKAGING: 21 in 1 BOTTLE PLASTIC

PACKAGING: 28 in 1 BOTTLE PLASTIC

PACKAGING: 15 in 1 BOTTLE PLASTIC

PACKAGING: 30 in 1 BOTTLE PLASTIC

Store at 20° to 25°C (68° to 77°F) excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

PROTECT FROM LIGHT

Repackaged and Distributed By:

Remedy Repack, Inc.

625 Kolter Dr. Suite #4 Indiana, PA 1-724-465-8762

Repackaged and Distributed By:

Remedy Repack, Inc.

625 Kolter Dr. Suite #4 Indiana, PA 1-724-465-8762

## **PRINCIPAL DISPLAY PANEL**

DRUG: Benzonatate

GENERIC: Benzonatate

DOSAGE: CAPSULE

ADMINISTRATION: ORAL

NDC: 70518-3542-0

NDC: 70518-3542-1

NDC: 70518-3542-2

NDC: 70518-3542-3

NDC: 70518-3542-4

COLOR: yellow

SHAPE: OVAL

SCORE: No score

SIZE: 11 mm

IMPRINT: 2

PACKAGING: 30 in 1 BLISTER PACK

PACKAGING: 21 in 1 BOTTLE PLASTIC

PACKAGING: 28 in 1 BOTTLE PLASTIC

PACKAGING: 15 in 1 BOTTLE PLASTIC

PACKAGING: 30 in 1 BOTTLE PLASTIC

ACTIVE INGREDIENT(S):

- BENZONATATE 200mg in 1

INACTIVE INGREDIENT(S):

- D&C YELLOW NO. 10
- GELATIN, UNSPECIFIED
- GLYCERIN
- METHYLPARABEN
- PROPYLPARABEN

- TITANIUM DIOXIDE

## Benzonatate Capsule

NDC #: 70518-3542-00

Expires:

Protect from light.

LOT #:

Org NDC: 51224-0001-60

MFG: TAGI Pharma, South Beloit, IL 61080

200 mg

QTY: 30 Capsules

WARNING: Do not break, chew, dissolve, cut, or crush Benzonatate capsules

Oval YELLOW 2

Usual Dosage: See Insert



RX ONLY

Repackaged By:  
RemedyRepack Inc.,  
Indiana, PA 15701, 724.465.8762

Keep this and all medication out of the reach of children  
Store at 20-25°C (68-77°F);  
excursions permitted to 15-30°C (59-86°F) [See USP]

LOT: 180101  
EXP: 12/2020  
NDC: 70518-3542-00  
NDC: 51224-0001-60  
MFG: TAGI PHARMA  
SOUTHBELoit, IL 61080  
REPACKED BY:  
REMEDY REPACK  
INDIANA, PA 15701  
724.465.8762

## Benzonatate Capsule

NDC #: 70518-3542-04

Expires:

Protect from light.

LOT #:

Org NDC: 51224-0001-60

MFG: TAGI Pharma, South Beloit, IL 61080

200 mg

QTY: 30 Capsules

WARNING: Swallow benzonatate capsules USP whole. Do not break, chew, dissolve, cut, or crush benzonatate capsules USP.

Oval YELLOW 2

Usual Dosage: See Insert



RX ONLY

Repackaged By:  
RemedyRepack Inc.,  
Indiana, PA 15701, 724.465.8762

Keep this and all medication out of the reach of children  
Store at 20-25°C (68-77°F);  
excursions permitted to 15-30°C (59-86°F) [See USP]

LOT: 180101  
EXP: 12/2020  
NDC: 70518-3542-00  
NDC: 51224-0001-60  
MFG: TAGI PHARMA  
SOUTHBELoit, IL 61080  
REPACKED BY:  
REMEDY REPACK  
INDIANA, PA 15701  
724.465.8762

## BENZONATATE

benzonatate capsule

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70518-3542(NDC:51224-001)
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZONATATE (UNII: 5P4DHS6ENR) (BENZONATATE - UNII:5P4DHS6ENR)	BENZONATATE	200 mg

Inactive Ingredients				
Ingredient Name			Strength	
GLYCERIN (UNII: PDC6A3C0OX)				
METHYLPARABEN (UNII: A2I8C7HI9T)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)				
Product Characteristics				
Color	yellow	Score	no score	
Shape	OVAL	Size	11mm	
Flavor		Imprint Code	2	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70518-3542-0	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	09/29/2022	
2	NDC:70518-3542-1	21 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/03/2023	03/21/2025
3	NDC:70518-3542-2	28 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/15/2023	04/29/2025
4	NDC:70518-3542-3	15 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/16/2023	04/29/2025
5	NDC:70518-3542-4	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/05/2023	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA202765	09/29/2022	

**Labeler** - REMEDYREPACK INC. (829572556)

Revised: 7/2025

REMEDYREPACK INC.