

MUCINEX INSTASOOTHE SORE THROAT PLUS COUGH RELIEF-
dextromethorphan hydrobromide and hexylresorcinol lozenge
RB Health (US) LLC

MUCINEX® INSTASOOTHE™
SORE THROAT PLUS COUGH RELIEF

Drug Facts

Active ingredients (in each lozenge)

Dextromethorphan HBr 5 mg

Hexylresorcinol 2 mg

<i>Active ingredients (in each lozenge)</i>	<i>Purposes</i>
Dextromethorphan HBr 5 mg	Cough suppressant
Hexylresorcinol 2 mg	Oral pain reliever

Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold
- occasional irritation and pain due to a sore mouth or sore throat
- pain associated with canker sores

Warnings

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in children under 6 years of age

When using this product do not exceed recommended dosage

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor or dentist if

- sore mouth symptoms do not improve in 7 days

- irritation, pain, or redness persists or worsens
- swelling develops
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

dose as follows or as directed by a doctor or dentist

adults and children 12 years of age and older	take 2 lozenges (one immediately after the other) and allow each lozenge to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 12 lozenges in any 24-hour period
children 6 to under 12 years of age	take 1 lozenge and allow product to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 6 lozenges in any 24-hour period
children under 6 years of age	do not use

Other information

- store at 20-25°C (68-77°F)
- protect contents from humidity
- keep lozenge in outer pack until use. Reseal outer pack after use.

Inactive ingredients

corn syrup, D&C yellow no. 10, ethyl lactate, FD&C blue no. 1, flavor, propylene glycol, purified water, sucrose.

Questions?

1-866-MUCINEX (1-866-682-4639)

**Dist. by: RB Health (US)
Parsippany,
NJ 07054-0224
Made in Thailand**

PRINCIPAL DISPLAY PANEL - 20 Drop Pouch Bag

NDC 72854-263-20

Mucinex®
INSTASOOTHE™

SORE THROAT
+ COUGH RELIEF

Dextromethorphan HBr 5.0 mg | Cough Suppressant
Hexylresorcinol 2.0 mg | Oral Pain Reliever

3 IN 1 FORMULA

- ✓ Up to 4 Hours Cough Relief
- ✓ Cooling Comfort
- ✓ Fast Acting

See back
panel for
actual size

Winter Mint Flavor

20 MEDICATED DROPS

NDC 72854-263-20

Mucinex[®]

INSTA»SOOTHE[®]

SORE THROAT + COUGH

Dextromethorphan HBr 5 mg » Cough Suppressant
Hexylresorcinol 2 mg » Oral Pain Reliever



20 MEDICATED
DROPS



FRESH
MINT
FLAVOR

Mucinex®

INSTA-SOOTHE®

Choose from our range of Mucinex® INSTASOOTHE® products for powerful symptom relief!



SORE THROAT
HONEY & ECHINACEA FLAVOR

COUGH RELIEF
MIXED BERRY FLAVOR

SORE THROAT
ELDERBERRY & WILD CHERRY FLAVOR

Drug Facts

Active ingredients (in each lozenge) **Purposes**
Dextromethorphan HBr 5 mg.....Cough suppressant
Hexylresorcinol 2 mg.....Oral pain reliever

Uses temporarily relieves:

- cough due to minor throat and bronchial irritation as may occur with the common cold
- occasional irritation and pain due to a sore mouth or sore throat
- pain associated with canker sores

Warnings
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in children under 6 years of age

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product do not exceed recommended dosage

Stop use and ask a doctor or dentist if

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling develops

*Helps relieve these symptoms day or night

Drug Facts (continued)

■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ dose as follows or as directed by a doctor or dentist

adults and children 12 years of age and older	take 2 lozenges (one immediately after the other) and allow each lozenge to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 12 lozenges in any 24-hour period
children 6 to under 12 years of age	take 1 lozenge and allow product to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 6 lozenges in any 24-hour period
children under 6 years of age	do not use

Other information

- store at 20-25°C (68-77°F)
- protect contents from humidity
- keep lozenge in outer pack until use. Reseal outer pack after use.

Inactive ingredients
 corn syrup, D&C yellow no. 10, ethyl lactate, FD&C blue no. 1, flavor, propylene glycol, purified water, sucrose

Questions? 1-866-MUCINEX (1-866-682-4639)



Dist. by: RB Health (US)
 Parsippany, NJ 07054-0224
 Made in Thailand
 ©2025 RB Health
 Patents: www.reckitt.com/patents
www.mucinex.com
 021225

Keep outer pack for full information



ACTUAL SIZE

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



3 63824 21572 3



50063844

MUCINEX

MUCINEX INSTASOOthe SORE THROAT PLUS COUGH RELIEF

dextromethorphan hydrobromide and hexylresorcinol lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72854-263
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg
HEXYLRESORCINOL (UNII: R9QTB5E82N) (HEXYLRESORCINOL - UNII:R9QTB5E82N)	HEXYLRESORCINOL	2 mg

Inactive Ingredients

Ingredient Name	Strength
CORN SYRUP (UNII: 9G5L16BK6N)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ETHYL LACTATE (UNII: F3P750VW8I)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SUCROSE (UNII: C151H8M554)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	turquoise	Score	no score
Shape	ROUND	Size	18mm
Flavor	MINT	Imprint Code	S
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-263-20	20 in 1 BAG	07/01/2021	
1		1 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:72854-263-40	40 in 1 BAG	07/01/2021	
2		1 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
------------------	--	------------------------	----------------------

Category	Citation	Date	Date
OTC Monograph Drug	M012	07/01/2021	

Labeler - RB Health (US) LLC (081049410)

Revised: 4/2025

RB Health (US) LLC