

**DAYTIME NIGHTTIME COLD FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl**  
**FSA Store Inc. (Caring Mills)**

-----

**DRUG FACTS**

**Active ingredients for Daytime (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Active ingredients for Nighttime (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose for Daytime**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Purpose for Nighttime**

**Pain reliever/fever reducer**

Cough suppressant

Antihistamine

**Uses**

**DAYTIME**

- temporarily relieves common cold and flu symptoms
  - cough due to minor throat and bronchial irritation
  - fever
  - sore throat
  - nasal congestion and sinus pressure
  - headache
  - minor ache and pains

**NIGHTTIME**

- temporarily relieves common cold and flu symptoms
  - cough due to minor throat and bronchial irritation
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing

## Warnings

### DAYTIME

**Liver warning:** These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using these products

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### NIGHTTIME

**Liver warning:** This product contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or following by fever, headache, rash, nausea, vomiting, consult a doctor promptly.

## Do not use

### DAYTIME

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **NIGHTTIME**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

#### **DAYTIME**

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

#### **NIGHTTIME**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

#### **DAYTIME**

taking the blood thinning drug warfarin

#### **NIGHTTIME**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product,**

#### **DAYTIME**

**do not exceed recommended dosage.**

## **NIGHTTIME**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

## **Stop use and ask a doctor if**

### **DAYTIME**

- nervousness, dizziness or sleeplessness occur
- pain, cough, and nasal congestion gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

### **NIGHTTIME**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash headache that lasts.

These could be a signs of a serious condition.

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

### **DAYTIME**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over; take 2 softgels with water every 4 hours.
- children under 12 years: do not use

### **NIGHTTIME**

- do not take more than directed (see Overdose warning)

- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

### **Other information**

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

### **Inactive ingredients**

#### **DAYTIME**

FD&C red #40, FD&C yellow #6, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

#### **NIGHTTIME**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

### **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

\*\*Compare to Vicks® Dayquil™ and Nyquil™ Cold & Flu LiquiCaps™ active ingredients

#### **DAYTIME**

#### **Cold & Flu Relief**

Pain reliever, Fever Reducer,

Cough Suppressant, Nasal Decongestant

#### **Acetaminophen**

Dextromethorphan HBr

Phenylephrine HCl

Alcohol-free

Antihistamine-free

Softgels\*

(\*Liquid-filled capsules)

#### **NIGHTTIME**

## **Cold & Flu Relief**

Pain Reliever, Fever Reducer,  
Cough Suppressant, Antihistamine

## **Acetaminophen**

Dextromethorphan HBr  
Doxylamine Succinate  
Alcohol-free

**When using Daytime and Nighttime product, carefully read the labeling to ensure correct dosing.**

Softgels

(\*Liquid-filled capsules)

\*\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil™, NyQuil™, and LiquiCaps™ are registered trademarks of the Procter and Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Manufactured for FSA Stores Inc.

5473 Blair Rd

Suite 100

PMB 24308

Dallas, Texas 75231

**Product Label**

P.D.-5086C  
FD069833  
Lot No.:  
Exp. Date:

# caring mill™ Nighttime Cold & Flu Relief



NDC 101-06-15122-46

\*\*Compare to Vicks® DayQuil™ and NyQuil™ active ingredients

**Daytime  
Cold & Flu Relief**  
Pain Reliever, Fever Reducer,  
Cough Suppressant, Nasal Decongestant  
**Acetaminophen**  
Dextromethorphan HBr  
Phenylephrine HCl



Actual Size

Alcohol-free  
Antihistamine-free

**32 Softgels\***  
(\*Liquid-filled capsules)

**Nighttime  
Cold & Flu Relief**  
Pain Reliever, Fever Reducer,  
Cough Suppressant, Antihistamine  
**Acetaminophen**  
Dextromethorphan HBr  
Doxylamine succinate



Actual Size

Alcohol-free  
When using Daytime and Nighttime  
products, carefully read the labeling  
to ensure correct dosing.

**48 Total Softgels\***  
(\*Liquid-filled capsules)

**16 Softgels\***  
(\*Liquid-filled capsules)



3 59726 90148 2

PMB 24308  
Dallas, Texas 75231

## Daytime Cold & Flu Softgel

**Drug Facts**  
**Purposes** Pain reliever/fever reducer  
Acetaminophen 325 mg  
Cough suppressant  
Dextromethorphan HBr 10 mg  
Nasal decongestant  
Phenylephrine HCl 5 mg

**Uses** Temporarily relieves common cold and flu symptoms:  
cough due to minor throat and bronchial irritation ■ fever ■ sore throat  
■ nasal congestion and sinus pressure ■ headache ■ minor aches and pains

**Warnings** Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours  
■ 3 or more alcohol drinks every day while using the product  
■ with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Do not use** ■ if you are now taking a prescription medication called tricyclic (MAD) (certain drugs for depression, psychiatric, or medical conditions, or Parkinson's disease), or for 2 weeks after stopping the MAD drug, if you do not know if your prescription drug contains an MAD, ask a doctor or pharmacist before taking the product.

**Do not use** ■ if you are now taking a prescription medication called tricyclic (MAD) (certain drugs for depression, psychiatric, or medical conditions, or Parkinson's disease), or for 2 weeks after stopping the MAD drug, if you do not know if your prescription drug contains an MAD, ask a doctor or pharmacist before taking the product.

**Do not use** ■ if you are now taking a prescription medication called tricyclic (MAD) (certain drugs for depression, psychiatric, or medical conditions, or Parkinson's disease), or for 2 weeks after stopping the MAD drug, if you do not know if your prescription drug contains an MAD, ask a doctor or pharmacist before taking the product.

**Drug Facts (continued)**  
**Ask a doctor before use if you have** ■ liver disease ■ heart disease ■ diabetes ■ thyroid disease ■ high blood pressure ■ asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if** ■ nervousness, dizziness, or sleepiness occur ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ redness or swelling is present ■ new symptoms occur ■ fever gets worse or lasts more than 3 days

**Stop use and ask a doctor if** ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.** Overdose warnings: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ do not take more than 4 doses in 24 hours  
■ swallow whole, do not crush, chew, or dissolve  
■ adults and children 17 years and over: take 2 softgels with water every 4 hours  
■ children under 17 years: do not use

## Nighttime Cold & Flu Softgel

**Drug Facts**  
**Purposes** Pain reliever/fever reducer  
Acetaminophen 325 mg  
Cough suppressant  
Dextromethorphan HBr 10 mg  
Antihistamine  
Doxylamine succinate 5.25 mg

**Uses** Temporarily relieves common cold and flu symptoms:  
cough due to minor throat and bronchial irritation ■ sore throat  
■ headache ■ minor aches and pains ■ fever ■ runny nose & sneezing

**Warnings** Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours  
■ 3 or more alcohol drinks every day while using the product  
■ with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Do not use** ■ if you are now taking a prescription medication called tricyclic (MAD) (certain drugs for depression, psychiatric, or medical conditions, or Parkinson's disease), or for 2 weeks after stopping the MAD drug, if you do not know if your prescription drug contains an MAD, ask a doctor or pharmacist before taking the product.

**Do not use** ■ if you are now taking a prescription medication called tricyclic (MAD) (certain drugs for depression, psychiatric, or medical conditions, or Parkinson's disease), or for 2 weeks after stopping the MAD drug, if you do not know if your prescription drug contains an MAD, ask a doctor or pharmacist before taking the product.

**Do not use** ■ if you are now taking a prescription medication called tricyclic (MAD) (certain drugs for depression, psychiatric, or medical conditions, or Parkinson's disease), or for 2 weeks after stopping the MAD drug, if you do not know if your prescription drug contains an MAD, ask a doctor or pharmacist before taking the product.

**Drug Facts (continued)**  
**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers


**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if** ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.** Overdose warnings: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ do not take more than 4 doses in 24 hours  
■ swallow whole, do not crush, chew, or dissolve  
■ adults and children 17 years and over: take 2 softgels with water every 5 hours  
■ children under 17 years: do not use



## CARING MILL Daytime Nighttime Cold & Flu Flu

### DAYTIME NIGHTTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:81522-901
---------------------	----------------	---------------------------	---------------

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81522-901-48	1 in 1 KIT; Type 0: Not a Combination Product	01/31/2023	

#### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	16 BLISTER PACK	16
Part 2	32 BLISTER PACK	32

### Part 1 of 2

### NIGHT TIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

#### Product Information

<b>Item Code (Source)</b>	NDC:81522-899
<b>Route of Administration</b>	ORAL

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	

**Product Characteristics**

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	PC10
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1		16 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012	01/31/2023	

# DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

## Product Information

**Item Code (Source)** NDC:81522-898

**Route of Administration** ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

## Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	PC9
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		32 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/31/2023	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/31/2023	

**Labeler** - FSA Store Inc. (Caring Mills) (049283340)

Revised: 9/2025

FSA Store Inc. (Caring Mills)