

ONLY FOR FEVER- acetaminophen tablet
O4 Global Trading Usa, Llc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Only For- FEVER

DRUG FACTS

Active ingredient (in each caplet)

Acetaminophen 500mg

Purpose

Pain Reliever - Fever reducer

Uses:

- temporarily relieves minor aches and pains due to:
- the common cold • headache • backache • minor pain of arthritis • toothache • muscular aches • premenstrual and menstrual cramps • temporarily reduces fever

Warnings:

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 3,000mg of acetaminophen in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have
liver disease

Ask a doctor or a pharmacist before use if
you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse

If pregnant or breast-feeding,
ask a health professional before use.

Keep out of reach of children

Overdose Warning: In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions:

- take with a full glass of water
- **do not take more than directed (See Overdose Warning)**

adults and children 12 years of age & over	<ul style="list-style-type: none">• take 2 caplets every 6 hours while symptoms persist.• do not take more than 6 caplets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children under 12 years of age	ask a doctor

Other Information:

- Store at room temperature between 20-25°C (68-77°F).

Inactive Ingredients:

colloidal silicon dioxide, gelatin, glycerin, hydroxypropyl methyl cellulose, maize starch, magnesium stearate, purified talc, sodium starch glycolate.

Questions Or Comments ?

1-833-665-9444.

NDC: 73006-117-41

ADULT

PAIN RELIEVER-FEVER REDUCER

Distributed By: O4 GLOBAL TRADING USA LLC., 2241 S Watson RD, Suite 121 Arlington, TX 76010.

www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc., East Brunswick, NJ 08816.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF A BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGN OF TAMPERING.

Tear along perforation push product through foil

Packaging



NDC: 73006-117-41

ADULT

Only For

FEVER
PAIN RELIEVER- FEVER REDUCER

ACETAMINOPHEN 500mg / Pain Reliever - Fever reducer



4 CAPLETS



LOT: K009A

EXP: 06/21

DRUG FACTS

Active ingredient (in each caplet)	Purpose
Acetaminophen 500mg	Pain Reliever - Fever reducer

Uses:

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
 - temporarily reduces fever

Warnings:

Liver warning:
This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 3,000mg of acetaminophen in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:
Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have
liver disease

Ask a doctor or pharmacist before use if
you are taking the blood thinning drug warfarin

DRUG FACTS(Continued)

Stop use and ask a doctor if

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse

If pregnant or breast-feeding,
ask a health professional before use.

Keep out of reach of children

Overdose Warning: In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions:

- take with a full glass of water
- do not take more than directed (See Overdose Warning)

adults and children 12 years of age & over	<ul style="list-style-type: none"> take 2 caplets every 6 hours while symptoms persist. do not take more than 6 caplets in 24 hours, unless directed by a doctor do not use for more than 10 days unless directed by a doctor
children under 12 years of age	ask a doctor

Other Information:

- Store at room temperature between 20-25°C (68-77°F).

Inactive Ingredients:

colloidal silicon dioxide, gelatin, glycerin, hydroxypropyl methyl cellulose, maize starch, magnesium stearate, purified talc, sodium starch glycolate.

Questions Or Comments ?

1-833-665-9444.

Distributed By: O4 GLOBAL TRADING USA LLC., 2241 S Watson RD, Suite 121 Arlington, TX 76010.

www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc., East Brunswick, NJ 08816.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF A BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGN OF TAMPERING.



www.onlyfor.com

CT2FBR4-REV.102018

ONLY FOR FEVER

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73006-117
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)	
STARCH, CORN (UNII: O8232NY3SJ)	
MAGNESIUM PALMITOSTEARATE (UNII: R4OXA9G5BV)	
TALC (UNII: 7SEV7J4R1U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL (CAPLET)	Size	18mm
Flavor		Imprint Code	A5
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73006-117-41	1 in 1 CARTON	03/29/2019	
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	03/29/2019	

Labeler - O4 Global Trading Usa, Llc (081226861)