

STOMACH RELIEF, ULTRA STRENGTH- bismuth subsalicylate suspension
Marc Glassman, Inc.

Stomach Relief, Ultra Strength

Drug Facts

Active ingredient

(in each 30 mL dose cup or 2 tablespoons)

Bismuth subsalicylate 1050 mg

Purposes

Bismuth subsalicylate Upset stomach reliever
and antidiarrheal

Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach
due to overindulgence of food and drink including: ■ heart burn ■
indigestion ■ nausea ■ gas ■ belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering
from chicken pox or flu-like symptoms should not use this product.
When using this product, if changes in behavior with nausea and
vomiting occur, consult a doctor because these symptoms could be an
early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have ■ an ulcer
■ a bleeding problem ■ black or bloody stool

Ask a doctor before use if you have

- fever ■ mucus in stool

Ask a doctor or pharmacist before use if you are

taking any drug for ■ anticoagulation (thinning of the blood)
■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless, darkening
of the stool and/or tongue may occur

Stop use and ask a doctor if

- diarrhea lasts more than 2 days
- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs

If pregnant or breast feeding, ask health professional before use.

Keep out of reach of children. In case of overdose, get medical
help or contact a Poison Control Center right away.

Directions ■ Shake well before use

- use dose cup or tablespoon (TBSP)

- adults and children 12 years and over: 1 dose (30 mL or 2 TBSP) every 1 hour as needed
- do not exceed 4 doses (120 mL or 8 TBSP) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

■ **each 30 mL dose cup contains:** sodium 13 mg, salicylate 455 mg ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C) ■ low sodium

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, purified water, saccharin sodium, salicylic acid, sodium salicylate, xanthan gum

Marc's Product Label

NDC 68998-558-28

*** Compare to the active
ingredients in Pepto-Bismol™
Maximum Strength**

Marc's

ULTRA STRENGTH

Stomach Relief

**Antidiarrheal / Upset Stomach Reliever
Bismuth subsalicylate**

5 Symptoms Relief of

- Nausea
- Upset Stomach
- Heartburn
- Diarrhea
- Indigestion

12 FL OZ (354 mL)

TAMPER EVIDENT: Do not use if imprinted shrinkband is missing or broken

***This product is not manufactured or distributed by Procter & Gamble, Inc" distributor of Pepto Bismol™**

**Distributed by:
Marc Glassman, Inc.
West 130th Street
Cleveland, OH 44130**

LR-144

**Lot.
Exp.**

Drug Facts (continued)

Stop use and ask a doctor if

- diarrhea lasts more than 2 days
- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs

If pregnant or breast feeding, ask health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ Shake well before use

- use dose cup or tablespoon (TBSP)
- adults and children 12 years and over: 1 dose (30 mL or 2 TBSP) every 1 hour as needed
- do not exceed 4 doses (120 mL or 8 TBSP) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each 30 mL dose cup contains: sodium 13 mg, salicylate 455 mg ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C) ■ low sodium


Inactive ingredients benzoic acid, D&C red #22, D&C red # 28, flavor, purified water, saccharin sodium, salicylic acid, sodium salicylate, xanthan gum

*This product is not manufactured or distributed by Procter & Gamble, Inc., distributor of Pepto Bismol™

Distributed by:
Marc Glassman, Inc.
West 130th Street
Cleveland, OH 44130

LR-144

Lot.
Exp.



0 94442 00184 8

NDC 68998-558-28

* Compare to the active ingredient in Pepto-Bismol™ Maximum Strength

Marc's

ULTRA STRENGTH

Stomach Relief

Antidiarrheal / Upset Stomach Reliever

Bismuth subsalicylate

5 Symptoms Relief of:

- Nausea
- Upset Stomach
- Heartburn
- Diarrhea
- Indigestion

12 FL OZ (354 mL)

TAMPER EVIDENT: Do not use if imprinted shrinkband is missing or broken

Drug Facts

Active ingredient
(in each 30 mL dose cup or 2 tablespoons)

Bismuth subsalicylate 1050 mg

Purposes
Upset stomach reliever and antidiarrheal

Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence of food and drink including: ■ heart burn ■ indigestion ■ nausea ■ gas ■ belching

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have ■ an ulcer ■ a bleeding problem ■ black or bloody stool

Ask a doctor before use if you have ■ fever ■ mucus in stool

Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Drug Facts (continued)

Stop use and ask a doctor if

- diarrhea lasts more than 2 days
- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs

If pregnant or breast feeding, ask health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ Shake well before use

- use dose cup or tablespoon (TBSP)
- adults and children 12 years and over: 1 dose (30 mL or 2 TBSP) every 1 hour as needed
- do not exceed 4 doses (120 mL or 8 TBSP) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each 30 mL dose cup contains: sodium 13 mg, salicylate 455 mg ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C) ■ low sodium

Inactive ingredients benzoic acid, D&C red #22, D&C red # 28, flavor, purified water, saccharin sodium, salicylic acid, sodium salicylate, xanthan gum

*This product is not manufactured or distributed by Procter & Gamble, Inc., distributor of Pepto Bismol™

Distributed by:
Marc Glassman, Inc.
West 130th Street
Cleveland, OH 44130

LR-144

Lot.
Exp.



0 94442 00184 8

NDC 68998-558-28

* Compare to the active ingredient in Pepto-Bismol™ Maximum Strength

Marc's

ULTRA STRENGTH

Stomach Relief

Antidiarrheal / Upset Stomach Reliever

Bismuth subsalicylate

5 Symptoms Relief of:

- Nausea
- Upset Stomach
- Heartburn
- Diarrhea
- Indigestion

12 FL OZ (354 mL)

TAMPER EVIDENT: Do not use if imprinted shrinkband is missing or broken

Drug Facts

Active ingredient
(in each 30 mL dose cup or 2 tablespoons)

Bismuth subsalicylate 1050 mg

Purposes
Upset stomach reliever and antidiarrheal

Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence of food and drink including: ■ heart burn ■ indigestion ■ nausea ■ gas ■ belching

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have ■ an ulcer ■ a bleeding problem ■ black or bloody stool

Ask a doctor before use if you have ■ fever ■ mucus in stool

Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

res

STOMACH RELIEF, ULTRA STRENGTH

bismuth subsalicylate suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68998-558
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	1050 mg in 30 mL

Inactive Ingredients	
Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
SODIUM SALICYLATE (UNII: WQ1H85SYP)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color	pink	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68998-558-28	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/15/2020	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	12/15/2020	

Labeler - Marc Glassman, Inc. (094487477)

Revised: 12/2025

Marc Glassman, Inc.