

CIPROFLOXACIN- ciprofloxacin solution

Direct_Rx

CIPROFLOXACIN

Ciprofloxacin Ophthalmic Solution is a synthetic, sterile, multiple dose, antimicrobial for topical ophthalmic use. Ciprofloxacin is a fluoroquinolone antibacterial active against a broad spectrum of gram-positive and gram-negative ocular pathogens. It is available as the monohydrochloride monohydrate salt of 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinoline-carboxylic acid. It is a faint to light yellow crystalline powder with a molecular weight of 385.8. Its empirical formula is $C_{17}H_{18}FN_3O_3 \cdot HCl \cdot H_2O$ and its chemical structure is as follows:

[Ciprofloxacin hydrochloride]

Ciprofloxacin hydrochloride

Ciprofloxacin differs from other quinolones in that it has a fluorine atom at the 6-position, a piperazine moiety at the 7-position, and a cyclopropyl ring at the 1-position.

Each mL of Ciprofloxacin Ophthalmic Solution contains: Active: ciprofloxacin HCl 3.5 mg equivalent to 3 mg base. Preservative: benzalkonium chloride 0.006%. Inactives: sodium acetate, acetic acid, mannitol 4.6%, edetate disodium 0.05%, hydrochloric acid and/or sodium hydroxide (to adjust pH) and water for injection. The pH is approximately 4.5 and the osmolality is approximately 300 mOsm.

Systemic Absorption: A systemic absorption study was performed in which Ciprofloxacin Ophthalmic Solution was administered in each eye every two hours while awake for two days followed by every four hours while awake for an additional 5 days. The maximum reported plasma concentration of ciprofloxacin was less than 5 ng/mL. The mean concentration was usually less than 2.5 ng/mL.

Microbiology: Ciprofloxacin has in vitro activity against a wide range of gram-negative and gram-positive organisms. The bactericidal action of ciprofloxacin results from interference with the enzyme DNA gyrase which is needed for the synthesis of bacterial DNA.

Ciprofloxacin has been shown to be active against most strains of the following organisms both in vitro and in clinical infections (see Indications and Usage):

Gram-Positive:

Staphylococcus aureus

Staphylococcus epidermidis

Streptococcus pneumoniae

Streptococcus (Viridans Group)

Gram-Negative:

Haemophilus influenzae

Pseudomonas aeruginosa

Serratia marcescens

Ciprofloxacin has been shown to be active in vitro against most strains of the following organisms, however, the clinical significance of these data is unknown:

Gram-Positive:

Enterococcus faecalis (Many strains are only moderately susceptible)

Staphylococcus haemolyticus

Staphylococcus hominis

Staphylococcus saprophyticus

Streptococcus pyogenes

Gram-Negative

Acinetobacter calcoaceticus subsp. *anitratus*

Aeromonas caviae

Aeromonas hydrophila

Brucella melitensis

Campylobacter coli

Campylobacter jejuni

Citrobacter diversus

Citrobacter freundii

Edwardsiella tarda

Enterobacter aerogenes

Enterobacter cloacae

Escherichia coli

Haemophilus ducreyi

Haemophilus parainfluenzae

Klebsiella pneumoniae

Klebsiella oxytoca

Legionella pneumophila

Moraxella (*Branhamella*) *catarrhalis*

Morganella morganii

Neisseria gonorrhoeae

Neisseria meningitidis

Pasteurella multocida

Proteus mirabilis

Proteus vulgaris

Providencia rettgeri

Providencia stuartii

Salmonella enteritidis

Salmonella typhi

Shigella sonnei

Shigella flexneri

Vibrio cholera

Vibrio parahaemolyticus

Vibrio vulnificus

Yersinia enterocolitica

Other Organisms:

Chlamydia trachomatis (only moderately susceptible) and *Mycobacterium tuberculosis* (only moderately susceptible).

Most strains of *Pseudomonas cepacia* and some strains of *Pseudomonas maltophilia* are resistant to ciprofloxacin as are most anaerobic bacteria, including *Bacteroides fragilis* and *Clostridium difficile*.

The minimal bactericidal concentration (MBC) generally does not exceed the minimal inhibitory concentration (MIC) by more than a factor of 2. Resistance to ciprofloxacin in vitro usually develops slowly (multiple-step mutation).

Ciprofloxacin does not cross-react with other antimicrobial agents such as beta-lactams or aminoglycosides; therefore, organisms resistant to these drugs may be susceptible to ciprofloxacin.

Ciprofloxacin Ophthalmic Solution is indicated for the treatment of infections caused by susceptible strains of the designated microorganisms in the conditions listed below:

Corneal Ulcers:

Pseudomonas aeruginosa

Serratia marcescens *

Staphylococcus aureus

Staphylococcus epidermidis

Streptococcus pneumoniae

Streptococcus (Viridans Group) *

Conjunctivitis:

Haemophilus influenzae

Staphylococcus aureus

Staphylococcus epidermidis

Streptococcus pneumoniae

*Efficacy for this organism was studied in fewer than 10 infections.

A history of hypersensitivity to ciprofloxacin or any other component of the medication is a contraindication to its use. A history of hypersensitivity to other quinolones may also contraindicate the use of ciprofloxacin.

NOT FOR INJECTION INTO THE EYE.

Serious and occasionally fatal hypersensitivity (anaphylactic) reactions, some following the first dose, have been reported in patients receiving systemic quinolone therapy. Some reactions were accompanied by cardiovascular collapse, loss of consciousness, tingling, pharyngeal or facial edema, dyspnea, urticaria, and itching. Only a few patients had a history of hypersensitivity reactions. Serious anaphylactic reactions require immediate emergency treatment with epinephrine and other resuscitation measures, including oxygen, intravenous fluids, intravenous antihistamines, corticosteroids, pressor amines and airway management, as clinically indicated. Remove contact lenses before using.

General:

As with other antibacterial preparations, prolonged use of ciprofloxacin may result in overgrowth of nonsusceptible organisms, including fungi. If superinfection occurs, appropriate therapy should be initiated. Whenever clinical judgment dictates, the patient should be examined with the aid of magnification, such as slit lamp biomicroscopy and, where appropriate, fluorescein staining.

Ciprofloxacin should be discontinued at the first appearance of a skin rash or any other sign of hypersensitivity reaction.

In clinical studies of patients with bacterial corneal ulcer, a white crystalline precipitate located in the superficial portion of the corneal defect was observed in 35 (16.6%) of 210 patients. The onset of the precipitate was within 24 hours to 7 days after starting therapy. In one patient, the precipitate was immediately irrigated out upon its appearance. In 17 patients, resolution of the precipitate was seen in 1 to 8 days (seven within the first 24 to 72 hours), in five patients, resolution was noted in 10 to 13 days. In nine patients, exact resolution days were unavailable; however, at follow-up examinations, 18 to 44 days after onset of the event, complete resolution of the precipitate was noted. In three patients, outcome information was unavailable. The precipitate did not preclude continued use of ciprofloxacin, nor did it adversely affect the clinical course of the ulcer or visual outcome. (see Adverse Reactions).

Information for patients:

Do not touch dropper tip to any surface, as this may contaminate the solution.

Drug Interactions:

Specific drug interaction studies have not been conducted with ophthalmic ciprofloxacin. However, the systemic administration of some quinolones has been shown to elevate plasma concentrations of theophylline, interfere with the metabolism of caffeine, enhance the effects of the oral anticoagulant, warfarin, and its derivatives, and has been associated with transient elevations in serum creatinine in patients receiving cyclosporine concomitantly.

Carcinogenesis, Mutagenesis, Impairment of Fertility:

Eight in vitro mutagenicity tests have been conducted with ciprofloxacin and the test results are listed below:

Salmonella/Microsome Test (Negative)

E. coli DNA Repair Assay (Negative)

Mouse Lymphoma Cell Forward Mutation Assay (Positive)

Chinese Hamster V79 Cell HGPRT Test (Negative)

Syrian Hamster Embryo Cell Transformation Assay (Negative)

Saccharomyces cerevisiae Point Mutation Assay (Negative)

Saccharomyces cerevisiae Mitotic Crossover and Gene Conversion Assay (Negative)

Rat Hepatocyte DNA Repair Assay (Positive)

Thus, two of the eight tests were positive, but the results of the following three in vivo test systems gave negative results:

Rat Hepatocyte DNA Repair Assay

Micronucleus Test (Mice)

Dominant Lethal Test (Mice).

Long term carcinogenicity studies in mice and rats have been completed. After daily oral dosing for up to two years, there is no evidence that ciprofloxacin had any carcinogenic or tumorigenic effects in these species.

Pregnancy: Pregnancy Category C:

Reproduction studies have been performed in rats and mice at doses up to six times the usual daily human oral dose and have revealed no evidence of impaired fertility or harm to the fetus due to ciprofloxacin. In rabbits, as with most antimicrobial agents, ciprofloxacin (30 and 100 mg/kg orally) produced gastrointestinal disturbances resulting in maternal weight loss and an increased incidence of abortion. No teratogenicity was observed at either dose.

After intravenous administration, at doses up to 20 mg/kg, no maternal toxicity was produced and no embryotoxicity or teratogenicity was observed. There are no adequate and well controlled studies in pregnant women. Ciprofloxacin Ophthalmic Solution should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers:

It is not known whether topically applied ciprofloxacin is excreted in human milk. However, it is known that orally administered ciprofloxacin is excreted in the milk of lactating rats and oral ciprofloxacin has been reported in human breast milk after a single 500 mg dose. Caution should be exercised when Ciprofloxacin Ophthalmic Solution is administered to a nursing mother.

Pediatric Use:

The safety and effectiveness of Ciprofloxacin ophthalmic solution 0.3% have been

established in all ages. Use of Ciprofloxacin ophthalmic solution is supported by evidence from adequate and well controlled studies of Ciprofloxacin ophthalmic solution in adults, children and neonates [see Clinical Studies]. Although ciprofloxacin and other quinolones cause arthropathy in immature animals after oral administration, topical ocular administration of ciprofloxacin to immature animals did not cause any arthropathy and there is no evidence that the ophthalmic dosage form has any effect on the weight bearing joints.

Geriatric Use:

No overall differences in safety or effectiveness have been observed between elderly and younger patients.

The most frequently reported drug related adverse reaction was local burning or discomfort. In corneal ulcer studies with frequent administration of the drug, white crystalline precipitates were seen in approximately 17% of patients (see Precautions). Other reactions occurring in less than 10% of patients included lid margin crusting, crystals/scales, foreign body sensation, itching, conjunctival hyperemia and a bad taste following instillation. Additional events occurring in less than 1% of patients included corneal staining, keratopathy/keratitis, allergic reactions, lid edema, tearing, photophobia, corneal infiltrates, nausea and decreased vision.

To report SUSPECTED ADVERSE REACTIONS, contact LEADING PHARMA AT 1-844-740-7500 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

A topical overdose of ciprofloxacin Ophthalmic Solution may be flushed from the eye(s) with warm tap water.

Corneal Ulcers:

The recommended dosage regimen for the treatment of corneal ulcers is two drops into the affected eye every 15 minutes for the first six hours and then two drops into the affected eye every 30 minutes for the remainder of the first day. On the second day, instill two drops in the affected eye hourly. On the third through the fourteenth day, place two drops in the affected eye every four hours. Treatment may be continued after 14 days if corneal re-epithelialisation has not occurred.

Bacterial Conjunctivitis:

The recommended dosage regimen for the treatment of bacterial conjunctivitis is one or two drops instilled into the conjunctival sac(s) every two hours while awake for two days and one or two drops every four hours while awake for the next five days.

As a sterile ophthalmic solution: 2.5 mL, 5 mL and 10 mL in translucent LDPE bottle with insert cap assembly comprising of a tan colored HDPE screw cap over a LDPE nozzle with tamper evident LDPE dust cover sealing the bottle cap.

2.5 mL -

5 mL -

10 mL -

STORAGE CONDITIONS

Store at 25°C (77°F); excursions permitted 15°C to 30°C (59°F to 86°F) [See USP

controlled room temperature]. Retain in carton until contents are used and protect from light.

ANIMAL PHARMACOLOGY

Ciprofloxacin and related drugs have been shown to cause arthropathy in immature animals of most species tested following oral administration. However, a one month topical ocular study using immature Beagle dogs did not demonstrate any articular lesions.

CLINICAL STUDIES

Following therapy with Ciprofloxacin ophthalmic solution, 76% of the patients with corneal ulcers and positive bacterial cultures were clinically cured and complete re-epithelialisation occurred in about 92% of the ulcers.

In 3 and 7 day multicenter clinical trials, 52% of the patients with conjunctivitis and positive conjunctival cultures were clinically cured and 70-80% had all causative pathogens eradicated by the end of treatment.

In a randomized, double-masked, multicenter, parallel-group clinical trial of pediatric patients with bacterial conjunctivitis, between birth and 31 days of age, patients were dosed with Ciprofloxacin ophthalmic solution or another anti-infective agent. Clinical outcomes for the trial demonstrated a clinical cure rate of 80% at Day 9 and a microbiological eradication success rate of 85% at Day 9.

Please note that microbiologic eradication does not always correlate with clinical outcome in anti-infective trials.

Rx Only

Distributed by:

Leading Pharma, LLC.

Fairfield, NJ07004

Manufactured By:

FDC Limited,

B-8, MIDC Industrial Area,

Waluj, Aurangabad - 431 136,

Maharashtra, India

Revision: 12/2017

STERILE OPHTHALMIC SOLUTION

CIPROFLOXACIN OPHTHALMIC SOLUTION USP 0.3% as base

INSTRUCTIONS FOR USE

Read this Instruction for Use before you start using Ciprofloxacin ophthalmic solution and each time you get a refill. There may be new information. This information does not take the place of talking to your doctor about your medical condition or your treatment.

Use Ciprofloxacin ophthalmic solution as prescribed by your doctor. Ask your doctor if

you have questions about how to use Ciprofloxacin ophthalmic solution.

Important information:

Ciprofloxacin ophthalmic solution is for use in the eye only. Ciprofloxacin ophthalmic solution should not be injected into the eye.

If you use other medicines in your eye, wait at least 10 minutes between using Ciprofloxacin ophthalmic solution and your other eye medicines.

If you wear contact lenses, remove them before using Ciprofloxacin ophthalmic solution. Wait at least 15 minutes after using Ciprofloxacin ophthalmic solution before placing your contact lenses back in your eyes.

Do not touch your eye, fingers, or other surfaces with the tip of the Ciprofloxacin ophthalmic solution bottle. You may get bacteria on the tip of the bottle that can cause you to get an eye infection. This eye infection can lead to serious eye damage or vision loss. If you think you have gotten bacteria on the tip of the bottle or you have an eye infection, call your doctor right away.

Wash your hands before each use.

[Fig 1]

Before you use a bottle of Ciprofloxacin ophthalmic solution for the first time:

Check the expiration date on the bottle before use. Do not use Ciprofloxacin ophthalmic solution if the expiration date has passed.

Check that the seal on the dust cover is not broken or missing. Do not use the medicine if the seal is broken or missing. Call your doctor or pharmacist.

Each time you use Ciprofloxacin ophthalmic solution:

Step 1: Turn the dust cover clockwise to break the seal.

[fig 2]

Step 2: Remove the dust cover and throw it away.

[fig 2]

Step 3: Turn the tan coloured cap anti-clockwise and remove. Place the cap on a clean flat surface.

[fig 4]

Step 4: Tilt your head backwards.

[fig 5]

Step 5: Turn the bottle upside down and place the tip of the bottle close to your eye. Do not touch your eye with the tip. Pull your lower eyelid downward and look up. Gently squeeze the bottle and let 1 drop fall into the space between your lower eyelid and your eye. If a drop misses your eye, repeat Step 5.

[fig 6]

Step 6: Replace the tan coloured cap back on the bottle and turn it clockwise to close.

[fig 7]

Step 7: Remove any extra solution from skin around the eyes with a tissue.

[fig 8]

If your doctor has told you to use drops in both eyes, repeat Step 5 for your other eye. The Ciprofloxacin ophthalmic solution bottle tip is made to give a certain amount of medicine in 1 drop. Do not make the opening of the bottle tip bigger or you may get too much Ciprofloxacin ophthalmic solution.

After you use all of the doses of Ciprofloxacin ophthalmic solution that your doctor has prescribed, there will be some medicine left in the bottle. Do not use the extra medicine in the bottle. Throw the bottle away.

How should I store Ciprofloxacin ophthalmic solution?

Store at 25°C (77°F); excursions permitted 15°C to 30°C (59°F to 86°F) [See USP controlled room temperature].

Retain in carton until contents are used and protect from light

Keep Ciprofloxacin ophthalmic solution and all medicines out of the reach of children.

This Instruction for Use has been approved by the U.S. Food and Drug Administration.

Distributed by:

Leading Pharma, LLC.

Fairfield, NJ07004

Manufactured By:

FDC Limited,

B-8, MIDC Industrial Area,

Waluj, Aurangabad - 431 136,

Maharashtra, India

Revision: 12/2017

CIPROFLOXACIN OPTH SOLUTION 0.3% 5 mL

Generic For: **CILOXIN**

Each mL contains: Active: ciprofloxacin hcl 3.5mg equivalent to 3mg base

Lot# 17JY1908
Prod# 4144-003-05

Discard After: 2/28/21
61919-795-05
17JY1908
2/28/21 DAWSONVILLE,
AXMJN GA 30534

Caution: Federal law prohibits transfer of this drug to any person other than the patient for whom it was prescribed.

RX ONLY-KEEP OUT OF REACH OF CHILDREN

Dosage: See package insert. Store between 68-77 degrees F

AXMJN

NDC 61919-795-05

DIST BY: Leading Pharma, LLC
Fairfield, NJ 07004
NDC 69315-308-05

Mfg Lot: 089C071
PM 7/17/2019 3207856

Package and Distributed By: **DIRECT Rx**

CIPROFLOXACIN OPTH SOLUTION 0.3%
NDC 61919-795-05 5 mL
Lot 17JY1908 Exp Date 02/21
Mfg NDC 69315-308-05

CIPROFLOXACIN OPTH SOLUTION 0.3%
NDC 61919-795-05 5 mL
Lot 17JY1908 Exp Date 02/21
Mfg NDC 69315-308-05

CIPROFLOXACIN OPTH SOLUTION 0.3%
NDC 61919-795-05 5 mL
Lot 17JY1908 Exp Date 02/21
Mfg NDC 69315-308-05

CIPROFLOXACIN OPTH SOLUTION 0.3%
NDC 61919-795-05 5 mL
Lot 17JY1908 Exp Date 02/21
Mfg NDC 69315-308-05

Caution: Federal law prohibits transfer of this drug to any person other than the patient for whom it was prescribed. Dosage: See package insert. Store between 68-77 degrees F. For RX ONLY. Keep out of reach of children.

D NDC 61919-795-25

CIPROFLOXACIN OPTH SOLUTION

0.3% **2.5 mL**

Generic For: **CILOXAN**
 Each mL contains: 3.5mg ciprofloxacin hydrochloride equivalent to ciprofloxacin base 3 mg

Lot# 220C1914
 Prod# 4144-025-25
 Packaged and Distributed By: **DIRECT B**

Discard After: 3/31/21
 61919-795-25
 220C1914
 3/31/21
 AZXTB Dawsonville, GA 30534

Mfg Lot: 089D103
 8W 10/22/2019 1145557
 Mfg NDC 69315-308-02

CIPROFLOXACIN OPTH SOLUTION
 NDC 61919-795-25 2.5 mL
 Lot 220C1914 Exp 3/31/21
 Mfg NDC 69315-308-02

CIPROFLOXACIN OPTH SOLUTION
 NDC 61919-795-25 2.5 mL
 Lot 220C1914 Exp 3/31/21
 Mfg NDC 69315-308-02

CIPROFLOXACIN OPTH SOLUTION
 NDC 61919-795-25 2.5 mL
 Lot 220C1914 Exp 3/31/21
 Mfg NDC 69315-308-02

CIPROFLOXACIN OPTH SOLUTION
 NDC 61919-795-25 2.5 mL
 Lot 220C1914 Exp 3/31/21
 Mfg NDC 69315-308-02

Dist. By: Leading Pharma, LLC
 Fairfield, NJ 07004
 NDC 69315-308-02

CIPROFLOXACIN

ciprofloxacin solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:61919-795(NDC:69315-308)
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CIPROFLOXACIN HYDROCHLORIDE (UNII: 4BA73M5E37) (CIPROFLOXACIN - UNII:5E8K9I0O4U)	CIPROFLOXACIN	3 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
SODIUM ACETATE ANHYDROUS (UNII: NVG71ZZ7P0)	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
ACETIC ACID (UNII: Q40Q9N063P)	
MANNITOL (UNII: 3OWL53L36A)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61919-795-05	5 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/23/2019	
2	NDC:61919-795-25	25 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/23/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077568	07/23/2019	

Labeler - Direct_Rx (079254320)

Registrant - Direct_Rx (079254320)

Establishment

Name	Address	ID/FEI	Business Operations
Direct_Rx		079254320	repack(61919-795)

Revised: 5/2023

Direct_Rx