

IBUPROFEN CA- ibuprofen tablet, coated
TARGET CORPORATION

600R -TARGET IBUPROFEN 200 MG CAPLETS NDC 11673-990

ACTIVE INGREDIENT(S)

Ibuprofen 200 mg (NSAID)*

* nonstreoidal anti-inflammatory drug

SILICON DIOXIDE

CROSCARMELLOSE SODIUM

FERRIC OXIDE RED

MAGNESIUM STEARATE

CELLULOSE, MICROCRYSTALLINE

TALC

TITANIUM DIOXIDE

STARCH, PREGELATINIZED CORN

POLYVINYL ALCOHOL

POLYETHYLENE GLYCOL

PURPOSE

Pain reliever / fever reducer

USE(S)

temporarily relieves minor aches and pains due to :

backache

headache

menstrual cramps

minor pain of arthritis

muscular aches

the common cold

toothache

temporarily reduces fever

DIRECTIONS

do not take more than directed

the smallest effective dose should be used

adults and children 12 years and older: take 1 caplet every 4 to 6 hours while symptoms persist

if pain or fever does not respond to 1 caplet, 2 caplets may be used

do not exceed 6 caplets in 24 hours, unless directed by a doctor.

Children under 12 years: ask a doctor

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

WARNINGS

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- asthma (wheezing)
- blisters
- facial swelling
- hives
- rash
- shock
- skin reddening

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAID's except aspirin increases the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or longer than directed.

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-990
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	brown	Score	no score
Shape	CAPSULE ((CAPLET))	Size	15mm
Flavor		Imprint Code	117
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-990-42	24 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
2	NDC:11673-990-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091239	01/01/2020	

Labeler - TARGET CORPORATION (006961700)**Registrant** - TIME CAP LABORATORIES, INC (037005209)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LTD		925822975	manufacture(11673-990)

Revised: 5/2020

TARGET CORPORATION