

**ANTI DIARRHEAL ANTI GAS- loperamide hydrochloride, simethicone tablet**  
**Rite Aid Corporation**

-----  
**Rite Aid Corporation Anti-Diarrheal Anti-Gas Drug Facts**

**Active ingredients (in each caplet)**

Loperamide HCl 2 mg

Simethicone 125 mg

**Purposes**

Anti-diarrheal

Anti-gas

**Uses**

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

- if you have bloody or black stool
- if you have difficulty swallowing

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse

- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- **each caplet contains:** calcium 165 mg, sodium 3 mg
- store between 20°-25° C (68°-77° F). Protect from light.
- **do not use if printed blister unit is broken or torn**

**Inactive ingredients**

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin, microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

**Questions or comments?**

**1-800-719-9260**

**Package/Label Principal Display Panel**

Compare to the active ingredients of Imodium® Multi-Symptom Relief

ANTI-DIARRHEAL/ANTI-GAS

LOPERAMIDE HYDROCHLORIDE & SIMETHICONE TABLETS, 2 mg/125 mg

ANTI-DIARRHEAL/ANTI-GAS

Multi-symptom relief caplet

Relieves symptoms of diarrhea plus:

cramps & pressure

bloating/gas

ACTUAL SIZE

24 CAPLETS\*

\*CAPSULE-SHAPED TABLETS

CODE AREA

0R162 83 C1

**Drug Facts** (continued)

**Inactive ingredients:** asarilium epol asilium, crocarmellos sodium, dibasic calcium phosphate, maleate, microcrystalline cellulose, mannitol and hydroxypropyl methylcellulose.

**Questions or comments?** 1-800-771-9928



**Drug Facts**

**Active ingredients (in each caplet)** Loperamide HCl 2 mg, Simethicone, Antidiarrheal, Antigas

**Warnings**

- Do not use if you have bloody or black stool
- If you have difficulty swallowing
- Ask a doctor before use if you have a history of abnormal heart rhythm
- Ask a doctor before use if you are taking a prescription drug, Loperamide may interact with certain prescription drugs.
- With use, you may experience dizziness, weakness or drowsiness may occur. Be careful when driving or operating machinery.
- Stop use and ask a doctor if diarrhea lasts for more than 2 days
- Symptoms get worse
- You are having difficulty swallowing or you are having a serious condition
- You are having difficulty swallowing or you are having a serious condition

**Directions**

Take only on an empty stomach, 1 hour before or 2 hours after a meal.

Take with a full (8 oz) glass of water.

Take right dose once a day below. If possible, use a weight to dose.

Children and adults: 2 caplets after the first loose stool, but no more than 4 caplets in 24 hours.

Children 9-11 years: 1 caplet after the first loose stool, but no more than 3 caplets in 24 hours.

Children 6-8 years: 1 caplet after the first loose stool, but no more than 2 caplets in 24 hours.

Children 2-5 years: ask a doctor.

Children under 2 years: do not use.

Do not use if you are taking any other medicine for diarrhea.

**Other information**

Each caplet contains calcium 165 mg, sodium 3 mg, and contains 20% to 25% alcohol.

Do not use if you are taking any other medicine for diarrhea.

Do not use if you are taking any other medicine for diarrhea.

# ANTI-DIARRHEAL /ANTI-GAS

LOPERAMIDE HYDROCHLORIDE & SIMETHICONE TABLETS, 2 mg/125 mg

ANTI-DIARRHEAL/ANTI-GAS

NDC 11822-1003-3

Compare to the active ingredients of Imodium® Multi-Symptom Relief\*

# ANTI-DIARRHEAL /ANTI-GAS

LOPERAMIDE HYDROCHLORIDE & SIMETHICONE TABLETS, 2 mg/125 mg

ANTI-DIARRHEAL/ANTI-GAS

Multi-symptom relief caplet  
Relieves symptoms of diarrhea *plus*:  
cramps & pressure  
bloating/gas



ACTUAL SIZE

**24** CAPLETS\*\*  
\*\*CAPSULE-SHAPED TABLETS



\* This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® Multi-Symptom Relief.

**DISTRIBUTED BY:**  
RITE AID, 30 HUNTER LANE,  
CAMP HILL, PA 17011  
www.riteaid.com

**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.

# ANTI DIARRHEAL ANTI GAS

loperamide hydrochloride, simethicone tablet

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-1003
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	125 mg

## Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

## Product Characteristics

Color	WHITE	Score	2 pieces
Shape	OVAL	Size	17mm
Flavor		Imprint Code	L0R1;2;125
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-1003-2	12 in 1 CARTON	01/03/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-1003-1	1 in 1 CARTON	01/03/2019	
2		42 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:11822-1003-3	24 in 1 CARTON	02/24/2020	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209837	01/03/2019	

**Labeler** - Rite Aid Corporation (014578892)

Revised: 2/2020

Rite Aid Corporation