

ACETAMINOPHEN- acetaminophen tablet, coated
TARGET CORPORATION

341R-TGT APAP 500 MG 11673-838

Drug Facts

Drug Facts

Active Ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed.

See overdose warning

adults and children 12 years and over

-take 2 caplets every 6 hours while symptoms last

-do not take more than 6 caplets in 24 hours, unless directed by a doctor

-do not take for more than 10 days unless directed by a doctor

children under 12 years

-ask a doctor

Other information

- SODIUM FREE
- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

Inactive ingredients

hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate*
stearic acid, titanium dioxide

*may contain this ingredient

Questions or comments? Call 1-800-910-6874

Extra Strength Acetaminophen

NDC 11673-838-01

500 mg
Pain Reliever / Fever Reducer
For Adults



100 CAPLETS,** 500 mg EACH
(*CAPSULE-SHAPED TABLETS)

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

 If a skin reaction occurs, stop use and seek medical help right away.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have: liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if:

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed. See overdose warning

adults and children 12 years and over:

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor

children under 12 years: ask a doctor

Other information

■ SODIUM FREE

■ store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)

■ use by expiration date on package

Questions or comments? Call 1-800-910-6874

TM & ©2024

094 01 0103 R00

C-00262-01-034-0000

Dist. by Target Corp.

Mpls, MN 55403

341R 1023

Lot No.:

Exp. Date:

UNVARNISHED AREA

Coating/Print Omit Area



THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient **Purpose**
(in each caplet)

Acetaminophen 500 mg..... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

 If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Drug Facts (continued)

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed. See overdose warning

adults and children 12 years and over

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor

children under 12 years ask a doctor

Drug Facts (continued)

Other information

- SODIUM FREE
- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

Inactive ingredients

hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid, titanium dioxide
*may contain this ingredient

Questions or comments?

Call 1-800-910-6874

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Extra Strength Tylenol®

Satisfaction guaranteed - Love It or your money back.

Distributed by Target Corporation
Minneapolis, MN 55403
TM & ©2024 Target Brands, Inc.

341R 1023

Compare to active ingredient in Extra Strength Tylenol® NDC 11673-838-01

Extra Strength Acetaminophen

500 mg
Pain Reliever / Fever Reducer
For Adults



100 CAPLETS,** 500 mg EACH
(*CAPSULE-SHAPED TABLETS)

Actual Size
100 Caplets

Coating Omit

Coating/Print Omit Area



Extra Strength Acetaminophen
500 mg
Pain Reliever / Fever Reducer

Compare to active ingredient in Extra Strength Tylenol®†

Extra Strength Acetaminophen

500 mg
Pain Reliever / Fever Reducer
For Adults



Actual Size

500 CAPLETS* 500 mg EACH
(*CAPSULE-SHAPED TABLETS)

NDC 11673-838-50

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 500 mg.....Pain reliever/
fever reducer

Uses ■ temporarily relieves minor aches and pains due to: ■ the common cold ■ headache ■ backache ■ minor pain of arthritis ■ toothache ■ muscular aches ■ premenstrual and menstrual cramps ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing

Drug Facts (continued)

acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions Do not take more than directed. See overdose warning
adults and children 12 years and over ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not take for more than 10 days unless directed by a doctor

children under 12 years ask a doctor

Other information

■ SODIUM FREE ■ store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F) ■ use by expiration date on package

Inactive ingredients

hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid, titanium dioxide
*may contain this ingredient

Questions or comments?
Call 1-800-910-6874

094-01-0219-R00-C-00262-01-034-0000
The product is manufactured and distributed by Johnson & Johnson Consumer Inc., Medical Consumer Healthcare Division, over the registered trademark of Johnson & Johnson.
Dist. by Target Corp., 341R-1023
TM & © 2024
Target Brands, Inc.
Lot No.:
Exp. Date:
UNVARNISHED AREA
4 9483 34150 8

Extra Strength Acetaminophen

500 mg
Pain Reliever / Fever Reducer
For Adults



50 CAPLETS** 500 mg EACH
(*CAPSULE-SHAPED TABLETS)

NDC 11673-838-051

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Uses ■ temporarily relieves minor aches and pains due to:
■ the common cold ■ headache ■ backache
■ minor pain of arthritis ■ toothache ■ muscular aches
■ premenstrual and menstrual cramps
■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions Do not take more than directed. See overdose warning
adults and children 12 years and over ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not take for more than 10 days unless directed by a doctor
children under 12 years ask a doctor

Other information ■ SODIUM FREE ■ store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F) ■ use by expiration date on package
Questions or comments? Call 1-800-910-6874

094-01-0646-R00 Dist. by Target Corp. TM & © 2024 Target Brands, Inc.
C-00262-01-034-0000 Mpls., MN 55403
341R-1023

Lot No.:
Exp. Date:
UNVARNISHED AREA

Extra Strength Acetaminophen

500 mg
Pain Reliever / Fever Reducer
For Adults



225 CAPLETS** 500 mg EACH
(**CAPSULE-SHAPED TABLETS)

NDC 11673-838-26

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet)
Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses
temporarily relieves minor aches and pains due to:
the common cold ■ headache ■ backache
minor pain of arthritis ■ toothache ■ muscular aches
premenstrual and menstrual cramps
temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur

■ redness or swelling is present

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions Do not take more than directed. See overdose warning adults and children 12 years and over ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not take for more than 10 days unless directed by a doctor

Other information ■ SODIUM FREE ■ store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F) ■ use by expiration date on package

Questions or comments? Call 1-800-910-6874

08401004 R00 Dist. by Wm. S. Targis, Inc. TMS 06/2024

C-00226201-004-0000 JMS-16/95495 Target Brands, Inc. 344R 1023

UNVARNISHED AREA

Lot No.:
Exp. Date:

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each caplet)
Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses
temporarily relieves minor aches and pains due to:
the common cold ■ headache
backache ■ minor pain of arthritis
toothache ■ muscular aches
premenstrual and menstrual cramps
temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions
Do not take more than directed. See overdose warning

adults and children 12 years and over
■ take 2 caplets every 6 hours while symptoms last
■ do not take more than 6 caplets in 24 hours, unless directed by a doctor
■ do not take for more than 10 days unless directed by a doctor

children under 12 years
ask a doctor

Other information
■ SODIUM FREE
■ store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
■ use by expiration date on package

Inactive ingredients
hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid, titanium dioxide
*may contain this ingredient

Questions or comments?
Call 1-800-910-6874

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Extra Strength Tylenol.
344R 1023

Compare to active ingredient in Extra Strength Tylenol®

Extra Strength Acetaminophen

500 mg Pain Reliever / Fever Reducer
For Adults

up&up

50 CAPLETS** 500 mg EACH
**CAPSULE-SHAPED TABLETS

Actual Size
50 Caplets

Varnish/Print Omit Area

Varnish Omit

Varnish Omit

Lot No.:
Exp. Date:
Varnish Omit

094 01 06468 00 0-00226201-034-1000

8 497483 34705 3

Dist. by Wm. S. Targis, Inc. TMS 06/2024
M.P.S. 16/95495 Target Brands, Inc.

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING



ACETAMINOPHEN

acetaminophen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-838
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MINERAL OIL (UNII: T5L8T28FGP)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	TCL341
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-838-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
2	NDC:11673-838-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
3	NDC:11673-838-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
4	NDC:11673-838-26	225 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/01/2020	

Labeler - TARGET CORPORATION (006961700)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(11673-838)