

**ACETAMINOPHEN- acetaminophen tablet, coated**  
**TARGET CORPORATION**

-----  
**341R-TGT APAP 500 MG 11673-838**

***Drug Facts***

***Drug Facts***

***Active Ingredient (in each caplet)***

Acetaminophen 500 mg

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever.

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

***Do not use***

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

**Do not take more than directed.**

See overdose warning

**adults and children 12 years and over**

-take 2 caplets every 6 hours while symptoms last

-do not take more than 6 caplets in 24 hours, unless directed by a doctor

-do not take for more than 10 days unless directed by a doctor

**children under 12 years**

-ask a doctor

### ***Other information***

- SODIUM FREE
- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

### ***Inactive ingredients***

hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate\*  
stearic acid, titanium dioxide

\*may contain this ingredient

**Questions or comments? Call 1-800-910-6874**

# Extra Strength Acetaminophen

NDC 11673-838-01

500 mg  
Pain Reliever / Fever Reducer  
For Adults



100 CAPLETS,\*\* 500 mg EACH  
(\*\*CAPSULE-SHAPED TABLETS)

**RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**  
**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold ■ headache ■ backache
  - minor pain of arthritis ■ toothache
  - muscular aches ■ premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.  
**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have:** liver disease  
**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if** ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present

These could be signs of a serious condition.  
**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

**Do not take more than directed.** See overdose warning  
**adults and children 12 years and over:**

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor

**children under 12 years:** ask a doctor

**Other information** ■ SODIUM FREE  
■ store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)

■ use by expiration date on package

**Questions or comments?** Call 1-800-910-6874

TM & ©2024  
094 01 0103 R00

C-002262-01-034-0000  
Dist. by Target Corp.  
Mpls, MN 55403

Lot No.:

Exp. Date:

**UNVARNISHED AREA**

341R 1023



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

**RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**Drug Facts**

**Active ingredient** **Purpose**  
(in each caplet)  
Acetaminophen 500 mg.....Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold ■ headache
  - backache ■ minor pain of arthritis
  - toothache ■ muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
■ more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Drug Facts (continued)**

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

Do not take more than directed. See overdose warning

- adults and children 12 years and over**
  - take 2 caplets every 6 hours while symptoms last
  - do not take more than 6 caplets in 24 hours, unless directed by a doctor
  - do not take for more than 10 days unless directed by a doctor

**children under 12 years** ask a doctor

**Drug Facts (continued)**

**Other information**

- SODIUM FREE
- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

**Inactive ingredients**

hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate\*, stearic acid, titanium dioxide  
\*may contain this ingredient

**Questions or comments?**

Call 1-800-910-6874

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Extra Strength Tylenol®

Satisfaction guaranteed –  
Love it or your money back.

Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2024 Target Brands, Inc.

341R 1023

Compare to active ingredient in Extra Strength Tylenol®† NDC 11673-838-01

# Extra Strength Acetaminophen

500 mg  
Pain Reliever / Fever Reducer  
For Adults

Actual Size  
100 Caplets  
100 CAPLETS,\*\* 500 mg EACH  
(\*\*CAPSULE-SHAPED TABLETS)



Coating  
Omit

Coating/Print  
Omit Area



Extra Strength  
500 mg  
Acetaminophen  
Pain Reliever / Fever Reducer

Compare to active ingredient in Extra Strength Tylenol<sup>®</sup>\*\*

# Extra Strength Acetaminophen

500 mg  
Pain Reliever / Fever Reducer  
For Adults



up&up<sup>™</sup>

500 CAPLETS,\*\* 500 mg EACH  
(\*CAPSULE-SHAPED TABLETS)



Actual Size  
500 Caplets

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each caplet)</b> Acetaminophen 500 mg</p> <p><b>Purpose</b> Pain reliever/fever reducer</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> <li>the common cold</li> <li>headache</li> <li>backache</li> <li>minor pain of arthritis</li> <li>toothache</li> <li>muscular aches</li> <li>premenstrual and menstrual cramps</li> </ul> </li> <li>temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> <li>more than 4,000 mg of acetaminophen in 24 hours</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> <p><b>Allergy alert:</b> acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>itch</li> </ul> <p>If a skin reaction occurs, stop use and seek medical help right away.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are allergic to acetaminophen or any of the inactive ingredients in this product</li> </ul> <p><b>Ask a doctor before use if you have liver disease</b></p> <p><b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>pain gets worse or lasts more than 10 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>new symptoms occur</li> <li>redness or swelling is present</li> </ul> <p>These could be signs of a serious condition.</p> <p><b>If pregnant or breast-feeding, ask a health professional before use.</b></p> <p><b>Keep out of reach of children.</b></p> <p><b>Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not take more than directed (see overdose warning)</li> </ul> <p>adults and children: <ul style="list-style-type: none"> <li>take 2 caplets every 6 hours while symptoms last</li> <li>do not take more than 6 caplets in 24 hours, unless directed by a doctor</li> <li>do not take for more than 10 days unless directed by a doctor</li> </ul> <p>children under 12 years: <ul style="list-style-type: none"> <li>ask a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>SODIUM FREE</li> <li>store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)</li> <li>use by expiration date on package</li> </ul> <p><b>Inactive ingredients</b> hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid, titanium dioxide *may contain this ingredient</p> <p><b>Questions or comments?</b> Call 1-800-910-6874</p> </p></p>
--	--	--

\*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Extra Strength Tylenol<sup>®</sup>.

094-01-0120 R00  
NDC 11673-888-50

**NOT FOR RESALE**

Satisfaction guaranteed—  
Love it or your money back.

LOT:  
EXP:..

Varnish Omit Area

Compare to active ingredient in Extra Strength Tylenol<sup>®</sup>\*\*

# Extra Strength Acetaminophen

Twin Pack

500 mg  
Pain Reliever / Fever Reducer  
For Adults



up&up<sup>™</sup>

2 BOTTLES, 2 x 500 CAPLETS\*\*  
(\*CAPSULE-SHAPED TABLETS)



Actual Size  
500+500  
1,000  
Caplets

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each caplet)</b> Acetaminophen 500 mg</p> <p><b>Purpose</b> Pain reliever/fever reducer</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> <li>the common cold</li> <li>headache</li> <li>backache</li> <li>minor pain of arthritis</li> <li>toothache</li> <li>muscular aches</li> <li>premenstrual and menstrual cramps</li> </ul> </li> <li>temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> <li>more than 4,000 mg of acetaminophen in 24 hours</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul> <p><b>Allergy alert:</b> acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> <li>skin reddening</li> <li>blisters</li> <li>itch</li> </ul> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</p> <p><b>If you are allergic to acetaminophen or any of the inactive ingredients in this product, ask a doctor before use if you have liver disease</b></p> <p><b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>pain gets worse or lasts more than 10 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>new symptoms occur</li> <li>redness or swelling is present</li> </ul> <p>These could be signs of a serious condition.</p> <p><b>If pregnant or breast-feeding, ask a health professional before use.</b></p> <p><b>Keep out of reach of children.</b></p> <p><b>Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not take more than directed (see overdose warning)</li> </ul> <p>adults and children 12 years and over: <ul style="list-style-type: none"> <li>take 2 caplets every 6 hours while symptoms last</li> <li>do not take more than 6 caplets in 24 hours, unless directed by a doctor</li> <li>do not take for more than 10 days unless directed by a doctor</li> </ul> <p>children under 12 years: <ul style="list-style-type: none"> <li>ask a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>SODIUM FREE</li> <li>store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)</li> <li>use by expiration date on package</li> </ul> <p><b>Inactive ingredients</b> hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid, titanium dioxide *may contain this ingredient</p> <p><b>Questions or comments?</b> Call 1-800-910-6874</p> </p></p>
--	--

\*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Extra Strength Tylenol<sup>®</sup>.

Satisfaction guaranteed—  
Love it or your money back.

LOT:  
EXP:..

094-01-0120 R00

**NOT FOR RESALE**

Satisfaction guaranteed—  
Love it or your money back.

LOT:  
EXP:..

Varnish Omit Area

**RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**Drug Facts**

**Active ingredient** **Purpose**  
(in each caplet) **Pain reliever/fever reducer**

**Acetaminophen 500 mg**.....


**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

- **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take
  - more than 4,000 mg of acetaminophen in 24 hours
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks every day while using this product.
- Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
  - skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Varnish Omit  
Lot No.:  
Exp. Date:



TAMPER EVIDENCE DO NOT USE THIS PRODUCT IF THE UNPRINTED FOLDS SEAL OVER THE MOUTH OF THE BOTTLES CUT, TORN, BROKEN OR MISSING

Dist. by Targent Corp.  
Mpls., MN 55403

Targent Brands, Inc.

**Drug Facts (continued)**

- Do not use
  - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
  - if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- Do not take more than directed. See overdose warning.
- adults and children 12 years and over
  - take 2 caplets every 6 hours while symptoms last
  - do not take more than 6 caplets in 24 hours, unless directed by a doctor
  - do not take for more than 10 days unless directed by a doctor
- children under 12 years
  - ask a doctor

**Other information**

- SODIUM FREE
- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

**Inactive ingredients**

hydroxytoluene, mineral oil, polydioxane, pregelatinized starch, sodium starch glycolate\*, stearic acid, titanium dioxide \*may contain this ingredient

**Questions or comments?**

Call 1-800-910-8874  
This product is not manufactured or distributed by Johnson & Johnson. Contact J&J Consumer Relations, P.O. Box 100, New Brunswick, NJ 08903 for more information. ©2011 J&J. All rights reserved.

50-868-638-2-01T1:001

Compare to active ingredient in Extra Strength Tylenol®

**Extra Strength Acetaminophen**

500 mg Pain Reliever / Fever Reducer  
For Adults



50 CAPLETS,\*\* 500 mg EACH  
\*\*CAPSULE-SHAPED TABLETS



Actual Size  
50 Caplets

Varnish/Print Omit Area

Varnish Omit

Varnish Omit

**Extra Strength Acetaminophen**  
500 mg  
Pain Reliever / Fever Reducer  
d&D

# Extra Strength Acetaminophen

500 mg  
Pain Reliever / Fever Reducer  
For Adults



50 CAPLETS\* 500 mg EACH  
(\*\*CAPSULE-SHAPED TABLETS)

NDC 11673-838-05

**RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**  
**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Uses** ■ temporarily relieves minor aches and pains due to:  
■ the common cold ■ headache ■ backache  
■ minor pain of arthritis ■ toothache ■ muscular aches  
■ premenstrual and menstrual cramps  
■ temporarily reduces fever

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
■ more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash ■ if a skin reaction occurs, stop use and seek medical help right away.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease**  
**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if** ■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days ■ new symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** Do not take more than directed. See overdose warning adults and children 12 years and over ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not take for more than 10 days unless directed by a doctor children under 12 years ask a doctor

**Other information** ■ SODIUM FREE ■ store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F) ■ use by expiration date on package

**Questions or comments?** Call 1-800-910-6874  
094.01.0646.R00 Dist. by Target Corp. TM & ©2024 Target Brands, Inc.  
C10282.01.034.000 Mpls., MN 55403  
3-41R.1023

Lot No.:  
Exp. Date:

**UNVARNISHED AREA**

**RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**Drug Facts**  
**Active ingredient (in each caplet) Purpose**  
Acetaminophen 500 mg Pain reliever/fever reducer

**Uses**  
■ temporarily relieves minor aches and pains due to:  
■ the common cold ■ headache  
■ backache ■ minor pain of arthritis  
■ toothache ■ muscular aches  
■ premenstrual and menstrual cramps  
■ temporarily reduces fever

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
■ more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product.  
**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash  
■ if a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have liver disease**  
**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if**  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ new symptoms occur  
■ redness or swelling is present  
These could be signs of a serious condition.

**Drug Facts (continued)**  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children.  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
Do not take more than directed. See overdose warning  
adults and children 12 years and over ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not take for more than 10 days unless directed by a doctor  
children under 12 years ■ ask a doctor

**Other information**  
■ SODIUM FREE  
■ store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)  
■ use by expiration date on package

**Inactive ingredients**  
hydrophilic, mineral oil, povidone, pregelatinized starch, sodium starch glycolate\*, stearic acid, titanium dioxide  
\*may contain this ingredient

**Questions or comments?**  
Call 1-800-910-6874

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Extra Strength Tylenol®.

Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2024 Target Brands, Inc.

Exp. Date:  
Lot No.:  
Coating/Print Omit Area

094 01 001 4 R00  
C-002-282-01-01-034-000  
4948354126

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Compare to active ingredient in Extra Strength Tylenol®  
**Extra Strength Acetaminophen**

500 mg  
Pain Reliever / Fever Reducer

Compare to Active Ingredient in Extra Strength Tylenol®  
**Extra Strength Acetaminophen**

500 mg  
Pain Reliever / Fever Reducer

225 CAPLETS\*\* 500 mg EACH  
(\*\*CAPSULE-SHAPED TABLETS)

Coating Omit

Coating Omit

Coating/Print Omit Area

Actual Size  
225 Caplets

NDC 11673-838-05

**Extra Strength Acetaminophen**  
500 mg  
Pain Reliever / Fever Reducer

NDC 11673-838-26

# Extra Strength Acetaminophen

500 mg  
Pain Reliever / Fever Reducer  
For Adults



225 CAPLETS\*\* 500 mg EACH  
(\*\*CAPSULE-SHAPED TABLETS)

**RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**  
**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each caplet)**  
Acetaminophen 500 mg.....Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - minor pain of arthritis
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** this product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours, or with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash.

If a skin reaction occurs, stop use and seek medical help right away. Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if:**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

 These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** Do not take more than directed. See overdose warning adults and children 12 years and over Take 2 caplets every 6 hours while symptoms last Do not take more than 6 caplets in 24 hours, unless directed by a doctor Do not take for more than 10 days unless directed by a doctor

**Other information**

- SODIUM FREE
- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

**Questions or comments?** Call 1-800-910-6874

34LR 1023  
 Questions or comments? Call 1-800-910-6874  
 Dist. by Target Corp. TM8 02024  
 84001004 R00  
 Mpls., MN 55403  
 ©2014 Target Brands, Inc.  
 C-0026261-004-0000

Lot No.:  
Exp. Date:

**UNVARNISHED AREA**

Compare to active ingredient in Extra Strength Tylenol®†

# Extra Strength Acetaminophen

500 mg  
Pain Reliever / Fever Reducer  
For Adults




500 CAPLETS\* 500 mg EACH  
(\*CAPSULE-SHAPED TABLETS)

NDC 11673-838-50

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each caplet)**  
Acetaminophen 500 mg.....Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - the common cold
  - headache
  - backache
  - minor pain of arthritis
  - toothache
  - muscular aches
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours, or with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash.

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** with any other drug containing

**Drug Facts (continued)**

acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if:**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

 These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions** Do not take more than directed. See overdose warning adults and children 12 years and over Take 2 caplets every 6 hours while symptoms last Do not take more than 6 caplets in 24 hours, unless directed by a doctor Do not take for more than 10 days unless directed by a doctor

**Other information**

- SODIUM FREE
- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

**Inactive ingredients** hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate\*, stearic acid, titanium dioxide

\*may contain this ingredient

**Questions or comments?** Call 1-800-910-6874

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McKee Consumer Healthcare Division, owner of the registered trademark Extra Strength Tylenol®.  
 Dist. by Target Corp. 34LR 1023  
 Mpls., MN 55403  
 ©2014 Target Brands, Inc.  
 C-0026261-004-0000

094 01 0219 R00 C-002282-01-034-0000

Lot No.:  
Exp. Date:

**UNVARNISHED AREA**

ACETAMINOPHEN			
acetaminophen tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-838
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg	
Inactive Ingredients			
Ingredient Name	Strength		
HYPROMELLOSES (UNII: 3NXW29V3WO)			
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)			

<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL341
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-838-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
2	NDC:11673-838-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
3	NDC:11673-838-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
4	NDC:11673-838-26	225 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
5	NDC:11673-838-10	2 in 1 CARTON	01/01/2020	
5	NDC:11673-838-50	500 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/01/2020	

**Labeler** - TARGET CORPORATION (006961700)

**Registrant** - TIME CAP LABORATORIES, INC (037052099)

### Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(11673-838)