

**ASPIRIN- aspirin tablet, coated**  
**TARGET CORPORATION**

-----  
**481R-TARGET - LOW DOSE ASPIRIN 81 MG DELAYED RELEASE 11673-181**

***Drug Facts***

***Active ingredient (in each tablet)***

Aspirin 81 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you

- you have a history of stomach problems, such as heartburn.
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away
- you experience any of the following signs of stomach bleeding:

-feel faint

-vomit blood

-have bloody or black stools

-have stomach pain that does not get better

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### ***Directions***

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

### **OTHER INFORMATION**

#### ***Other information***

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package
- avoid excessive heat above 40°C (104°F)

***Inactive ingredients*** anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

Questions or comments? Call 1-800-910-6874

Adhesive Area

**Drug Facts (continued)**

- take more or for a longer time than directed
- Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients
- Ask a doctor before use if
  - stomach bleeding warning applies to you
  - you have a history of stomach problems, such as heartburn
  - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
  - you are taking a diuretic
  - you have asthma
- Ask a doctor or pharmacist before use if you are taking a prescription drug for
  - gout ■ diabetes ■ arthritis
- Stop using and ask a doctor if
  - an allergic reaction occurs. Seek medical help right away.
  - you experience any of the following signs of stomach bleeding:
    - feel faint ■ vomit/blood ■ have bloody or black stools
    - have stomach pain that does not get better
    - pain gets worse or lasts more than 10 days
    - fever gets worse or lasts more than 3 days
    - redness or swelling is present
    - new symptoms occur
    - ringing in the ears or a loss of hearing occurs
- If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package ■ avoid excessive heat above 40°C (104°F)

**Inactive ingredients** anhydrous indigo, cantharidin wax, colloidal silicon dioxide, croscarmellose sodium, D8C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, poly(oxo)octadecyl, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

**Questions or comments?** Call 1-800-910-6874

Compare to active ingredient in Bayer® Low Dose Aspirin®

# Low Dose Aspirin

81 mg Delayed-Release Tablets  
Pain Reliever (NSAID)  
Safety Coated Aspirin Regimen\*\*




Actual Size

500 ENTERIC-COATED TABLETS

NDC 11673-181-50

\*Talk to your doctor or other healthcare provider before using this product for your heart.  
**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\* ..... Pain reliever  
nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:
 

- hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

**Drug Facts (continued under label)**

This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin.

094 01 0678 800 C-002262-01-034-0000



3 49483 48150 1

Dist. by Target Corporation  
Minneapolis, MN 55403  
TM & ©2024  
Target Brands, Inc.  
481R 1023

Lot No.:  
Exp. Date:


SEE HERE

Varnish Omit Area

Compare to active ingredient in Bayer® Low Dose Aspirin®

# Low Dose Aspirin

81 mg Delayed-Release Tablets  
Pain Reliever (NSAID)  
Safety Coated Aspirin Regimen\*\*



120 ENTERIC-COATED TABLETS

NDC 11673-181-12

\*Talk to your doctor or other healthcare provider before using this product for your heart.  
**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\* ..... Pain reliever  
nonsteroidal anti-inflammatory drug

**Purpose**  
Pain reliever

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction which may include:
 

- hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.

**breast-feeding, ask a health professional before use. Keep out of reach of children.**

**Directions**

■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

**Questions or comments?** Call 1-800-910-6874 481R 1023  
Dist. by Target Corp.  
094 01 0362 R00 Mpls., MN 55403  
C-002262-01-034-0000

Lot No.:  
Exp. Date:

Varnish Omit Area



## ASPIRIN

aspirin tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-181
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

### Inactive Ingredients

#### Drug Facts (continued)

bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ● pain gets worse or lasts more than 10 days ● fever gets worse or lasts more than 3 days ● redness or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs

#### Drug Facts (continued)

**if pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### Directions

■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

#### Other information

■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package ■ avoid excessive heat above 40°C (104°F)

#### Inactive ingredients

anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

**Questions or comments?** call 1-800-910-6874

**Satisfaction guaranteed –  
Love it or your money back.**  
\*\*Talk to your doctor or other healthcare provider before using this product for your heart.  
Distributed by Target Corporation 4891 1023  
Minneapolis, MN 55403  
TM & ©2024 Target Brands, Inc.  
The product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer®.

Compare to active ingredient in Bayer® Low Dose Aspirin®  
**Low Dose Aspirin**

81 mg Delayed-Release Tablets  
Pain Reliever (NSAID)  
Safety Coated Aspirin Regimen\*\*



Actual Size  
120 Tablets

120 ENTERIC-COATED TABLETS

Coating/Print Omit Area

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A</b> (UNII: NX76LV5T8J)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND (bi-convex)	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	HEART;RAISED
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-181-12	1 in 1 CARTON	01/01/2020	
1		120 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11673-181-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/01/2020	

**Labeler** - TARGET CORPORATION (006961700)

**Registrant** - TIME CAP LABORATORIES, INC (037052099)

### Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

TIME CAP LABORATORIES, INC.

037052099

manufacture(11673-181)

Revised: 9/2024

TARGET CORPORATION