### BISACODYL- biscodyl tablet, coated TARGET CORPORATION

-----

#### Target Bisacodyl Tablets 5 mg

### Drug Facts

### Active ingredient (in each tablet)

Bisacodyl USP, 5 mg

### Purpose

Stimulant Laxative

#### Uses

- for relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6-12 hours

#### Warnings

#### WARNINGS

Warning: Do not use if you cannot swallow without chewing.

Ask a doctor before use if you have: stomach pain, nausea or vomiting; A sudden change in bowel habits that lasts more than two weeks.

When using this product: Do not chew or crush tablet(s); Do not use within 1 hour after taking an antacid or milk; Do not use this product if you have stomach discomfort, faintness or cramps.

#### Do not use

• if you cannot swallow without chewing

### Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- noticed a sudden change in bowel habits that lasts more than 2 weeks

#### When using this product

- it may cause stomach discomfort, faintness and cramps
- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk

#### Stop use and ask a doctor if

• you have rectal bleeding or fail to have a bowel movement after using this product.

These could be signs of a serious condition.

• you need to use a laxative for more than 1 week

# If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

# Other information

- each tablet contains: magnesium 5 mg
- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package
- protect from excessive humidity

# Directions

take with a glass of water

- adults and children 12 years of age and over: 1 to 3 tablets in a single daily dose
- children 6 to under 12 years of age: 1 tablet in a single daily dose
- children under 6 years of age: ask a doctor

# Inactive ingredients

acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow (iron oxide ochre), magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide

### **Questions or comments?**

Call **1-800-910-6874** 

Adhesive Area	(PEG) 400, polyrinyl acetate phtmatate, povydone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide Questions or comments? Call 1-800-910-6874	<b>Inactive ingredients</b> acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide microcrystalline cellulose, polyethylene dycol	<ul> <li>Other information</li> <li>each tablet contains: magnesium 5 mg</li> <li>store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)</li> <li>use by expiration date on package</li> <li>protect from excessive humidity</li> </ul>	children 6 to under 12 1 tablet in a single years of age daily dose children under 6 years of age ask a doctor	Directions ■ take with a glass of water adults and children 12 1 to 3 tablets in a years of age and over single daily dose	<ul> <li>Drug Facts (Continued)</li> <li>Stop use and ask a doctor if         <ul> <li>you have rectal beeding or fail to have a bowel movement after using this product. These could be signs of a serious condition.</li> <li>you need to use a laxitive for more than 1 week</li> <li>If pregnant or breast-feeding, ask a health professional before use.</li> <li>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</li> </ul> </li> </ul>
Compare to active ingredient in Dulcolax® Laxative Extra Gentle Bisacodyl Delayed-Release USP, 5 mg Stimulant laxative	TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF The imprinted foil seal over the mouth of the Bottle is cut, torn, broken or missing	Drug Facts Active ingredient Purpose (in each tablet) Bisacodyl USP, 5 mgStimulant laxative Uses	<ul> <li>for relief of occasional constipation and irregularity</li> <li>this product generally produces bowel movement in 6-12 hours</li> <li><i>Warnings</i></li> <li>not use</li> <li>if you cannot swallow without chewing</li> </ul>	Ask a doctor before use if you have ■ stomach pain, nausea or vomiting ■ noticed a sudden change in bowel habits that lasts more than 2 weeks	When using this product The it may cause stomach discomfort, faintness and cramps do not chew or crush tablet(s) do not use within 1 hour after taking an antacid or milk	This product is not manufactured or distributed by Sanofi Consumer This product is not manufactured or distributed by Sanofi Consumer Health care, distributor of Duckar*Laxative. Tages 66.65549 Rd 26.66.5549 Rd 27.000 10.17,1

	Drug Facts (continued) Inactive ingredients eccia, anhydrous calcium: silicon dioxide, corn starch, D&C yellow #10 aluminum iron oxide black, iron oxide yellow (iron oxide ochre), m polyethylene glycol (PEG) 400, polyviny acetate phihala polyethylene glycol (FEG) 400, polyethylene glycol polyethylene glycol (FEG) 400, polytiny acetate phihala polyethylene glycol (FEG) 400, polytiny acetate phihala polytiny acetate phihala polytene glycol (FEG) 400, polytiny acetate phihala polytene glycol (FEG) 400, polytiny acetate phihala polytene glycol (FEG) 400, polytiny acetate phihala polytene glycol (FEG) 400, polytene	ake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, ake, FD&C yellow #6 aluminum lake, gelatin, iron oxide,		
mofi Consumerited thase, distributor of Duts olax* laastive -910-6874 -0000 14R 0324 3 0003557	Warnings Do not use Ask a doctor before use if you have a fryou cannot swallow without chewing Ask a doctor before use if you have a connach pain, nausea or vomiting more than 2 weeks more than 2 weeks do not use within 1 hour after taking an antacid or milk When using this product a do not use within 1 hour after taking an antacid or milk milk milk milk milk milk milk milk	Directions ■ take with a glass of water aduits and children 12 1 to 3 tablets in a years of age and over 12 1 tablet in a single children to under 12 1 tablet in a single years of age daily dose children under 6 years ask a doctor of age each tablet contains: magnesium 5 mg each tablet contains 1 mg each tablet	TAMPER EVIDENT: EACH TABLET IS SAFETY SEALED NOT USE IF BLISTER OR FOIL IS BROKEN OR MISSI	LOT #: EXP. DATE:
otmanufactured or distributedby S <i>or comments</i> ? Call 1-81 D. Target Cor pora tion , MN 554.03 4 Target Brands, Inc.	Active ingredient Purpose (in each tablet) Bisscodyl USP, 5 mgStimulant laxative USes I fris product generatify produces bowel movement in this product generatify produces bowel movement in 6-12 hours	signs of a serious condition. ■ you need to use a laxative for more than 1 week If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, keep out of reach of children. In case of overdose, tweet out of reach of children. In case of overdose, tweet out of reach of children. In case of overdose, tweet out of reach of children. In case of overdose, the set of overdose, the set of the set of the set of overdose, professional professional set of the set of overdose, the set of the set	IS SAFETY SEALED. BROKEN OR MISSIN	Varnish Omit Are



BISACODYL					
biscodyl tablet, coated					
Product Information					
Product Type	HUMAN OTC DRUG	Item Code (So	urce)	NDC:116	73-182
Route of Administration	ORAL				
Active Ingredient/Active	Moiety				
Ingredient Name Basis of Strength Streng					Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)			BISACODYL		5 mg

Ingredient Name	Strength
ACACIA (UNII: 5C5403N260)	
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: 08232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
POVIDONE (UNII: FZ989GH94E)	
GELATIN (UNII: 2G86QN327L)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYVINYL ACETATE PHTHALATE (UNII: 58QVG85GW3)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SHELLAC (UNII: 46N107B71O)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	

Product Characteristics						
Color	orange	Score	no score			
Shape	ROUND	Size	5mm			
Flavor		Imprint Code	TCL003			
Contains						

_	-	
Dar	kan	Ind
Pac	Ray	JIIIG

#	ltem Code	Package Description	Marketing Start Date	Marketing End Date		
		25 in 1 BLISTER PACK; Type 0: Not a Combination Product	02/04/2020			
2	NDC:11673- 182-01         100 in 1 BOTTLE; Type 0: Not a Combination Product         06/15/2024					
Marketing Information						
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
ОТ	C Monograph Dru	g M007	02/04/2020			

# **Registrant -** TIME CAP LABORATORIES, INC (037052099)

Establishment					
Name	Address	ID/FEI	<b>Business Operations</b>		
TIME CAP LABORATORIES, INC.		037052099	manufacture(11673-182)		

Revised: 7/2024

TARGET CORPORATION