

**TUSSIN CF NON DROWSY MULTI SYMPTOM- dextromethorphan hbr,  
guaifenesin, phenylephrine hcl liquid  
WinCo Foods, LLC**

-----

**Drug Facts**

**Active ingredients (in each 10 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 200 mg

Phenylephrine HCl 10 mg

**Purposes**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm ( mucus)
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

taking any other oral nasal decongestant or stimulant.

**When using this product,  
do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1800-222-122) right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adults product is not intended for use in children under 12 years of age
- adult and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

**Other information**

- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

**Questions?**

Call **800-824-1706** Monday-Friday 9am-4pm MST

**Principal Display Panel**

\*Compare to the active ingredients in Robitussin® Peak Cold Multi-Symptom Cold CF

NON-DROWSY

**Tussin CF**

Multi-Symptom

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

GUAIFENESIN / EXPECTORANT

PHENYLEPHRINE HCl / NASAL DECONGESTANT

Relieves: Cough, Stuffy Nose, Chest Congestion & Mucus

Adult Formula, For Ages 12 & Over

Alcohol Free

FL OZ (mL)

\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributors of Robitussin® Peak Cold Multi-Symptom Cold CF.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY: WINCO FOODS, LLC, BOISE, ID 83704

[www.wincofoods.com](http://www.wincofoods.com)

**Package Label**

**Drug Facts (continued)**

**Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

**Other information**

- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

**Questions?** Call 800-824-1706  
Monday - Friday 9am - 4pm MST

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**DISTRIBUTED BY:** WINCO FOODS, LLC, BOISE, ID 83704

\*This product is not manufactured or distributed by PFIZER CONSUMER HEALTHCARE, distributor of ROBITUSSIN® Peak Cold Multi-Symptom Cold CF.

The Supermarket Low Price Leader®

**100% SATISFACTION GUARANTEE**

Comments & Questions?  
800-824-1706 - www.wincofoods.com



PLEASE RECYCLE

\*Compare to the active ingredients in ROBITUSSIN®  
Peak Cold Multi-Symptom Cold CF



DEXTROMETHORPHAN HBr/  
COUGH SUPPRESSANT  
GUAIFENESIN/EXPECTORANT  
PHENYLEPHRINE HCl/  
NASAL DECONGESTANT

Relieves: Cough, Stuffy Nose,  
Chest Congestion & Mucus

Adult Formula, For Ages 12 & Over

Alcohol Free



4 FL OZ (118 mL)

\*Compare to the active ingredients in ROBITUSSIN®  
Peak Cold Multi-Symptom Cold CF



DEXTROMETHORPHAN HBr/  
COUGH SUPPRESSANT  
GUAIFENESIN/EXPECTORANT  
PHENYLEPHRINE HCl/  
NASAL DECONGESTANT

Relieves: Cough, Stuffy Nose,  
Chest Congestion & Mucus

Adult Formula, For Ages 12 & Over

Alcohol Free



4 FL OZ (118 mL)

**Drug Facts**

Active ingredients (in each 10 mL)	Purposes
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold
  - nasal congestion
  - cough due to minor throat and bronchial irritation

**Warnings**

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.**

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.



FC004101 PLD-0322C

Lot No.:

Exp. Date:

**WINCO FOODS Multi Symptom Tussin CF**

## TUSSIN CF NON DROWSY MULTI SYMPTOM

dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67091-266
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL

<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL
---	-----------------------------	-------------------

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>LACTIC ACID</b> (UNII: 33X04XA5AT)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-266-18	1 in 1 BOX	12/31/2014	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/31/2014	

**Labeler** - WinCo Foods, LLC (056098817)

Revised: 5/2024

WinCo Foods, LLC