

ANTI-DIARRHEAL- loperamide hcl tablet
H E B

HEB 44-375

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use

if you have bloody or black stool.

Ask a doctor before use if you have

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product,

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store between 20°-25°C (68°-77°F)
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

Compare to *Imodium® A-D* active ingredient*

NDC 37808-375-02

H-E-B®

Anti-Diarrheal

Loperamide Hydrochloride 2 mg
Anti-Diarrheal

Diarrhea Relief

Controls The Symptoms of Diarrhea,
Including Travelers' Diarrhea

24 CAPLETS

Each caplet contains 2 mg loperamide hydrochloride

actual
size

1. To open, tear
at perforations and
remove a section.

2. Use □ to cut
through plastic
for caplet.

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the
registered trademark Imodium® A-D.

50844 REV0619A37508

**MADE WITH PRIDE AND CARE FOR
H-E-B®, SAN ANTONIO, TX 78204**

H-E-B®

100%

GUARANTEE

promise

If you aren't completely pleased
with this product, we'll be happy to
replace it or refund your money.

You have our word on it.

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark "Product" A-0.

Drug Facts (continued)
 ■ store between 20°-25°C (68°-77°F) ■ see end flap for expiration date and lot number
Inactive ingredients corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dryhydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide
Questions or comments? Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Other information
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool but no more than 2 caplets in 24 hours
children 2-5 years (24-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
 ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

Warnings
 Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.
 Heart alert: Taking more than directed can cause serious heart problems or death.
 Do not use if you have bloody or black stool.
 Ask a doctor before use if you have ■ a fever ■ mucus in the stool ■ a history of liver disease
 ■ a history of abnormal heart rhythm
 Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

Use controls symptoms of diarrhea, including travelers' diarrhea
 Loperamide HCl 2 mg Anti-diarrheal

Drug Facts
Active ingredient (in each caplet)
 Loperamide HCl 2 mg Anti-diarrheal

Purpose
 Loperamide HCl 2 mg Anti-diarrheal

Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
 ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.
 ■ pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.
 Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days
 ■ you get abdominal swelling or bloating. These may be signs of a serious condition.
 When using this product, dizziness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.
 Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

Other information
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN



1. To open, tear at perforations and remove a section.
 2. Use $\frac{1}{2}$ to cut through plastic for caplet.

MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TX 78204

100% GUARANTEE
 If you aren't completely pleased with this product, we'll be happy to replace it or refund your money.
 You have our word on it.



Anti-Diarrheal

Loperamide Hydrochloride 2 mg/Anti-Diarrheal

Diarrhea Relief

Compare to *Imodium® A-D* active ingredient*

NDC 37808-375-08



Anti-Diarrheal

Loperamide Hydrochloride 2 mg


B-0712-375-08-R
 REV0619A37508

No print/No varnish
 Lot & Exp date

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

50844 REV0619A37508 13564-1911

0 41220 54777 7



44-375

ANTI-DIARRHEAL

loperamide hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-375
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	44;375
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-375-02	2 in 1 CARTON	05/03/2005	02/18/2022
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:37808-375-08	4 in 1 CARTON	05/03/2005	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:37808-375-22	1 in 1 CARTON	05/03/2005	04/30/2020
3		48 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	05/03/2005	

Labeler - HEB (007924756)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(37808-375)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(37808-375)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(37808-375)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(37808-375)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(37808-375)