**ACETAMINOPHEN- acetaminophen tablet, film coated, extended release**
**WALGREENS**

---

**8-Hour**
**Pain Reliever**
**ACETAMINOPHEN EXTENDED-RELEASE TABLETS USP, 650 mg**
**PAIN RELIEVER / FEVER REDUCER**

**- Relieves Minor Muscle Pain for up to 8 hours**

**CONTAINS NO ASPIRIN**

## Active ingredient

(in each caplet)
Acetaminophen USP, 650 mg

## Purpose

Pain reliever/fever reducer

## Uses

■ temporarily relieves minor aches and pains due to:

■ muscular aches

■ backache

■ minor pain of arthritis

■ toothache

■ premenstrual and menstrual cramps

■ headache

■ the common cold

■ temporarily reduces fever

## Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

■ more than 6 caplets in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

**Allergy alert**

acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening

■ blisters

■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have**

liver disease

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

■ pain gets worse or lasts more than 10 days

■ fever gets worse or lasts more than 3 days

■ new symptoms occur

■ redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding**

ask a health professional before use.

**Keep out of reach of children**

Keep out of reach of children

**Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

■ d **o not take more than directed (see overdose warning)**

adults and children 12 years of age and over

■ take 2 caplets every 8 hours with water

■ swallow whole; do not crush, chew, split or dissolve

■ do not take more than 6 caplets in 24 hours

■ do not use for more than 10 days unless directed by a doctor

children under 12 years

■ do not use

**Other information**

■ store between 20-25°C (68-77°F)

■ **do not use if foil inner seal is broken or missing**

**Inactive ingredients**

hydroxy ethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

**Questions or comments?**

Contact **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST.




To simulate a printed label, fold along dotted line.

Inside (adhesive side)




COATING FREE AREA
200000005025
700000001958
Drug Facts (continued)
Ingredients in this product
Ask a doctor before use if you have liver disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
Stop use and ask a doctor if
pain gets worse or lasts more than 10 days
fever gets worse or lasts more than 3 days
new symptoms occur
redness or swelling is present
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
do not take more than directed (see overdose warning)
adults and children 12 years of age and over
take 2 caplets every 8 hours with water
swallow whole; do not crush, chew, split or dissolve
do not take more than 6 caplets in 24 hours
do not use for more than 10 days unless directed by a doctor
children under 12 years
do not use
Other information
store between 20-25°C (68-77°F)
Drug Facts (continued)
do not use if bottle inner seal is broken or missing
Inactive ingredients
hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid.
Questions or comments?
contact 1-877-778-0783 Mon-Fri 8:00 AM EST to 9:00 PM PST
*Contains less than 20ppm of gluten
†We invite you to compare to national brands. This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Tylenol®.
NDC 0363-0982-24
PAPER BOX
PLASTIC BOTTLE
DISTRIBUTED BY:
WALGREEN CO.
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2023 Walgreen Co.
MADE IN INDIA
REV-0324
WFPORG0324-F
200000005025
700000001958
COATING FREE AREA
Walgreens free+pure
8-Hour Pain Reliever
Acetaminophen
extended-release tablets USP 650 mg,
Pain reliever / fever reducer
Walgreens free+pure
8-Hour Pain Reliever
Acetaminophen extended-release tablets USP 650 mg,
pain reliever / fever reducer
Compare to the active ingredient in Tylenol® 8 HR Muscle Aches & Pain†
8 hour | Muscle aches & pain
For up to 8 hours relief of minor muscle aches & pain
†Free + Pure is:
• No artificial flavors
• No synthetic colors
• Gluten free*
• Non-GMO
CONTAINS NO ASPIRIN
Actual size
24 caplets*
(*capsule-shaped bi-layer tablets)
ITEM 442621 W10383-0324-L
3 11917 21079 7
Lot
Exp.
COATING FREE AREA
IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING. KEEP THE CARTON FOR IMPORTANT INFORMATION.
DO NOT USE IF FOIL INNER SEAL IS BROKEN OR MISSING
Drug Facts
Active ingredient (in each caplet) Purpose
Acetaminophen USP 650 mg ..... Pain reliever/fever reducer
Uses
temporarily relieves minor aches and pains due to:
muscular aches
backache
minor pain of arthritis
toothache
premenstrual and menstrual cramps
headache
the common cold
temporarily reduces fever
Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
more than 6 caplets in 24 hours, which is the maximum daily amount
with other drugs containing acetaminophen
3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
skin reddening
blisters
rash
If a skin reaction occurs, stop use and seek medical help right away.
Do not use
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
if you are allergic to acetaminophen or any of the inactive




# ACETAMINOPHEN

acetaminophen tablet, film coated, extended release

| Product Information | | | |
|---|---|---|---|
| **Product Type** | HUMAN OTC DRUG | **Item Code (Source)** | NDC:0363-9992 |
| **Route of Administration** | ORAL | | |

| Active Ingredient/Active Moiety | | |
|---|---|---|
| **Ingredient Name** | **Basis of Strength** | **Strength** |
| **ACETAMINOPHEN** (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 650 mg |

| Inactive Ingredients | |
|---|---|
| **Ingredient Name** | **Strength** |

| | |
|---|---|
| **HYPROMELLOSE 2910 (6 MPA.S)** (UNII: 0WZ8WG20P6) | |
| **HYDROXYETHYL CELLULOSE (140 CPS AT 5%)** (UNII: 8136Y38GY5) | |
| **MICROCRYSTALLINE CELLULOSE** (UNII: OP1R32D61U) | |
| **STARCH, CORN** (UNII: O8232NY3SJ) | |
| **SODIUM STARCH GLYCOLATE TYPE A CORN** (UNII: AG9B65PV6B) | |
| **STEARIC ACID** (UNII: 4ELV7Z65AP) | |
| **MAGNESIUM STEARATE** (UNII: 70097M6I30) | |
| **POLYETHYLENE GLYCOL 400** (UNII: B697894SGQ) | |
| **POVIDONE K30** (UNII: U725QWY32X) | |

## Product Characteristics

| | | | |
|---|---|---|---|
| **Color** | white (White to off white colored) | **Score** | no score |
| **Shape** | OVAL (Capsule shaped, biconvex intact film coated tablets) | **Size** | 19mm |
| **Flavor** | | **Imprint Code** | G;650 |
| **Contains** | | | |

## Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:0363-9992-24 | 24 in 1 BOTTLE; Type 0: Not a Combination Product | 09/28/2019 | |
| **2** | NDC:0363-9992-20 | 200 in 1 BOTTLE; Type 0: Not a Combination Product | 09/28/2019 | |
| **3** | NDC:0363-9992-10 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 09/28/2019 | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| ANDA | ANDA211544 | 09/28/2019 | |

## Labeler - WALGREENS (008965063)

Revised: 4/2024 WALGREENS