

**IBUPROFEN- ibuprofen tablet, coated**  
**RETAIL BUSINESS SERVICES , LLC.**

-----  
**1110C-CAR-2025-1118**

***Drug Facts***

**Active ingredient (in each caplet)**

Ibuprofen 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance

is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Heart attack and stroke warning**

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain

- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and older:
  - take 1 caplet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 caplet, 2 caplets may be used
  - do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

## **Other information**

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

## **Inactive ingredients**

colloidal silicon dioxide, corn starch, iron oxide red, polyethylene glycol, polyvinyl alcohol, povidone K30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

## **Questions or comments?**

1-844-705-4384

**PRINCIPAL DISPLAY PANEL**

CAREONE®

NDC 72476-110-02

†Compare to the active ingredient in Advil® Ibuprofen Caplets

IBUPROFEN

Tablets, 200 mg

Pain Reliever/

Fever Reducer (NSAID)

Actual Size

50 CAPLETS\*\*

\*\*Capsule-Shaped Tablets

INK AND COATING FREE  
FOR LOT AND  
EXPIRATION STAMPING

3 41520 31050 8  
T-0007 1025 F1110C02CAR\_R1

DO NOT USE IF IMPRINTED SEAL  
UNDER CAP IS BROKEN OR MISSING

This product is not manufactured or distributed by  
GSK CH, distributor of Advil®

**Drug Facts**

**Active ingredient (in each caplet)** **Purposes**  
Ibuprofen 200 mg (NSAID) — Pain reliever/fever reducer  
Nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to:  
■ muscular aches ■ headache  
■ toothache ■ backache  
■ the common cold ■ menstrual cramps  
■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheezing)  
■ shock ■ skin redness ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs combining prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack.

**Drug Facts (continued)**

heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery  
**Ask a doctor before use if** ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug  
**When using this product** ■ take with food or milk if stomach upset occurs ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke:  
■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area ■ any new symptoms appear

**Drug Facts (continued)**  
**Questions or comments?**  
1-844-705-4384

**Important:** Read all product information before using. Keep this box for important information.

DISTRIBUTED BY: ADVISA DISTRIBUTION, LLC  
SALISBURY, NC 28147  
For product questions or concerns, contact us at 1-833-992-3872.  
Quality guaranteed or your money back.  
Made in India.

RECYCLED PAPER BOX  
RECYCLED PAPER CAPSULES

**Drug Facts (continued)**

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions** ■ do not take more than directed ■ the smallest effective dose should be used  
adults and children 12 years and older  
■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor  
children under 12 years ■ ask a doctor

**Other information** ■ read all warnings and directions before use ■ store between 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

**Inactive ingredients** colloidal silicon dioxide, corn starch, iron oxide red, polyethylene glycol, polyvinyl alcohol, polydioxane K30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

NDC 72476-110-02  
*'Compare to the active ingredient in Advil®'*

**CAREone®**

**IBUPROFEN**  
Tablets, 200 mg  
Pain Reliever/  
Fever Reducer (NSAID)

**50 COATED CAPLETS\*\***  
\*\*Capsule-Shaped Tablets

Actual Size

COATING FREE AREA

<b>IBUPROFEN</b> ibuprofen tablet, coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72476-110
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>STARCH, PREGELATINIZED CORN</b> (UNII: O8232NY3SJ)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	15mm
<b>Flavor</b>		<b>Imprint Code</b>	G2
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:72476-110-02	1 in 1 CARTON	11/01/2020	
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:72476-110-03	1 in 1 CARTON	02/01/2020	
2		100 in 1 PACKAGE; Type 0: Not a Combination Product		
3	NDC:72476-110-04	1 in 1 CARTON	02/01/2020	
3		250 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:72476-110-05	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/01/2020	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA079174	02/01/2020	

---

**Labeler** - RETAIL BUSINESS SERVICES , LLC. (967989935)

Revised: 11/2025

RETAIL BUSINESS SERVICES , LLC.