

PAIN TERMINATOR EXTRA STRENGTH PAIN RELIEF - caffeine, acetaminophen, aspirin capsule

Wildman Business Group

ACTIVE INGREDIENT

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin (NSAID*) 250 mg

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

PURPOSE

Purpose

Pain reliever/fever reducer

Pain reliever/fever reducer

Pain reliever aid

INDICATIONS & USAGE

Uses

temporarily relieves minor aches and pains due to

- headache
- muscular aches
- arthritis pain
- a cold
- toothache
- premenstrual and menstrual cramps

WARNINGS

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin (NSAIDs) may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat

DO NOT USE

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you ever have had an allergic reaction to acetaminophen, aspirin or any other pain relievers/ fever reducers

ASK DOCTOR

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have a history of stomach problems such as heartburn
- you have asthma
- you have liver disease

ASK DOCTOR/PHARMACIST

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for gout, diabetes, or arthritis
- taking any other drug, or under a doctor's care for any serious condition

STOP USE

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- any new symptoms appear

These could be signs of a serious condition.

PREGNANCY OR BREAST FEEDING

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

KEEP OUT OF REACH OF CHILDREN

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

DOSAGE & ADMINISTRATION

Directions

- **do not use more than directed**
- drink a full glass of water with each dose

Adults and children: (12 years and over) Take 2 caplets every 6 hours as needed; not more than 8 caplets in 24 hours

Children under 12 years: Ask a doctor

OTHER SAFETY INFORMATION

Other information

- read all product information before using. Keep box for important information.
- store at room temperature 68°-77°F (20°-25°C)

- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets
- **each caplet contains: Magnesium 0.58 mg**

INACTIVE INGREDIENT

Inactive ingredients

colloidal silicon dioxide, hypromellose, microcrystalline cellulose, polyvinyl pyrrolidone, polyethylene glycol, pregelatinized starch, starch 1500, stearic acid, talc

QUESTIONS

Questions or comments? (888) 602-0288

Principal Display Panel - 100 Caplet Box Label

2533

PAIN TERMINATOR

Advanced Severe Pain Formula

Pull to Open

Fast Relief of:

- Headaches
- Muscle Aches
- Back Aches
- Arthritis Pain

The Provision

First Aid

Line™

BY WILDMAN

100 Caplets

2 Per Packet



- Fast Relief of:**
- Headaches
 - Muscle Aches
 - Back Aches
 - Arthritis Pain



PAIN TERMINATOR

Advanced Severe Pain Formula

2533

2533

2533

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 250 mg	Pain reliever/fever reducer
Aspirin (NSAID*) 250 mg	Pain reliever/fever reducer
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains due to

- headache
- muscular aches
- arthritis pain
- a cold
- toothache
- premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin (NSAIDs) may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you ever have had an allergic reaction to acetaminophen, aspirin or any other pain relievers/fever reducers

(continued on opposite panel)

PAIN TERMINATOR

Advanced Severe Pain Formula

Acetaminophen 250 mg
Aspirin (NSAID) 250 mg
Caffeine 65 mg

Pull to Open



Fast Relief of:

- Headaches
- Muscle Aches
- Back Aches
- Arthritis Pain



Retain carton for complete product information

100 Caplets

2 Per Packet

Drug Facts (continued)**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you are taking a diuretic
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have a history of stomach problems such as heartburn
- you have liver disease

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for gout, diabetes, or arthritis
- taking any other drug, or under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
 - you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - ringing in the ears or loss of hearing occurs
 - painful area is red or swollen
 - any new symptoms appear
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- drink a full glass of water with each dose

Adults and children: (12 years and over) Take 2 caplets every 6 hours as needed; not more than 8 caplets in 24 hours

Children under 12 years: Ask a doctor

Other information

- read all product information before using. Keep box for important information.
- store at room temperature 68°-77°F (20°-25°C)
- tamper evident sealed packets
- each caplet contains: Magnesium 0.58 mg
- avoid excessive heat and humidity
- do not use any opened or torn packets

Inactive ingredients

colloidal silicon dioxide, hypromellose, microcrystalline cellulose, polyvinyl pyrrolidone, polyethylene glycol, pregelatinized starch, starch 1500, stearic acid, talc

Questions or comments? (888) 602-0288 Manufactured for: Provision by Wildman, Warsaw, IN 46580

PAIN TERMINATOR

Advanced Severe Pain Formula

Acetaminophen 250 mg
Aspirin (NSAID) 250 mg
Caffeine 65 mg

**Fast Relief of:**

- Headaches
- Muscle Aches
- Back Aches
- Arthritis Pain



Retain carton for complete product information

100 Caplets

2 Per Packet

Manufactured for Provision by Wildman
Warsaw, IN 46580



PAIN TERMINATOR EXTRA STRENGTH PAIN RELIEF

caffeine, acetaminophen, aspirin capsule

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84269-2533
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	

Product Characteristics

Color	white (white to off white with specks)	Score	no score
Shape	CAPSULE (capsule shaped, biconvex)	Size	18mm
Flavor		Imprint Code	S53
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84269-2533-2	100 in 1 BOX	07/01/2024	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:84269-2533-1	50 in 1 BOX	07/01/2024	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/01/2024	

Labeler - Wildman Business Group (016677338)

Establishment

Name	Address	ID/FEI	Business Operations
Elysium Pharmaceuticals LTD.		863182240	MANUFACTURE(84269-2533)

Establishment

Name	Address	ID/FEI	Business Operations
Medique Products		086911794	PACK(84269-2533)

Establishment

Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	PACK(84269-2533)

Revised: 7/2024

Wildman Business Group