

## **ASPIRIN LOW DOSE- aspirin tablet, chewable**

### **Chain Drug Consortium**

-----  
**Premier Value 44-218**

#### ***Active ingredient (in each chewable tablet)***

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

#### ***Purpose***

Pain reliever

#### ***Uses***

for the temporary relief of minor aches and pains or as recommended by your doctor

#### ***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems

#### **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, dextrans hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

### ***Questions or comments?***

**1-800-426-9391**

### ***Principal Display Panel***

#### **VALUE PACK**

108 Tablets

(3 bottles of 36 tablets each)

†COMPARE TO THE ACTIVE INGREDIENT IN  
BAYER® CHEWABLE LOW DOSE ASPIRIN

#### ***Premier Value®***

#### ***Low Dose***

***Aspirin 81 mg***

***Chewable Tablets***

**PAIN RELIEVER (NSAID)**

Aspirin regimen\*\*

actual  
size

**108** Orange flavored  
**Chewable Tablets**

**Chew or crush tablets completely  
before swallowing.**

**\*\*Talk to your doctor or other healthcare provider  
before using this product for your heart.**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY  
SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer AG, owner of the registered  
trademark Bayer®

Chewable Low Dose Aspirin.      50844      REV0122B21807

Distributed By:  
Pharmacy Value Alliance, LLC  
407 East Lancaster Avenue  
Wayne, PA 19087

**If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.**

**Drug Facts**

**Active ingredient (in each chewable tablet)**  
Aspirin 81 mg (NSAID)  
Pain reliever

**Purpose**  
Nonsteroidal anti-inflammatory drug

**Uses**  
For the temporary relief of minor aches and pains or as recommended by your doctor

**Warnings**  
Check on and be aware of anyone who has or is recovering from chicken pox or the like. Aspirin should not be used in children because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**VALUE PACK**  
108 Tablets  
(3 bottles of 36 tablets each)



**Low Dose Aspirin 81 mg Chewable Tablets**  
**PAIN RELIEVER (NSAID)**

Aspirin regimen\*\*

**108 Orange flavored Chewable Tablets**  
Chew or crush tablets completely before swallowing.

**COMPARE TO THE ACTIVE INGREDIENT IN BAYER® CHEWABLE LOW DOSE ASPIRIN**

**108 Orange flavored Chewable Tablets**  
Chew or crush tablets completely before swallowing.

**Drug Facts (continued)**

**Inactive ingredients**  
as in strength, does not include 750 mg of aspirin.

**Questions or comments?**  
1-800-428-9381

**Other Information**  
Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F). See end flap for expiration date and lot number.

**Directions**  
Do not take more than directed.

Drink a full glass of water with each dose.

Chew or crush tablets completely before swallowing. Do not swallow tablets whole.

Adults and children 12 years and over: take 4 to 6 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor.

Children under 12 years: do not use unless directed by a doctor.

**Disturbed By:**  
Pharmacy Value Alliance, LLC  
407 East Lancaster Avenue  
Wayne, PA 19087

**Aspirin 81 mg Chewable Tablets**  
**PAIN RELIEVER (NSAID)**

**108 Orange flavored Chewable Tablets**  
Chew or crush tablets completely before swallowing.

**108 Orange flavored Chewable Tablets**  
Chew or crush tablets completely before swallowing.

**B-1590-218A-07-TB**  
**REV0122821807**

**no print / no varnish area**  
lot no. & exp. date

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*\*Talk to your doctor or other healthcare provider before using this product for your heart.



8 40986 03221 9

**Drug Facts (continued)**

**Warning:** Aspirin may cause a severe allergic reaction, which may include hives, facial swelling, shock, asthma (wheezing), stomach bleeding warning. This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you take blood thinning (anticoagulant) or steroid drug, are age 60 or older, take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others), have 2 or more alcohol drinks every day while using this product, have had stomach ulcers or bleeding problems, take more or for a longer time than directed.

**Directions:** Take as directed. Do not use if you are allergic to aspirin or any other pain reliever. Review product information for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor.

**Other important information:** If you have ever had an allergic reaction to this product or any of its ingredients, ask a doctor before use. If you have a stomach bleeding warning apply to you, ask a doctor before use. If you have a history of stomach problems, such as heartburn, you have a high blood pressure, heart disease, liver disease, or kidney disease, you have asthma, you are taking a diuretic, you have a history of stomach ulcers or bleeding problems, you are taking a prescription drug for gout, diabetes, or HIV, ask a doctor or pharmacist before use. If you are taking a prescription drug for any of these conditions, ask a doctor before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless directed to do so by a doctor.

**Other information:** These could be signs of a serious condition: new symptoms occur, redness or swelling is present, fever gets worse or lasts more than 3 days, pain gets worse or lasts more than 10 days, ringing in the ears or a loss of hearing occurs, you experience any of the following signs of stomach bleeding: vomit blood, have bloody or black stools, feel faint, have stomach pain that does not get better.

**Keep this and other medicine out of the reach of children.**

## Premier Value 44-218

### ASPIRIN LOW DOSE

aspirin tablet, chewable

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68016-690
<b>Route of Administration</b>	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

#### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

#### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	44;218
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-690-36	1 in 1 CARTON	09/12/1990	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:68016-690-18	3 in 1 CARTON	09/12/1990	
2		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/12/1990	

**Labeler** - Chain Drug Consortium (101668460)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(68016-690)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(68016-690)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(68016-690)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(68016-690)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(68016-690)

Revised: 9/2024

Chain Drug Consortium