

**VICKS DAYQUIL SEVERE AND VICKS NYQUIL SEVERE COLD AND FLU-
acetaminophen, phenylephrine hydrochloride, dextromethorphan
hydrobromide, guaifenesin and doxylamine succinate
The Procter & Gamble Manufacturing Company**

**Vicks ® DayQuil™ Severe and Vicks ® NyQuil™ Severe Cold & Flu LiquiCaps™
Convenience Pack**

DayQuil™ SEVERE Cold & Flu

Drug Facts

Active ingredients (in each LiquiCap)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/Fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning:

This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 8 LiquiCaps in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough get worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 LiquiCaps per 24 hrs

adults & children 12 yrs & over	2 LiquiCaps with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- do not exceed 25°C

Inactive ingredients

ascorbic acid, FD&C Blue no. 1, FD&C Red No. 40, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, shellac, simethicone, sorbitan, sorbitol, titanium dioxide, water

Questions?

1-800-362-1683

NyQuil™ SEVERE Cold & Flu

Drug Facts

Active ingredients (in each LiquiCap)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Doxylamine succinate 6.25 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/Fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 LiquiCaps in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain

drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, & tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 LiquiCaps per 24 hours

adults & children 12 yrs & over	2 LiquiCaps with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- do not exceed 25°C

Inactive ingredients

ascorbic acid, FD&C Blue No. 1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, shellac, simethicone, sorbitan, sorbitol, titanium dioxide, water

Questions?

1-800-362-1683

TAMPER EVIDENT: This package is safety-sealed & child resistant. Use only if blisters are intact. If difficult to open, use scissors.

Made in Canada

DIST. BY PROCTER & GAMBLE, CINCINNATI OH 45202

PRINCIPAL DISPLAY PANEL - 24 LIQUICAPS Convenience Pack

MAX

STRENGTH

Vicks ®DayQuil™ Severe and Vicks ®NyQuil™ Severe Cold & Flu LiquiCaps™

VICKS®

DayQuil™

SEVERE

COLD& FLU

Acetaminophen, Guaifenesin, Phenylephrine HCl, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Chest Congestion, Thins & Loosens Mucus

Nasal Congestion & Sinus Pressure

Cough

Non-Drowsy

8 DayQuil LiquiCaps

NyQuil TM

SEVERE

COLD& FLU

Acetaminophen, PhenylephrineHCl, Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Sneezing, Runny Nose

Cough

Nighttime Relief

16 DAYQUIL LIQUICAPS

8 NYQUIL LIQUICAPS

24 TOTAL LIQUICAPS

Drug Facts (continued) If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	Drug Facts (continued) If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions • take only as directed • do not exceed 8 LiquiCaps per 24 hrs adults & children 12 yrs & over 2 LiquiCaps with water every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use	Directions • take only as directed • do not exceed 8 LiquiCaps per 24 hrs adults & children 12 yrs & over 2 LiquiCaps with water every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use
Other information • store at no greater than 25°C	Other information • store at no greater than 25°C
Inactive ingredients ascorbic acid, FD&C Blue No. 1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, shellac, simethicone, sorbitan, sorbitol, titanium dioxide, water	Inactive ingredients ascorbic acid, FD&C Blue No. 1, FD&C Red No. 40, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, shellac, simethicone, sorbitan, sorbitol, titanium dioxide, water
Questions? 1-800-362-1683	Questions? 1-800-362-1683

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

TAMPER EVIDENT: This package is safety-sealed & child resistant. Use only if blisters are intact. If difficult to open, use scissors.

**MAX
STRENGTH**



DayQuil™

**SEVERE
COLD & FLU**

Acetaminophen, Guaifenesin,
Phenylephrine HCl, Dextromethorphan HBr

Headache, Fever, Sore Throat,
Minor Aches & Pains
Chest Congestion, Thins & Loosens Mucus
Nasal Congestion, Sinus Pressure
Cough

Non-Drowsy

16 DAYQUIL LIQUICAPS

NyQuil™

**SEVERE
COLD & FLU**

Acetaminophen, Phenylephrine HCl,
Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat,
Minor Aches & Pains
Nasal Congestion, Sinus Pressure
Sneezing, Runny Nose
Cough

Nighttime Relief

24 TOTAL
LIQUICAPS™

8 NYQUIL LIQUICAPS:



Made in Canada
DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202

P&G

Patents:
www.pg.com/patents

21328713



Drug Facts	NyQuil™ SEVERE Cold & Flu	Drug Facts	DayQuil™ SEVERE Cold & Flu
Active ingredients (in each LiquiCap)	Purpose	Active ingredients (in each LiquiCap)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer	Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant	Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Expectorant	Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant	Phenylephrine HCl 5 mg	Nasal decongestant
Uses	temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • cough to help you sleep • minor aches & pains • headache • fever • sore throat • runny nose & sneezing • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage	Uses	temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • minor aches & pains • headache • fever • sore throat • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage • helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.
Warnings	Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product	Warnings	Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Drug Facts (continued) Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin When using this product • do not use more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	Drug Facts (continued) which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland Ask a doctor or pharmacist before use if you are • taking the blood thinning drug warfarin When using this product, do not use more than directed. Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
---	--

VICKS DAYQUIL SEVERE AND VICKS NYQUIL SEVERE COLD AND FLU

acetaminophen, phenylephrine hydrochloride, dextromethorphan hydrobromide, guaifenesin and

doxylamine succinate kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37000-934
--------------	----------------	--------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37000-934-12	1 in 1 PACKAGE	07/04/2019	
2	NDC:37000-934-24	1 in 1 PACKAGE	07/04/2019	
3	NDC:37000-934-48	1 in 1 PACKAGE	07/04/2019	
4	NDC:37000-934-72	1 in 1 PACKAGE	07/04/2019	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	8
Part 2	1 BLISTER PACK	4

Part 1 of 2

VICKS DAYQUIL SEVERE COLD AND FLU

acetaminophen, phenylephine hcl, dextromethorphan hydrobromide, and guaifenesin capsule, liquid filled

Product Information

Item Code (Source)	NDC:37000-517
--------------------	---------------

Route of Administration	ORAL
-------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
SHELLAC (UNII: 46N107B71O)	
SORBITOL (UNII: 506T60A25R)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
MANNITOL (UNII: 3OWL53L36A)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	orange	Score	no score
Shape	BULLET	Size	16mm
Flavor		Imprint Code	DS
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 in 1 CARTON		
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/10/2018	

Part 2 of 2

VICKS NYQUIL SEVERE COLD AND FLU

acetaminophen, phenylephrine hydrochloride, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled

Product Information

Item Code (Source)	NDC:37000-518
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	green	Score	no score
Shape	BULLET	Size	16mm
Flavor		Imprint Code	NS
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 in 1 CARTON		
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/10/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/04/2019	

Revised: 8/2025

The Procter & Gamble Manufacturing Company