

**ULINE ACETAMINOPHEN EXTRA STRENGTH- acetaminophen tablet, film coated**  
**Uline**

-----  
**Uline Acetaminophen Extra Strength**

***Drug Facts***

***Active ingredient (in each tablet)***

Acetaminophen 500 mg

***Purpose***

Pain reliever/fever reducer

***Uses***

Temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- minor arthritis pain
- backache
- the common cold
- toothache
- premenstrual and menstrual cramps

Temporarily reduces fever.

***Warnings***

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug

contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have**

- liver disease

**Ask a doctor or pharmacist before use if you have**

- liver disease

**Stop use and ask a doctor if**

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- **do not use more than directed (see overdose warning)**

**Adults and children: (12 years and over)**

Take 2 tablets every 6 while symptoms last. Do not take more than 6 tablets in 24 hours, unless directed by a doctor. Do not use for more than 10 days unless directed by a doctor.

**Children under 12 years:**

Ask a doctor.

***Other information***

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

### Inactive ingredients

corn starch\*, hypromellose\*, polyethylene glycol\*, povidone (K-30)\*, pregelatinized starch\*, purified water\*, sodium starch glycolate, stearic acid, titanium dioxide\*

\* May contain

### Questions or comments?

1-800-295-5510

### Uline Acetaminophen Extra Strength Label

ULINE

Tamper evident sealed packets:

Do not use if packet is open or torn.

This package is for households without young children

Acetaminophen

Pull to Open

Extra Strength

500 mg

- Pain Reliever
- Fever Reducer

50 Packets

2 tablets Each

**ACETAMINOPHEN EXTRA STRENGTH 500 mg**

**ULINE**

Tamper evident sealed packets. Do not use if packet is open or torn.

This package is for households without young children

**ULINE**

**ACETAMINOPHEN EXTRA STRENGTH 500 mg**

Pull to Open

• Pain Reliever • Fever Reducer

50 PACKETS (2 TABLETS EACH)

Reorder No. S-18593  
Distributed by ULINE  
13375 Uline Drive  
Recorder, PA, WI 53158  
Rev 11-4-23 - Mfg. 1.24.24

Active ingredient (in each tablet)	Purpose
Acetaminophen 500 mg	Pain reliever/fever reducer

**Uses**  
Temporarily relieves minor aches and pains due to:  
 ■ headache ■ muscular aches ■ minor arthritis pain ■ backache ■ the common cold ■ toothache  
 ■ premenstrual and menstrual cramps  
 Temporarily reduces fever.

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
 ■ more than 4,000 mg in 24 hours, which is the maximum daily amount  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have** ■ liver disease

**Ask a doctor or pharmacist before use if** ■ you are taking the blood thinning drug warfarin

**Stop use and ask a doctor if**  
 ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur  
 ■ redness or swelling is present  
 These could be signs of a serious condition.

**Drug Facts (continued)**  
 If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children.  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children (even if you do not notice any signs or symptoms).

**Directions**  
 ■ do not use more than directed (see overdose warning)  
**Adults and children:** Take 2 tablets every 6 hours while symptoms last. Do not take more than 6 tablets in 24 hours, unless directed by a doctor. Do not use for more than 10 days unless directed by a doctor.  
**Children under 12 years:** Ask a doctor.

**Other information**  
 ■ store at room temperature 59°-86°F (15°-30°C)  
 ■ tamper-evident sealed packets  
 ■ do not use any opened or torn packets

**Inactive ingredients**  
 corn starch\*, hypromellose\*, polyethylene glycol\*, povidone (K-30)\*, pregelatinized starch\*, purified water\*, sodium starch glycolate, stearic acid, titanium dioxide\* \*may contain

**Questions or comments?** 1-800-295-5510

Retain carton for complete product information

### Uline Acetaminophen Extra Strength Label

ULINE

Tamper evident sealed packets:

Do not use if packet is open or torn.

This package is for households without young children

Acetaminophen

Pull to Open

Extra Strength

500 mg

- Pain Reliever
- Fever Reducer

50 Packets

2 tablets Each

**ACETAMINOPHEN**  
EXTRA STRENGTH  
500 mg

**ULINE**

Tamper evident sealed packets. Do not use if packet is open or torn.

This package is for households without young children

**ACETAMINOPHEN**  
EXTRA STRENGTH  
500 mg

Pull to Open

•Pain Reliever •Fever Reducer

50 PACKETS (2 TABLETS EACH)

50 PACKETS (2 TABLETS EACH)

Reorder No. 5-18593  
Distributed by: ULINE  
12875 Uline Drive  
Pleasant Prairie, WI 53158  
Rev 11-6-23 - Mfg. 1.24.44

Active ingredient (in each tablet)	Purpose
Acetaminophen 500 mg	Pain reliever/fever reducer

**Uses**  
Temporarily relieves minor aches and pains due to:  
■ headache ■ muscular aches ■ minor arthritis pain ■ backache ■ the common cold ■ toothache  
■ premenstrual and menstrual cramps  
Temporarily reduces fever.

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
■ more than 4,000 mg in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have** ■ liver disease

**Stop use and ask a doctor if** ■ you are taking the blood thinning drug warfarin  
■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur  
■ redness or swelling is present  
These could be signs of a serious condition.

**Drug Facts (continued)**  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children.  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ do not use more than directed (see overdose warning)

**Adults and children (12 years and over)**  
Take 2 tablets every 6 hours while symptoms last. Do not take more than 8 tablets in 24 hours unless directed by a doctor. Do not use for more than 10 days unless directed by a doctor.

**Children under 12 years:**  
Ask a doctor.

**Other information**  
■ store at room temperature 59°-86°F (15°-30°C)  
■ tamper-evident sealed packets  
■ do not use any opened or torn packets

**Inactive ingredients**  
corn starch, hypromellose, polyethylene glycol, povidone (K-301), pregelatinized starch, purified water, sodium starch glycolate, stearic acid, titanium dioxide. \*may contain

**Questions or comments?** 1-800-295-5510

Retain carton for complete product information

## Uline Acetaminophen Extra Strength Label

ULINE

Tamper evident sealed packets:

Do not use if packet is open or torn.

This package is for households without young children

Acetaminophen

Pull to Open

Extra Strength

500 mg

- Pain Reliever
- Fever Reducer

50 Packets

2 tablets Each



ULINE

Tamper evident sealed packets:

Do not use if packet is open or torn.

This package is for households without young children

Acetaminophen

Pull to Open


Extra Strength

500 mg

- Pain Reliever
- Fever Reducer

50 Packets

2 tablets Each



# ULINE ACETAMINOPHEN EXTRA STRENGTH

acetaminophen tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69790-173
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

## Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	44;148
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69790-173-33	50 in 1 BOX	01/30/2024	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/30/2024	

# ULINE ACETAMINOPHEN EXTRA STRENGTH

acetaminophen tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69790-804
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	AZ;235
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69790-804-33	50 in 1 BOX	10/07/2019	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	10/07/2019	

**ULINE ACETAMINOPHEN EXTRA STRENGTH**

acetaminophen tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69790-126
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	FR;33
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69790-126-33	50 in 1 BOX	10/07/2019	11/01/2023
1		2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/07/2019	11/01/2023

## ULINE ACETAMINOPHEN EXTRA STRENGTH

acetaminophen tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69790-154
---------------------	----------------	---------------------------	---------------

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE K30 (UNII: U725QWY32X)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

### Product Characteristics

<b>Color</b>	white ((White to Off-White))	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	G552
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69790-154-33	50 in 1 BOX	01/30/2024	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/30/2024	

**Labeler** - Uline (039612668)

**Registrant** - Unifirst First Aid Corporation (832947092)