

QUALITY CHOICE EFFERVESCENT COLD RELIEF PLUS- aspirin, chlorpheniramine maleate, and phenylephrine bitartrate tablet, effervescent
Chain Drug Marketing Association

Effervescent Cold Relief Tablets

Drug Facts

Active ingredients (in each effervescent tablet)	Purpose
Aspirin 325 mg (NSAID)*.	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Phenylephrine Bitartrate 7.8 mg	Nasal decongestant

*Nonsteroidal anti-inflammatory drug

Uses

temporarily relieves these symptoms of the common cold:

- fever ■ sneezing ■ nasal congestion ■ sore throat ■ headache
- minor aches and pains ■ runny nose
- sinus congestion and pressure

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. (See note below)

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are allergic to aspirin

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic. ■ you are taking sedatives or tranquilizers
You have ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

■ presently taking a prescription drug.

■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product

■ do not take more than 8 tablets (adults and children 12 years and over) in a 24-hour period or as directed by doctor.

■ do not use more than directed

■ you may get drowsy

■ avoid alcoholic drinks

■ excitability may occur, especially in children

■ alcohol, sedatives and tranquilizers may increase drowsiness

■ be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding

■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

■ an allergic reaction occurs. Seek medical help right away.

■ pain or nasal congestion gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ new symptoms occur

■ redness or swelling is present

■ ringing in the ears or a loss of hearing occurs

■ nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin during the last 3 months of

pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions


- do not use more than directed
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 ounces of water every 4-6 hours
- do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Inactive ingredients:

acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium

benzoate, sodium bicarbonate

Fast Relief of: Nasal Congestion, Runny Nose, Sneezing, Headache, Sinus Pain & Pressure



Drug Facts
(In each effervescent tablet) Purpose
Combining acetaminophen 325 mg, pseudoephedrine hydrochloride 120 mg, and phenylephrine hydrochloride 10 mg provides fast relief of pain, fever, and nasal congestion.

Warnings
Children and teenagers who have an asthma attack should not use this product. When using this product, it changes in behavior with stress and worry may occur, causing dizziness, a rash, or other symptoms, a rash that causes blisters, or a rash that causes blisters, or a rash that causes blisters.

Directions
Temporarily relieve these symptoms of the common cold: sore throat, nasal congestion, sinus pain and pressure, sinus congestion and pressure.

Other Information
Keep out of reach of children. This product contains a non-aspirin salicylate. Do not use this product if you are taking aspirin or other drugs containing aspirin, or if you are taking any other drug containing aspirin, or if you are taking any other drug containing aspirin, or if you are taking any other drug containing aspirin.

Drug Facts (continued)
Some of the most common side effects are: drowsiness, dizziness, or lightheadedness. If you experience any of these symptoms, stop taking this product and contact your doctor.

Other Information
Keep out of reach of children. This product contains a non-aspirin salicylate. Do not use this product if you are taking aspirin or other drugs containing aspirin, or if you are taking any other drug containing aspirin, or if you are taking any other drug containing aspirin.

OC QUALITY CHOICE
Effervescent Cold Relief Plus Original Flavor

OC QUALITY CHOICE
Effervescent Cold Relief Plus Original Flavor
Aspirin (NSAID) | Pain reliever- fever reducer
Chlorpheniramine Maleate | Antihistamine
Phenylephrine Bitartrate | Nasal Decongestant

Fast Relief of:
Nasal Congestion | Runny Nose | Sneezing | Headache | Sinus Pain & Pressure | Rummy Nose

20 Effervescent Tablets

Drug Facts (continued)
Some of the most common side effects are: drowsiness, dizziness, or lightheadedness. If you experience any of these symptoms, stop taking this product and contact your doctor.

Other Information
Keep out of reach of children. This product contains a non-aspirin salicylate. Do not use this product if you are taking aspirin or other drugs containing aspirin, or if you are taking any other drug containing aspirin, or if you are taking any other drug containing aspirin.

Drug Facts (continued)
Other Information
Keep out of reach of children. This product contains a non-aspirin salicylate. Do not use this product if you are taking aspirin or other drugs containing aspirin, or if you are taking any other drug containing aspirin, or if you are taking any other drug containing aspirin.

Inactive Ingredients
acesulfame potassium, aspartame, citric acid, docusate sodium, mannitol, povidone, sodium bicarbonate, sodium benzoate.

© Distributed by C.D.M.A., Inc.
4237 N.W. 187th Street
Miami Lakes, FL 33054
Questions: 248-449-9300
DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED.

PN:87360 QC C3
REV:01/14

6 35515 93704 6
DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED.

QUALITY CHOICE EFFERVESCENT COLD RELIEF PLUS

aspirin, chlorpheniramine maleate, and phenylephrine bitartrate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-228
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
PHENYLEPHRINE BITARTRATE (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ASPARTAME (UNII: Z0H242BBR1)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
DOCUSATE SODIUM (UNII: F05Q2T2JA0)	
MANNITOL (UNII: 3OWL53L36A)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor	LEMON	Imprint Code	CF
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-228-20	10 in 1 CARTON	01/01/2014	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug M012

01/01/2014

Labeler - Chain Drug Marketing Association (011920774)

Revised: 11/2024

Chain Drug Marketing Association