

ISOXSUPRINE HYDROCHLORIDE- isoxsuprine hydrochloride tablet
Vedco dba Valdar


Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Isosuprine Hydrochloride Tablets, USP

Rx Only

DESCRIPTION

Each tablet taken orally contains Isoxsuprine Hydrochloride, USP with the following chemical structure:



p-Hydroxy- α [1-[(methyl-2-phenoxy-ethyl)amino]ethyl]benzyl alcohol hydrochloride.

QUANTITATIVE INGREDIENT INFORMATION

Each tablet taken orally contains 20mg Isoxsuprine Hydrochloride.

PHARMACOLOGICAL CLASS

Peripheral Vasodilator

INDICATIONS

Based on a review of this drug by the National Academy of Sciences-National Research and/or other information, the FDA has classified the indications as follows:

Possibly Effective

1. For the relief of symptoms associated with cerebrovascular insufficiency.
2. In peripheral vascular disease of arteriosclerosis obliterans, thromboangitis obliterans (Buerger's disease) and Raynaud's disease.

Final classification of the less-than-effective indications requires further investigation.

CONTRAINDICATIONS

There are no known contraindications to oral use when administered in recommended doses.

Isoxsuprine Hydrochloride, USP should not be given immediately postpartum or in the presence of arterial bleeding.

PRECAUTIONS

Pediatric Use

Safety and effectiveness in pediatric patients has not been established.

ADVERSE REACTIONS

On rare occasion oral administration of the drug has been associated in time with the occurrence of hypotension, tachycardia, chest pain, nausea, vomiting, dizziness, abdominal distress, and severe rash. If rash appears, the drug should be discontinued.

Although available evidence suggests a temporal association of these reactions with Isoxsuprine Hydrochloride, a causal relationship can be neither confirmed nor refuted.

Beta Adrenergic receptor stimulants such as Isoxsuprine Hydrochloride have been used to inhibit pre-term labor.

Maternal and fetal tachycardia may occur under such use.

Hypocalcemia, hypoglycemia, hypotension and ileus have been reported to occur in infants whose mothers received Isoxsuprine Hydrochloride. Pulmonary edema has been reported in mothers treated with beta stimulants. Isoxsuprine Hydrochloride is neither approved nor recommended for use in the treatment of premature labor.

DOSAGE AND ADMINISTRATION

Oral: 10 to 20 mg, three or four times daily.

HOW SUPPLIED

Isoxsuprine HCl tablets, USP 20 mg

Bottles of 1000 NDC 63549-919-53

COMPOSITION

Isoxsuprine HCl 20mg tablets: These tablets contain the following inactive ingredients: corn starch, lactose monohydrate, magnesium stearate (vegetable), microcrystalline cellulose.

Manufactured For:

Valdar

St. Joseph, MO 64507

PRINCIPAL DISPLAY PANEL - 20 mg Tablet Bottle Label

NDC 63549-919-53

**Isoxsuprine
Hydrochloride
Tablets**

**USP
20 mg**

1000 Tablets

Rx Only

VALDAR

NDC 63549-919-53

**Isoxsuprine
Hydrochloride
Tablets
USP
20 mg**

**1000 Tablets
Rx Only**

VALDAR

Each tablet contains:
Isoxsuprine Hydrochloride, USP 20 mg

Inactive Ingredients: Corn Starch, Lactose Monohydrate, Magnesium Stearate, Microcrystalline Cellulose.

Usual Dosage: 1 tablet 3 or 4 times daily.
See package insert.

Store at controlled room temperature 15°-30°C (59°-86°F).
Dispense in a tight, light-resistant container.

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

Manufactured for
Valdar, St. Joseph, MO 64507

Iss. 07/2011



Lot No.:
Exp. Date:

ISOXSUPRINE HYDROCHLORIDE

isoxsuprine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63549-919
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
isoxsuprine hydrochloride (UNII: V74TEQ36CO) (Isoxsuprine - UNII:R15UIB245N)	isoxsuprine hydrochloride	20 mg

Inactive Ingredients

Ingredient Name	Strength
Lactose Monohydrate (UNII: EWQ57Q8I5X)	
Magnesium Stearate (UNII: 70097M6I30)	
Cellulose, Microcrystalline (UNII: OP1R32D61U)	
Starch, Corn (UNII: O8232NY3SJ)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	8mm
Flavor		Imprint Code	20
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63549-919-53	1000 in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		08/23/2011	

Labeler - Vedco dba Valdar (021634266)

Revised: 1/2012

Vedco dba Valdar