

**GOOD NEIGHBOR PHARMACY LORATADINE D- loratadine, pseudoephedrine sulfate tablet, film coated, extended release**

**Amerisource Bergen**

-----

**Amerisource Bergen Loratadine-D Drug Facts**

**Active ingredients (in each tablet)**

Loratadine 5 mg

Pseudoephedrine sulfate 120 mg

**Purpose**

Antihistamine

Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- sneezing
- itchy, watery eyes
- runny nose
- itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland

- liver or kidney disease. Your doctor should determine if you need a different dose.

**When using this product do not take more than directed.**

Taking more than directed may cause drowsiness.

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

**Other information**

- **each tablet contains:** calcium 25 mg
- do not use if blister unit is broken or torn
- store between 20° to 25°C (68° to 77°F)
- keep in a dry place

**Inactive ingredients**

croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

**Questions or comments?**

**1-800-719-9260**

**Package/Label Principal Display Panel**

Compare to Claritin-D® 12 Hour

active ingredients

12 HOUR

Loratadine-D

pseudoephedrine sulfate 120 mg/Loratadine 5 mg

extended release tablets

nasal decongestant/antihistamine

Extended Release Tablets

Relief of: Nasal and Sinus Congestion Due to Colds or Allergies

Sneezing

Runny Nose

Itchy, Watery Eyes

Itchy Throat or Nose Due to Allergies

indoor & outdoor allergies

allergy & congestion

Non-Drowsy\*

actual size

\*When taken as directed.

See Drug Facts Panel.

10 tablets

# Loratadine-D

pseudoephedrine sulfate 120 mg/loratadine 5 mg  
extended release tablets  
nasal decongestant/antihistamine

▶ 12 HOUR

**Extended Release Tablets**



Compare to Claritin-D® 12 Hour  
active ingredients\*\*

NDC 46122-455-52

# Loratadine-D

pseudoephedrine sulfate 120 mg/loratadine 5 mg  
extended release tablets  
nasal decongestant/antihistamine

▶ 12 HOUR

**Extended Release Tablets**

Relief of: Nasal and Sinus Congestion Due to Colds or Allergies

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose Due to Allergies

indoor & outdoor allergies  
allergy & congestion

**Non-Drowsy\***

\*When taken as directed.  
See Drug Facts Panel.

actual  
size



10 tablets

## Drug Facts

<b>Active ingredients (in each tablet)</b>	<b>Purpose</b>
Loratadine 5 mg.....	Antihistamine
Pseudoephedrine sulfate 120 mg.....	Nasal decongestant

### Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - sneezing
  - runny nose
  - itching of the nose or throat
  - itchy, watery eyes
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

### Warnings

- Do not use**
- if you have ever had an allergic reaction to this product or any of its ingredients
  - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- heart disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.
- thyroid disease
- diabetes

## Drug Facts (continued)

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

### Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center rightaway (1-800-222-1222).

### Directions

- do not divide, crush, chew or dissolve the tablet
- adults and children 12 years and over: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: ask a doctor
- consumers with liver or kidney disease: ask a doctor

### Other information

- each tablet contains: calcium 25 mg
- do not use if blister unit is broken or torn
- store between 20° to 25° C (68° to 77° F)
- keep in a dry place

### Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

Questions or comments? 1-800-71-9-9260

7U046 29 C1

OPEN OTHER END

CONVENIENT RECLOSING TAB

ABC# 1 018 26 85



\*\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Claritin-D® 12 hour.

**GLUTEN FREE**

Distributed By AmerisourceBergen  
1300 Morris Drive, Chesterbrook, PA 19087



**GOOD NEIGHBOR PHARMACY LORATADINE D**

loratadine, pseudoephedrine sulfate tablet, film coated, extended release

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:46 122-455
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJ03BO7QN) (LORATADINE - UNII:7AJ03BO7QN)	LORATADINE	5 mg
PSEUDOEPHEDRINE SULFATE (UNII: Y9DL7QPE6B) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE SULFATE	120 mg

**Inactive Ingredients**

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

**Product Characteristics**

<b>Color</b>	WHITE (to off-white)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	7U0
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46 122-455-52	10 in 1 CARTON	04/23/20 18	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076050	04/23/2018	

