

IBUPROFEN- ibuprofen capsule, liquid filled
Sam's West, Inc.

Ibuprofen capsules, 200 mg

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)** (present as the free acid and potassium salt)

**nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use.
- store at 20°-25°C (68°-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

FD&C green #3, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution, and white ink

Questions or comments ?

Call toll free: **1-855-215-8180 Monday-Friday 8AM-8PM EST**

PRINCIPAL DISPLAY PANEL - Bottle Label

Sam's Club Ibuprofen Capsules, 200 mg

NDC: 68196-981-41

400 Softgels

Compare to Advil active ingredient Liqui-Gels

NDC 68196-981-41

Member's Mark™

Ibuprofen
Capsules, 200 mg mg

pain reliever/
fever reducer
(NSAID)

actual size

400
SOFTGELS+
Liquid Filled
Capsules

Do Not Use If seal under bottle cap is imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Drug Facts

Active ingredient (in each capsule)
Sulfobutyl isobutyrates equal to 200 mg ibuprofen (NSAID)**
(present as the free acid and potassium salt)
**nonsteroidal anti-inflammatory drug

Purpose
Pain reliever/fever reducer

Uses
temporarily relieves minor aches and pains due to:
 ■ headache ■ toothache ■ backache ■ menstrual cramps
 ■ the common cold ■ muscle aches ■ minor pain of arthritis
 ■ temporarily reduces fever

Warnings
 Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
 ■ skin redness/itching ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
 Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen)
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen) or blood thinning (anticoagulant) or steroid drug
 ■ take more or for a longer time than directed
 ■ take 3 or more alcoholic drinks every day while using this product
 ■ have heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery

Ask a doctor before use if
 ■ stomach bleeding warning applies to you
 ■ you are taking other pain relievers or fever reducers
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure/heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 ■ you are taking a diuretic

Drug Facts (continued under label)

DISTRIBUTED BY:
 SAMS CLUB DRUG
 BENTONVILLE, AR 72727
 Made in China
 SA 1802-00

For comments or suggestions,
 call us toll-free at 1-855-215-8180
 from 8:00 am. to 8:00 pm. EST
 Monday - Friday

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk, if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- you have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains potassium 20 mg
- read all warnings and directions before use
- store at 20°-25° C (68°-77° F)
- avoid excessive heat above 40° C (104° F)

Inactive ingredients FD&C green #3, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitol, sorbitan solution and white ink

Questions or comments? 1-855-215-8180 Monday-Friday 8AM-8PM EST

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark, Advil Liqui-Gels®

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68196-981
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C GREEN NO. 3 (UNII: 3P3ONR601S)	
GELATIN (UNII: 2G86QN327L)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0K00R)	

Product Characteristics

Color	green (Clear)	Score	no score
Shape	CAPSULE (Oblong)	Size	18mm
Flavor		Imprint Code	PC16
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68196-981-41	400 in 1 BOTTLE; Type 0: Not a Combination Product	04/16/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206568	04/16/2018	

Labeler - Sam's West, Inc. (051957769)

Establishment

Name	Address	ID/FEI	Business Operations
Humanwell PuraCap Pharmaceutical (Wuhan) Co., Ltd.		421293287	manufacture(68196-981)

Revised: 11/2019

Sam's West, Inc.