DAYTIME NIGHTTIME COLD FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl Care One (Retail Business Services, LLC.)

DRUG FACTS

Active ingredients for Daytime (in each softgel) Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Active ingredients for Nighttime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose for Daytime

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Purpose for Nighttime

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

DAYTIME

- temporarily relieves common cold and flu symptoms
 - cough due to minor throat and bronchial irritation
 - nasal congestion
 - headache
 - minor aches and pains
 - fever
 - sore throat

NIGHTTIME

- temporarily relieves common cold and flu symptoms
 - cough due to minor throat and bronchial irritation
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose and sneezing

Warnings

DAYTIME

Liver warning: These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using these products

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

NIGHTTIME

Liver warning: This product contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Alergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or following by fever, headache, rash, nausea, vomiting, consult a doctor promptly.

Do not use

DAYTIME

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

NIGHTTIME

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

DAYTIME

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

NIGHTTIME

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

DAYTIME

taking the blood thinning drug warfarin

NIGHTTIME

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product,

DAYTIME

do not exceed recommended dosage.

NIGHTTIME

- do not exceed recommended dosage
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

DAYTIME

- nervousness, dizziness or sleeplessness occur
- pain, cough, and nasal congestion gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- · cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

NIGHTTIME

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash headache that lasts.

These could be a signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

DAYTIME

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over; take 2 softgels with water every 4 hours.
- children under 12 years: do not use

NIGHTTIME

• do not take more than directed (see Overdose warning)

- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

Other information

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

Inactive ingredients

DAYTIME

FD&C red #40, FD&C yellow #6, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

NIGHTTIME

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Vicks $\$ Dayquil $\$ and Nyquil $\$ Cold $\$ Flu LiquiCaps $\$ $\$

DAYTIME

MULTI-SYMPTOM

COLD / FLU RELIEF

Pain reliever / Fever Reducer - Acetaminophen

Cough Suppressant - Dextromethorphan HBr

Nasal Decongestant - Phenylephrine HCI

Relieves Major cold & Flu Symptoms:

Aches & Fever, Sore Throat, Coughing, Congestion

SOFTGELS

NIGHTTIME

MULTI-SYMPTOM

COLD / FLU RELIEF

Pain Reliever/ Fever Reducer - Acetaminophen

Cough Suppressant - Dextromethorphan HBr

Antihistamine - Doxylamine succinate

Relieves Cold & Flu Symptoms

So You Can Rest

SOFTGELS

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil, NyQuil®, and LiquiCaps® are registered trademarks of the Procter and Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOW SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.

DISTRIBUTED BY
FOODHOLD U.S.A., LLC
LANDOVER, MD 20785
1-877-846-9949

Product Label

children under 12 years; do not use

- and children 12 years and over, take 2 softgels with water every
 - swallow whole; do not crush, chew, or dissolve
 - m do not take more than 4 doses in 24 hours
 - do not take more than directed (see Overdose warming)

Directions

notice any signs of symptoms. medical attention is critical for adults as well as for children even if you do not help or contact a Poison Control Center (1-800-222-1222) right away. Quick secommended dose can cause liver damage, in case of overdose, get medical If pregnant or breast-feeding, sak a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the

These could be signs of a serious condition.

- condu course pack or occurs with rash or headache that lasts.
 - Lequese or swelling is present new symptoms occur
 - fever gets worse or lasts more than 3 days
 - pain or cough gets worse or lasts more than 7 days
 - Stop use and ask a doctor If
- be careful when driving a motor vehicle or operating machinery
- a)cohol, sedatives, and tranquilizers may increase drowsiness
 - marked drowsiness may occur
 avoid alcoholic drinks
- excitability may occur, especially in children
- When using this product 🔳 do not exceed recom agesob babnan
- enestillupriert no eerstebee grisket m m taking the blood thinning drug warfarin Ask a doctor or pharmacist before use if you are
 - brisig elisienq begraine ne of eub gritarinu elduori ...
 - sathma, chronic bronchitis, or emphysema

Drug Facts (continued)

- s prestring problem or chronic cough that lasts such as occurs with smoking,
 - condu quet occruz wigu po wricu bujediu (wricna)
 - Ask a doctor before use if you have milver disease m glaucoma

Nighttime Cold & Plu Softgel

When using this product, do not exceed recommended dosage.

■ pain, cough, or nasal congestion gets worse or lasts more than 7 days

adults and children 12 years and over; take 2 sofigets with water every

medical attention is critical for adults as well as for children even if you do not

help or contact a Polson Control Center (1-800-222-1222) right away. Quick recommended dose can cause liver damage. In case of overdose, get medical Keep out of reach of children. Overdose warning: Taking more than the

If pregnant or breast-feeding, ask a feelth professional before use.

condu comes pack or occurs with useh or headache that lasts.

пілетем дить Ask a doctor or pharmacist before use if you are taking the blood thinning

Stop use and ask a doctor if

Drug Facts (continued)

SEK 9 GOCJOL OL DUSILUSCIEC

pharmacist before taking this product.

■ children under 12 years: do not use

swellow whole; do not crush, chew, or dissolve

do not take more than directed (see Overdose warning)

a do not take more than 4 doses in 24 hours

These could be signs of a serious condition.

■ nervousness, dizziness, or sleeplessness occur

■ Jaker gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur

SURROU D

Directions

nooce any signs or symptoms.

- persistent or chronic cough such as occurs with smoking, asthma,
 - m trouble unnafing due to an enlarged prostate gland
 - m condit that occurs with too much philegm (mucus)
- sesseib bionyth a sesseib a euresaud poorg ubiu ... ■ Tiver disease ■ heart disease Ask a doctor before use if you have

not know if your prescription drug contains an MAOI, ask a doctor or

(certain drugs for depression, psychiatric, or emotional conditions, or

If you are now taking a prescription monoamine oxidase inhibitor (MMOI).

with any other drug containing acetaminophen (prescription or

If a skin reaction occurs, stop use and seek medical help right away.

occur If you take: m more than 4,000 mg of acetaminophen in 24 hours

■ headache ■ minor aches and pains ■ fever ■ sore throat

nees temporarly reflexes common cold and flu symptoms:

m condu que po univor turos; suq proucujej jurisspou m uesej condespou

■ 3 or more alcoholic drinks every day while using this product

include: 🖷 sidn reddening 🗯 blisters 👅 rash

■ with other drugs containing acetaminophen

Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do

nonprescription). If you are not sure whether a drug contains acetaminophen,

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, rauses, or vomiting, consult

Altergy alert: Acetaminophen may cause severe skin reactions. Symptoms may

Daytime Cold & Flu Softgel

pharmacist before taking this product. not know if your prescription drug contains an MAOI, sak a doctor or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do (cartain drugs for depression, psychiatric, or emotional conditions, or

(KDAM) ratidirfní esebbo enimeonom molitriosenq a gnislat won ena uoy fi 🔳 sax a qoctor or praemactar.

nonprescription). If you are not sure whether a drug contains acetaminophen, with any other drug containing acetaminophen (prescription or

Sew fon oil

я досцо, былифай

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult Sore throat warming: If sore throat is severe, persists for more than 2 days, is It a skin reaction occurs, stop use and seek medical help right away. include: m skin neddening m blisters m rash

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may ■ 3 or more alcoholic drinks every day while using this product

■ with other drugs containing acetaminophen

Active ingredients (in each softgel)

occur if you take: m more than 4,000 mg of acetaminophen in 24 hours Liver warning: This product contains acetaminophen. Severe liver damage may Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may SEULUS

> ■ peadache ■ minor aches and pains ■ fever ■ runny nose & sneezing acough due to minor throat and bronchial inflation ■ sore throat **ПЗВЗ** ■ решбокачућ изјемег сошшои соја вид уп аћшћроша:

DUILLIESSULLIV condu anbbueseaur

sasoding

Pain reliever/fever reducer

Dextromethorphan HBr 10 mg. Acetaminophen 325 mg.

Phenylephrne HCl 5 mg

Pain reliever/fever reducer Active ingredients (in each softgel)

Nighttime Cold & Flu Softgel

jueosabuocap jesen

Cough suppressant

Purposes

Drug Facts Daytime Cold & Flu Softgel

NDC 72476-901-48

Drug Facts

Doxylamine succinate 6.25 mg.

Dextromethorphan HBr 15 mg.


Acetaminophen 325 mg.

Compare to the active ingredients in Vicks® DayQuil® and NyQuil® Cold & Flu LiquiCaps® †

NIGHTTIME MULTI SYMPTOM **COLD/FLU RELIEF**

CARE**ONE**

DAYTIME **MULTI SYMPTOM COLD/FLU RELIEF**



CAREONE Daytime Nighttime Multi-Symptom Cold & Flu Relief

DAYTIME NIGHTTIME COLD FLU RELIEF

Package Quantity

Part #

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Inform	Product Information				
Product Type	HUMAN OTC DRUG Item Code (Source) NDC:72476-901				
Packaging					
i ackaging					
# Item Code	Package Description	n	Marketing Start Date	Marketing End Date	
1 NDC:72476-901-	1 in 1 KIT; Type 0: Not a Combinati Product	tion	09/30/2019		
Quantity of Parts					
Qualitity of Faits					

Total Product Quantity

Part 1	16 BLISTER PACK	16
Part 2	32 BLISTER PACK	32

Part 1 of 2

NIGHTTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg		
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg		

Inactive Ingredients	
Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ 989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6092ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
MINERAL OIL (UNII: T5L8T28FGP)	

Product Characteristics				
Color	green	Score	no score	
Shape	CAPSULE	Size	21mm	
Flavor		Imprint Code	PC10	
Contains				

Packaging

#	t Item Code	Package Description	Marketing Start Date	Marketing End Date
1	L	16 in 1 CARTON		
1	L	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	09/30/2019		

Part 2 of 2

DAYTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients	
Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ 989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6092ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics				
Color	orange	Score	no score	
Shape	CAPSULE	Size	20mm	
Flavor		Imprint Code	PC9	
Contains				

Pa	Packaging				
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date	
1		32 in 1 CARTON			
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	09/30/2019		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/30/2019	

Labeler - Care One (Retail Business Services, LLC.) (967989935)

Revised: 3/2024 Care One (Retail Business Services, LLC.)