

DOCUSATE MINI ENEMA- docusate sodium liquid
Major Pharmaceuticals

MAJOR®

DOCUSATE

MINI ENEMA

30 Single Use

5 mL Disposable Tubes

Stool Softener Evacuant

Drug Facts

Active Ingredient (per delivered-dose)

Docusate Sodium USP 283 mg

Purpose

Stool Softener Laxative

Uses

- For relief of occasional constipation (irregularity).
- This product generally produces a bowel movement in 2 to 15 minutes.

Warnings

For rectal use only. Drug Interaction Precaution: Do not take this product if you are presently taking mineral oil, unless directed by a doctor.

Do not use

- Laxative products when abdominal pain, nausea or vomiting are present.
- Laxative products for a period longer than 1 week unless directed by a doctor.

Stop use and ask a doctor if you:

- Rectal bleeding.
- Noticed a sudden change in bowel habits that persists over a period of 2 weeks.
- Failed to have a bowel movement after use. This may indicate a serious condition.

Pregnant or Lactating Women

Ask a healthcare professional before use.

Keep out of reach of children

In case of accidental ingestion, get medical help or contact a Poison Control Center immediately.

Directions

Adults and children 12 years of age and older (with adult supervision) one to three units daily. Children under 12 years of age, consult a doctor prior to use.

- Twist off and remove tip.
- Lubricate tip prior to insertion: Place a few drops of the tube's liquid content on the shaft prior to insertion. Also apply liquid content or lubricant to the anus before inserting the mini-enema.
- Positioning: For best results, lay on the left side with knees bent. Insert slowly into rectum, up to shoulder of tube. Alternate Positions: Administer while seated on the toilet. Or, kneel, then lower head and chest forward until side of face is resting on the surface.
- With steady pressure, gently insert the tube into the rectum with care to prevent damage to the rectal wall. Squeeze to empty the contents. Keep the tube squeezed until it is removed from the rectum. After the contents have been emptied, remove the disposable tube and discard. A small amount of liquid may remain in the unit after use.

Other Information

Other Information Store at room temperature 15°-30° C (59°-86° F)

RETAIN BOTTLE FOR COMPLETE DRUG FACTS

Inactive Ingredients

Glycerin USP and Polyethylene Glycol

Questions or Comments: (800) 616-2471

MAJOR®

NDC# 0904-6920-93

DOCUSATE**MINI ENEMA**

30 Single Use
5ml Disposable Tubes
 Stool Softener Evacuant

LR120611E

FOR YOUR PROTECTION THIS BOTTLE HAS AN IMPRINTED SEAL AROUND THE NECK.

DOCUSATE MINI ENEMA

docusate sodium liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-6920
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	283 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 300 (UNII: 5655G9Y8AQ)	
GLYCERIN (UNII: PDC6A3C0OX)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-6920-93	30 in 1 JAR	08/26/2019	
1		5 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	08/26/2019	

Labeler - Major Pharmaceuticals (191427277)

Revised: 12/2025

Major Pharmaceuticals