

**NIGHTTIME COLD AND FLU MULTI SYMPTOM- acetaminophen
dextromethorphan hydrobromide doxylamine succinate capsule, liquid filled
WinCo Foods, LLC**

Safeway Nighttime Cold & Flu

Active Ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine Succinate 6.25 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

Other Information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive Ingredients

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or Comments?

Call **1-800-824-1706** Monday-Friday 9AM-4PM MST

Principal Display Panel

*Compare to the active ingredients in VICKS® NYQUIL® Cold & Flu LiquiCap®

MULTI-SYMP TOM

Cold & Flu Relief

Nighttime

ACETAMINOPHEN /

PAIN RELIEVER - FEVER REDUCER

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

DOXYLAMINE SUCCINATE / ANTIHISTAMINE

Relieves: Aches, Fever, Sore Throat, Cough, Sneezing & Runny Nose

Alcohol-Free

Softgels**

**Liquid-Filled Capsules

†This product is not manufactured or distributed by THE PROCTER & GAMBLE COMPANY. VICKS®, NyQuil®, and LiquiCaps® are registered trademarks of THE PROCTER & GAMBLE COMPANY.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY:

WINCO FOODS, LLC,

BOISE, ID 83704

Package Label

<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease glaucoma trouble urinating due to an enlarged prostate gland a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema cough that occurs with too much phlegm (mucus) <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking the blood thinning drug warfarin taking sedatives or tranquilizers <p>When using this product</p> <ul style="list-style-type: none"> do not exceed recommended dosage excitability may occur, especially in children avoid alcoholic drinks marked drowsiness may occur alcohol, sedatives, and tranquilizers may increase drowsiness be careful when driving a motor vehicle or operating machinery 	<p>Drug Facts (continued)</p> <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease glaucoma trouble urinating due to an enlarged prostate gland a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema cough that occurs with too much phlegm (mucus) <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking the blood thinning drug warfarin taking sedatives or tranquilizers <p>When using this product</p> <ul style="list-style-type: none"> do not exceed recommended dosage excitability may occur, especially in children avoid alcoholic drinks marked drowsiness may occur alcohol, sedatives, and tranquilizers may increase drowsiness be careful when driving a motor vehicle or operating machinery
<p>Directions</p> <ul style="list-style-type: none"> do not take more than 4 doses in 24 hours adults and children 12 years and over: take 2 softgels with water every 6 hours children under 12 years: do not use swallow whole; do not crush, chew, or dissolve <p>When using other Daytime or Nighttime products, carefully read each label to ensure correct dosing</p> <p>Other information</p> <ul style="list-style-type: none"> store between 15-30°C (59-86°F) avoid excessive heat <p>Inactive ingredients</p> <p>D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, propylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p>Questions or comments?</p> <p>Call 800-824-1706 Monday - Friday 9am - 4pm MST</p>	<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> pain or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>Keep out of reach of children.</p> <p>Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Do not take more than directed (see Overdose warning)</p>

<p>Drug Facts (continued)</p> <p>Warnings</p> <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours occur if you take: with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Drug Facts (continued)</p> <p>Purposes</p> <ul style="list-style-type: none"> Pain reliever/fever reducer Cough suppressant Antihistamine <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves common cold and flu symptoms: sore throat headache minor aches and pains fever 	<p>Drug Facts</p> <p>Active ingredients (in each softgel)</p> <ul style="list-style-type: none"> Acetaminophen 325 mg Dextromethorphan HBr 15 mg Doxylamine succinate 6.25 mg Antihistamine
--	---



MULTI-SYMPTOM Cold & Flu Relief

Nighttime

**ACETAMINOPHEN/
PAIN RELIEVER - FEVER REDUCER**

DEXTROMETHORPHAN HBr/COUGH SUPPRESSANT

DOXYLAMINE SUCCINATE/ANTIHISTAMINE

24

Softgels **

**Liquid-Filled Capsules

Relieves: Aches, Fever, Sore Throat, Cough, Sneezing & Runny Nose

.....

Alcohol-free

*Compare to the active ingredients in VICKS® NYQUIL® Cold & Flu LiquiCaps®

Lot No.:

Exp. Date:

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

* This product is not manufactured or distributed by THE PROCTER & GAMBLE COMPANY. VICKS®, NyQuil®, and LiquiCaps® are registered trademarks of THE PROCTER & GAMBLE COMPANY.

DISTRIBUTED BY:
WINCO FOODS, LLC,
BOISE, ID 83704
PRODUCT OF CHINA



The Supermarket Low Price Leader®

100% SATISFACTION GUARANTEE

Comments & Questions?
800-824-1706 · www.wincofoods.com

FC007769 PLD-E585B WF2110



WINCO FOODS Nighttime Multi-Symptom Cold & Flu Relief

NIGHTTIME COLD AND FLU MULTI SYMPTOM

acetaminophen dextromethorphan hydrobromide doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67091-395
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics

Color	green (Clear)	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	PC10
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-395-24	24 in 1 CARTON	10/29/2021	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/29/2021	

Labeler - WinCo Foods, LLC (056098817)

Revised: 5/2024

WinCo Foods, LLC