

**ED A-HIST- chlorpheniramine maleate and phenylephrine hydrochloride tablet, coated**  
**EDWARDS PHARMACEUTICALS, INC.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**ED A-HIST TABLETS**

**Drug Facts**

<b>Active Ingredients (in each tablet)</b>	<b>Purpose</b>
Chlorpheniramine Maleate 4 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal Decongestant

**Uses**

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

**Warnings**

- **Do not exceed recommended dosage.**

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are taking sedatives or**

## tranquilizers

### When using this product

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by a fever
- new symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

### Directions

**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	1 tablet every 4 hours, not to exceed 6 tablets in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age:	1/2 tablet every 4 hours, not to exceed 3 tablets in 24 hours, or as directed by a doctor
Children under 6 years of age:	Consult a doctor.

### Inactive ingredients

FD&C Blue #2, FD&C Yellow #5, FD&C Yellow #6, Magnesium Stearate, Microcrystalline Cellulose, Sodium Starch Glycolate. Contains FD&C Yellow No. 5 (tartrazine) as a color additive

### Questions or Comments?

Call 1-800-543-9560 Rev.

**PRINCIPAL DISPLAY PANEL - 100 Tablet Bottle Label**

NDC 0485-0254-01

## **ED A-HIST TABLETS**

Antihistamine • Nasal Decongestant

### **Each tablet contains:**

Chlorpheniramine Maleate 4 mg

Phenylephrine HCl 10 mg

Store at 59°-86°F (15°-30°C) [see USP Controlled Room Temperature].

Tamper evident by foil seal under cap.

Do not use if foil seal is broken or missing.

### **Manufactured for:**

***EDWARDS***

**Pharmaceuticals, Inc.**

**Ripley, MS 38663**

100 tablets



NDC 0485-0254-01

# ED A-HIST TABLETS

Antihistamine • Nasal Decongestant

**Each tablet contains:**

- Chlorpheniramine Maleate ..... 4 mg
- Phenylephrine HCl ..... 10 mg

Store at 59°-86°F (15°-30°C) [see USP Controlled Room Temperature].  
 Tamper evident by foil seal under cap.  
 Do not use if foil seal is broken or missing.

Manufactured for:  
**EDWARDS**  
 Pharmaceuticals, Inc.  
 Ripley, MS 38663

100 tablets



## Drug Facts

Active Ingredients	Purpose
<i>(in each tablet)</i>	
Chlorpheniramine Maleate 4 mg .....	Antihistamine
Phenylephrine HCl 10 mg .....	Nasal Decongestant

**Uses** temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes
- nasal congestion ■ reduces swelling of nasal passages

Lot:  
Exp. Date:

Peel Here

### Drug Facts (continued)

#### Warnings

- Do not exceed recommended dosage.

#### Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma ■ heart disease
- high blood pressure ■ thyroid disease ■ diabetes mellitus
- difficulty in urination due to enlargement of the prostate gland

#### Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

#### When using this product

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages

### Drug Facts (continued)

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by a fever ■ new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use.

#### Keep out of reach of children.

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.


#### Directions

#### Do not exceed recommended dosage.

Adults and children 12 years of age and over:	1 tablet every 4 hours, not to exceed 6 tablets in 24 hours, or as directed by a doctor
Children 6 to under 12 years of age:	1/2 tablet every 4 hours, not to exceed 3 tablets in 24 hours, or as directed by a doctor
Children under 6 years of age:	Consult a doctor.

#### Inactive ingredients

FD&C Blue #2, FD&C Yellow #5, FD&C Yellow #6, Magnesium Stearate, Microcrystalline Cellulose, Sodium Starch Glycolate.



## ED A-HIST

chlorpheniramine maleate and phenylephrine hydrochloride tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0485-0254
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)	
<b>FD&amp;C YELLOW NO. 5</b> (UNII: I753WB2F1M)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	

### Product Characteristics

<b>Color</b>	brown (WHEAT)	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	13mm
<b>Flavor</b>		<b>Imprint Code</b>	E;1
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0485-0254-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/08/2011	

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
unapproved drug other		10/08/2011	

**Labeler** - EDWARDS PHARMACEUTICALS, INC. (195118880)

Revised: 11/2024

EDWARDS PHARMACEUTICALS, INC.