

# **GLIPIZIDE- glipizide tablet, extended release**

## **Northstar Rx LLC.**

-----

### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use GLIPIZIDE EXTENDED-RELEASE TABLETS safely and effectively. See full prescribing information for GLIPIZIDE EXTENDED-RELEASE TABLETS.**

**GLIPIZIDE extended-release tablets, for oral use**  
**Initial U.S. Approval: 1994**

### **INDICATIONS AND USAGE**

Glipizide extended-release tablets is a sulfonylurea indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus. Limitations of Use: Not for treatment of type 1 diabetes or diabetic ketoacidosis. (1)

### **DOSAGE AND ADMINISTRATION**

- Recommended starting dose is 5 mg once daily. Dose adjustment can be made based on the patient's glycemic control. Maximum recommended dose is 20 mg once daily (2.1).
- Administer with breakfast or the first meal of the day (2.1).
- For combination therapy with other blood-glucose-lowering agents, initiate the agent at the lowest recommended dose, and observe patients for hypoglycemia (2.2).

### **DOSAGE FORMS AND STRENGTHS**

Tablets: 2.5 mg, 5 mg, 10 mg (3).

### **CONTRAINDICATIONS**

- Known hypersensitivity to glipizide or any of the product's ingredients (4)
- Hypersensitivity to sulfonamide derivatives (4)

### **WARNINGS AND PRECAUTIONS**

- Hypoglycemia: May be severe. Ensure proper patient selection, dosing, and instructions, particularly in at-risk populations (e.g., elderly, renally impaired) and when used with other anti-diabetic medications (5.1).
- Hemolytic Anemia: Can occur if glucose 6-phosphate dehydrogenase (G6PD) deficient. Consider a non-sulfonylurea alternative (5.2).
- Potential Increased Risk of Cardiovascular Mortality with Sulfonylureas: Inform patient of risks, benefits and treatment alternatives (5.3).
- Macrovascular Outcomes: No clinical studies have established conclusive evidence of macrovascular risk reduction with glipizide extended-release tablets or any other anti-diabetic drug (5.4).

### **ADVERSE REACTIONS**

Most common adverse reactions (incidence > 3%) are dizziness, diarrhea, nervousness, tremor, hypoglycemia and flatulence (6.1).

**To report SUSPECTED ADVERSE REACTIONS, contact Northstar Rx LLC at 1-800-206-7821 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### **DRUG INTERACTIONS**

- Certain medications may affect glucose metabolism, requiring glipizide extended-release tablets dose adjustment and close monitoring of blood glucose (7.1)
- Miconazole: Monitor patients closely. Severe hypoglycemia can occur when glipizide and oral miconazole are used concomitantly (7.1, 12.3).
- Fluconazole: Monitor patients closely. An increase in glipizide AUC was seen after fluconazole administration (7.3, 12.3).
- Colesevelam: Glipizide extended-release tablets should be administered at least 4 hours prior to colesevelam (7.4, 12.3).

### **USE IN SPECIFIC POPULATIONS**

- Geriatric, Hepatically Impaired Patients: At risk for hypoglycemia with glipizide extended-release tablets. Use caution in dose selection and titration, and monitor closely (8.5, 8.6).

**See 17 for PATIENT COUNSELING INFORMATION, FDA-approved patient labeling and FDA-**

---

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

1.1 Limitations of Use

**2 DOSAGE AND ADMINISTRATION**

2.1 Recommended Dosing

2.2 Use with Other Glucose Lowering Agents

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

5.1 Hypoglycemia

5.2 Hemolytic Anemia

5.3 Increased Risk of Cardiovascular Mortality with Sulfonylureas

5.4 Macrovascular Outcomes

5.5 Gastrointestinal Obstruction

**6 ADVERSE REACTIONS**

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

**7 DRUG INTERACTIONS**

7.1 Drugs Affecting Glucose Metabolism

7.2 Miconazole

7.3 Fluconazole

7.4 Colesevelam

**8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

**10 OVERDOSAGE**

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**15 REFERENCES**

**16 HOW SUPPLIED/STORAGE AND HANDLING**

**17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Glipizide extended-release tablets are indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.

#### **1.1 Limitations of Use**

Glipizide extended-release tablets are not recommended for the treatment of type 1 diabetes mellitus or diabetic ketoacidosis.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Recommended Dosing**

Glipizide extended-release tablets should be administered orally with breakfast or the first main meal of the day.

The recommended starting dose of glipizide extended-release tablets is 5 mg once daily. Start patients at increased risk for hypoglycemia (e.g. the elderly or patients with hepatic insufficiency) at 2.5 mg [*see Use in Specific Population (8.5, 8.6)*].

Dosage adjustment can be made based on the patient's glycemic control. The maximum recommended dose is 20 mg once daily.

Patients receiving immediate-release glipizide may be switched to glipizide extended-release tablets once daily at the nearest equivalent total daily dose.

#### **2.2 Use with Other Glucose Lowering Agents**

When adding glipizide extended-release tablets to other anti-diabetic drugs, initiate glipizide extended-release tablets at 5 mg once daily. Start patients at increased risk for hypoglycemia at a lower dose.

When colesevelam is coadministered with glipizide extended-release tablets, maximum plasma concentration and total exposure to glipizide is reduced. Therefore, glipizide extended-release tablets should be administered at least 4 hours prior to colesevelam.

### **3 DOSAGE FORMS AND STRENGTHS**

Glipizide extended-release tablets USP, 2.5 mg are yellow colored, round, biconvex film-coated tablets imprinted with "2" on one side with black ink and plain on the other side.

Glipizide extended-release tablets USP, 5 mg are orange colored, round, biconvex film-coated tablets imprinted with "3" on one side with black ink and plain on the other side.

Glipizide extended-release tablets USP, 10 mg are white colored, round, biconvex film-coated tablets imprinted with "4" on one side and plain on the other side.

### **4 CONTRAINDICATIONS**

Glipizide is contraindicated in patients with:

- Known hypersensitivity to glipizide or any of the product's ingredients.
- Hypersensitivity to sulfonamide derivatives.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Hypoglycemia**

All sulfonylurea drugs, including glipizide extended-release tablets, are capable of producing severe hypoglycemia [see *Adverse Reactions* (6)]. Concomitant use of glipizide extended-release tablets with other anti-diabetic medication can increase the risk of hypoglycemia. A lower dose of glipizide extended-release tablets may be required to minimize the risk of hypoglycemia when combining it with other anti-diabetic medications.

Educate patients to recognize and manage hypoglycemia. When initiating and increasing glipizide extended-release tablets in patients who may be predisposed to hypoglycemia (e.g., the elderly, patients with renal impairment, patients on other anti-diabetic medications) start at 2.5 mg. Debilitated or malnourished patients, and those with adrenal, pituitary, or hepatic impairment are particularly susceptible to the hypoglycemic action of anti-diabetic medications. Hypoglycemia is also more likely to occur when caloric intake is deficient, after severe or prolonged exercise, or when alcohol is ingested.

The patient's ability to concentrate and react may be impaired as a result of hypoglycemia. Early warning symptoms of hypoglycemia may be different or less pronounced in patients with autonomic neuropathy, the elderly, and in patients who are taking beta-adrenergic blocking medications or other sympatholytic agents. These situations may result in severe hypoglycemia before the patient is aware of the hypoglycemia.

These impairments may present a risk in situations where these abilities are especially important, such as driving or operating other machinery. Severe hypoglycemia can lead to unconsciousness or convulsions and may result in temporary or permanent impairment of brain function or death.

### **5.2 Hemolytic Anemia**

Treatment of patients with glucose 6-phosphate dehydrogenase (G6PD) deficiency with sulfonylurea agents, including glipizide extended-release tablets, can lead to hemolytic anemia. Avoid use of glipizide extended-release tablets in patients with G6PD deficiency. In post marketing reports, hemolytic anemia has also been reported in patients who did not have known G6PD deficiency.

### **5.3 Increased Risk of Cardiovascular Mortality with Sulfonylureas**

The administration of oral hypoglycemic drugs has been reported to be associated with increased cardiovascular mortality as compared to treatment with diet alone or diet plus insulin. This warning is based on the study conducted by the University Group Diabetes Program (UGDP), a long-term prospective clinical trial designed to evaluate the effectiveness of glucose-lowering drugs in preventing or delaying vascular complications in patients with type 2 diabetes mellitus. The study involved 823 patients who were

randomly assigned to one of four treatment groups.

UGDP reported that patients treated for 5 to 8 years with diet plus a fixed dose of tolbutamide (1.5 grams per day) had a rate of cardiovascular mortality approximately 2½ times that of patients treated with diet alone. A significant increase in total mortality was not observed, but the use of tolbutamide was discontinued based on the increase in cardiovascular mortality, thus limiting the opportunity for the study to show an increase in overall mortality. Despite controversy regarding the interpretation of these results, the findings of the UGDP study provide an adequate basis for this warning. The patient should be informed of the potential risks and advantages of glipizide and of alternative modes of therapy.

Although only one drug in the sulfonylurea class (tolbutamide) was included in this study, it is prudent from a safety standpoint to consider that this warning may also apply to other oral hypoglycemic drugs in this class, in view of their close similarities in mode of action and chemical structure.

#### **5.4 Macrovascular Outcomes**

There have been no clinical studies establishing conclusive evidence of macrovascular risk reduction with glipizide extended-release tablets or any other anti-diabetic drug.

#### **5.5 Gastrointestinal Obstruction**

There have been reports of obstructive symptoms in patients with known strictures in association with the ingestion of another drug with this non-dissolvable extended release formulation. Avoid use of glipizide extended-release tablets in patients with preexisting severe gastrointestinal narrowing (pathologic or iatrogenic).

*The 2.5 mg tablets contain FD&C Yellow No.5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No.5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.*

### **6 ADVERSE REACTIONS**

The following serious adverse reactions are discussed in more detail below and elsewhere in the labeling:

- Hypoglycemia [see *Warnings and Precautions* (5.1) ]
- Hemolytic anemia [see *Warnings and Precautions* (5.2) ]

#### **6.1 Clinical Trials Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In clinical trials, 580 patients from 31 to 87 years of age received glipizide extended-release tablets in doses from 5 mg to 60 mg in both controlled and open trials. The dosages above 20 mg are not recommended dosages. In these trials, approximately 180 patients were treated with glipizide extended-release tablets for at least 6 months.

Table 1 summarizes the incidence of adverse reactions, other than hypoglycemia, that

were reported in pooled double-blind, placebo-controlled trials in  $\geq 3\%$  of glipizide extended-release tablets-treated patients and more commonly than in patients who received placebo.

**Table 1 Incidence (%) of Adverse Reactions Reported in  $\geq 3\%$  of Patients Treated in Placebo-Controlled Clinical Trials and More Commonly in Patients Treated with glipizide extended-release tablets(Excluding Hypoglycemia)**

|                       | <b>Glipizide extended-release tablets (%)</b> | <b>Placebo (%)</b> |
|-----------------------|-----------------------------------------------|--------------------|
|                       | <b>( N=278)</b>                               | <b>(N=69)</b>      |
| <b>Adverse Effect</b> |                                               |                    |
| Dizziness             | 6.8                                           | 5.8                |
| Diarrhea              | 5.4                                           | 0.0                |
| Nervousness           | 3.6                                           | 2.9                |
| Tremor                | 3.6                                           | 0.0                |
| Flatulence            | 3.2                                           | 1.4                |

**Hypoglycemia:** Of the 580 patients that received glipizide extended-release tablets in clinical trials, 3.4% had hypoglycemia documented by a blood-glucose measurement  $<60$  mg/dL and/or symptoms believed to be associated with hypoglycemia and 2.6% of patients discontinued for this reason. Hypoglycemia was not reported for any placebo patients.

**Gastrointestinal Reactions:** In clinical trials, the incidence of gastrointestinal (GI) side effects (nausea, vomiting, constipation, dyspepsia), occurred in less than 3% of glipizide extended-release tablets-treated patients and were more common in glipizide extended-release tablets-treated patients than those receiving placebo.

### **Dermatologic Reactions**

In clinical trials, allergic skin reactions, i.e., urticaria occurred in less than 1.5% of treated patients and were more common in glipizide extended-release tablets treated patients than those receiving placebo. These may be transient and may disappear despite continued use of glipizide extended-release tablets; if skin reactions persist, the drug should be discontinued.

### **Laboratory Tests**

Mild to moderate elevations of ALT, LDH, alkaline phosphatase, BUN and creatinine have been noted. The relationship of these abnormalities to glipizide is uncertain.

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during post approval use of glipizide extended-release tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Abdominal pain
- Cholestatic and hepatocellular forms of liver injury accompanied by jaundice
- Leukopenia, agranulocytosis, thrombocytopenia, hemolytic anemia [*see Warnings and*

*Precautions* (5.2) ], aplastic anemia, pancytopenia

- Hepatic porphyria and disulfiram-like reactions
- Hyponatremia and the syndrome of inappropriate antidiuretic hormone (SIADH) secretion
- Rash
- There have been reports of gastrointestinal irritation and gastrointestinal bleeding with use of another drug with this non-dissolvable extended release formulation.

## **7 DRUG INTERACTIONS**

### **7.1 Drugs Affecting Glucose Metabolism**

A number of medications affect glucose metabolism and may require glipizide extended-release tablets dose adjustment and close monitoring for hypoglycemia or worsening glycemic control.

The following are examples of medication that may increase the glucose lowering effect of glipizide extended-release tablets, increase the susceptibility to and/or intensity of hypoglycemia: antidiabetic agents, ACE inhibitors, angiotensin II receptor blocking agents, disopyramide, fibrates, fluoxetine, monoamine oxidase inhibitors, pentoxifylline, pramlintide, propoxyphene, salicylates, somatostatin analogs (e.g., octreotide), sulfonamide antibiotics, nonsteroidal anti-inflammatory agents, chloramphenicol, probenecid, coumarins, voriconazole, H2 receptor antagonists, and quinolones. When these medications are administered to a patient receiving glipizide extended-release tablets, monitor the patient closely for hypoglycemia. When these medications are discontinued from a patient receiving glipizide extended-release tablets, monitor the patient closely for worsening glycemic control.

The following are examples of medication that may reduce the glucose-lowering effect of glipizide extended-release tablets, leading to worsening glycemic control: atypical antipsychotics (e.g., olanzapine and clozapine), corticosteroids, danazol, diuretics, estrogens, glucagon, isoniazid, niacin, oral contraceptives, phenothiazines, progestogens (e.g., in oral contraceptives), protease inhibitors, somatropin, sympathomimetic agents (e.g., albuterol, epinephrine, terbutaline), thyroid hormones, phenytoin, nicotinic acid, and calcium channel blocking drugs. When such drugs are administered to patients receiving glipizide extended-release tablets, monitor the patients closely for worsening glycemic control. When these medications are discontinued from patients receiving glipizide extended-release tablets, monitor the patient closely for hypoglycemia.

Alcohol, beta-blockers, clonidine, and reserpine may lead to either potentiation or weakening of the glucose-lowering effect. Increased frequency of monitoring may be required when glipizide extended-release tablets is co-administered with these drugs.

The signs of hypoglycemia may be reduced or absent in patients taking sympatholytic drugs such as beta-blockers, clonidine, guanethidine, and reserpine. Increased frequency of monitoring may be required when glipizide extended-release tablets is co-administered with these drugs.

### **7.2 Miconazole**

Monitor patients closely for hypoglycemia when glipizide extended-release tablets are co-

administered with miconazole. A potential interaction between oral miconazole and oral hypoglycemic agents leading to severe hypoglycemia has been reported [see *Clinical Pharmacology* (12.3)].

### **7.3 Fluconazole**

Monitor patients closely for hypoglycemia when glipizide extended-release tablets are co-administered with fluconazole. Concomitant treatment with fluconazole increases plasma concentrations of glipizide, which may lead to hypoglycemia [see *Clinical Pharmacology* (12.3)].

### **7.4 Colesevelam**

Glipizide extended-release tablets should be administered at least 4 hours prior to the administration of colesevelam. Colesevelam can reduce the maximum plasma concentration and total exposure of glipizide when the two are coadministered [see *Clinical Pharmacology* (12.3)].

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### ***Risk Summary***

Available data from a small number of published studies and postmarketing experience with glipizide extended-release tablets use in pregnancy over decades have not identified any drug associated risks for major birth defects, miscarriage, or adverse maternal outcomes. However, sulfonylureas (including glipizide) cross the placenta and have been associated with neonatal adverse reactions such as hypoglycemia. Therefore, glipizide extended-release tablets should be discontinued at least two weeks before expected delivery (see Clinical Considerations). Poorly controlled diabetes in pregnancy is also associated with risks to the mother and fetus (see Clinical Considerations). In animal studies, there were no effects on embryofetal development following administration of glipizide to pregnant rats and rabbits during organogenesis at doses 833 times and 8 times the human dose based on body surface area, respectively. However, increased pup mortality was observed in rats administered glipizide from gestation day 15 throughout lactation at doses 2 times the maximum human dose based on body surface area (see Data).

The estimated background risk of major birth defects is 6-10% in women with pre-gestational diabetes with a HbA1c >7 and has been reported to be as high as 20 to 25% in women with HbA1c >10. The estimated background risk of miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

#### ***Clinical Considerations***

##### ***Disease-Associated Maternal and/or Embryo/Fetal Risk***

Poorly-controlled diabetes in pregnancy increases the maternal risk for diabetic

ketoacidosis, pre-eclampsia, miscarriage, preterm delivery, stillbirth, and delivery complications. Poorly controlled diabetes increases the fetal risk for major birth defects, stillbirth, and macrosomia related morbidity.

### *Fetal/Neonatal Adverse Reactions*

Neonates of women with gestational diabetes who are treated with sulfonylureas during pregnancy may be at increased risk for neonatal intensive care admission and may develop respiratory distress, hypoglycemia, birth injury, and be large for gestational age. Prolonged severe hypoglycemia, lasting 4 to 10 days, has been reported in neonates born to mothers receiving a sulfonylurea at the time of delivery and has been reported with the use of agents with a prolonged half-life. Observe newborns for symptoms of hypoglycemia and respiratory distress and manage accordingly.

### *Dose adjustments during pregnancy and the postpartum period*

Due to reports of prolonged severe hypoglycemia in neonates born to mothers receiving a sulfonylurea at the time of delivery, glipizide extended-release tablets should be discontinued at least two weeks before expected delivery (see Fetal/Neonatal Adverse Reactions).

## **Data**

### *Animal Data*

In teratology studies in rats and rabbits, pregnant animals received daily oral doses of glipizide during the period of organogenesis at doses up to 2000 mg/kg/day and 10 mg/kg/day (approximately 833 and 8 times the human dose based on body surface area), respectively. There were no adverse effects on embryo-fetal development at any of the doses tested. In a peri- and postnatal study in pregnant rats, there was a reduced number of pups born alive following administration of glipizide from gestation day 15 throughout lactation through weaning at doses  $\geq 5$  mg/kg/day (about 2 times the recommended maximum human dose based on body surface area).

## **8.2 Lactation**

### **Risk Summary**

Breastfed infants of lactating women using glipizide extended-release tablets should be monitored for symptoms of hypoglycemia (see Clinical Considerations). Although glipizide was undetectable in human milk in one small clinical lactation study; this result is not conclusive because of the limitations of the assay used in the study. There are no data on the effects of glipizide on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for glipizide extended-release tablets and any potential adverse effects on the breastfed child from glipizide extended-release tablets or from the underlying maternal condition.

## **Clinical Considerations**

### *Monitoring for adverse reactions*

Monitor breastfed infants for signs of hypoglycemia (e.g., jitters, cyanosis, apnea, hypothermia, excessive sleepiness, poor feeding, seizures).

## 8.4 Pediatric Use

Safety and effectiveness in children have not been established.

## 8.5 Geriatric Use

There were no overall differences in effectiveness or safety between younger and older patients, but greater sensitivity of some individuals cannot be ruled out. Elderly patients are particularly susceptible to the hypoglycemic action of anti-diabetic agents.

Hypoglycemia may be difficult to recognize in these patients. Therefore, dosing should be conservative to avoid hypoglycemia. [see *Dosage and Administration* (2.1), *Warnings and Precautions* (5.1) and *Clinical Pharmacology* (12.3)]

## 8.6 Hepatic Impairment

There is no information regarding the effects of hepatic impairment on the disposition of glipizide. However, since glipizide is highly protein bound and hepatic biotransformation is the predominant route of elimination, the pharmacokinetics and/or pharmacodynamics of glipizide may be altered in patients with hepatic impairment. If hypoglycemia occurs in such patients, it may be prolonged and appropriate management should be instituted.

[see *Dosage and Administration* (2.1), *Warnings and Precautions* (5.1) and *Clinical Pharmacology* (12.3)]


## 10 OVERDOSAGE

Overdosage of sulfonylureas including glipizide extended-release tablets can produce severe hypoglycemia. Mild hypoglycemic symptoms without loss of consciousness or neurologic findings should be treated with oral glucose. Severe hypoglycemic reactions with coma, seizure, or other neurological impairment are medical emergencies requiring immediate treatment. The patient should be treated with glucagon or intravenous glucose. Patients should be closely monitored for a minimum of 24 to 48 hours since hypoglycemia may recur after apparent clinical recovery. Clearance of glipizide from plasma may be prolonged in persons with liver disease. Because of the extensive protein binding of glipizide, dialysis is unlikely to be of benefit.

## 11 DESCRIPTION

Glipizide extended-release tablets, USP are an oral sulfonylurea.

The Chemical Abstracts name of glipizide is 1-cyclohexyl-3-[[p-[2-(5-methylpyrazinecarboxamido) ethyl] phenyl]sulfonyl]urea. The molecular formula is  $C_{21}H_{27}N_5O_4S$ ; the molecular weight is 445.55; the structural formula is shown below:



Glipizide, USP is a white to off-white powder, with a pKa of 5.9. It is freely soluble in dimethylformamide, soluble in 0.1N sodium hydroxide and slightly soluble in methylene chloride. Glipizide extended-release tablets, USP are formulated as a once-a-day extended-release tablet for oral use and are designed to deliver 2.5 mg, 5 mg, or 10 mg of glipizide.

Each glipizide extended-release tablet, USP contains the following inactive ingredients: acetyltributyl citrate, colloidal silicon dioxide, hydroxyethyl cellulose, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, methacrylic acid copolymer and polyethylene glycol. Additionally each 2.5 mg tablet contains: FD&C yellow #5 aluminum lake and titanium dioxide. Each 5 mg tablet contains: FD&C yellow #6 aluminum lake and titanium dioxide. The tablet is imprinted with opacode black S-1-17823 which contains following ingredients: ammonium hydroxide, iron oxide black, isopropyl alcohol, n-butyl alcohol, propylene glycol and shellac.

### **System Components and Performance**

Glipizide extended-release tablets are formulated as once-a-day extended-release tablets and are designed to deliver glipizide at a controlled rate over approximately 20 hours. The dosage form is comprised of a hydrophilic cellulose polymer matrix tablet containing the drug which is surrounded by a seal coat followed by an enteric coating system. The enteric coat is insoluble in the low pH environment of the stomach. As the tablet passes through the stomach and enters in the higher pH environment of the small intestine, the enteric coating dissolves and/or erodes to expose the polymer matrix tablet which swells and releases the drug at a controlled rate via diffusion and/or erosion.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Glipizide primarily lowers blood glucose by stimulating the release of insulin from the pancreas, an effect dependent upon functioning beta cells in the pancreatic islets. Sulfonylureas bind to the sulfonylurea receptor in the pancreatic beta-cell plasma membrane, leading to closure of the ATP-sensitive potassium channel, thereby stimulating the release of insulin.

### **12.2 Pharmacodynamics**

The insulinotropic response to a meal is enhanced with glipizide extended-release tablets

administration in diabetic patients. The postprandial insulin and C-peptide responses continue to be enhanced after at least 6 months of treatment. In two randomized, double-blind, dose-response studies comprising a total of 347 patients, there was no significant increase in fasting insulin in all glipizide extended-release tablets-treated patients combined compared to placebo, although minor elevations were observed at some doses.

In studies of glipizide extended-release tablets in subjects with type 2 diabetes mellitus, once daily administration produced reductions in hemoglobin A1c, fasting plasma glucose and postprandial glucose. The relationship between dose and reduction in hemoglobin A1c was not established, however subjects treated with 20 mg had a greater reduction in fasting plasma glucose compared to subjects treated with 5 mg.

## **12.3 Pharmacokinetics**

### **Absorption**

The absolute bioavailability of glipizide was 100% after single oral doses in patients with type 2 diabetes mellitus. Beginning 2 to 3 hours after administration of glipizide extended-release tablets, plasma drug concentrations gradually rise reaching maximum concentrations within 6 to 12 hours after dosing. With subsequent once daily dosing of glipizide concentrations are maintained throughout the 24 hour dosing interval with less peak to trough fluctuation than that observed with twice daily dosing of immediate release glipizide.

The mean relative bioavailability of glipizide in 21 males with type 2 diabetes mellitus after administration of 20 mg glipizide extended-release tablets, compared to immediate release Glipizide (10 mg given twice daily), was 90% at steady-state. Steady-state plasma concentrations were achieved by at least the fifth day of dosing with glipizide extended-release tablets in 21 males with type 2 diabetes mellitus and patients younger than 65 years. No accumulation of drug was observed in patients with type 2 diabetes mellitus during chronic dosing with glipizide extended-release tablets.

Administration of glipizide extended-release tablets with food has no effect on the 2 to 3 hour lag time in drug absorption. In a single dose, food effect study in 21 healthy male subjects, the administration of glipizide extended-release tablets immediately before a high fat breakfast resulted in a 40% increase in the glipizide mean  $C_{max}$  value, which was significant, but the effect on the AUC was not significant. There was no change in glucose response between the fed and fasting state. Markedly reduced GI retention times of the glipizide extended-release tablets over prolonged periods (e.g., short bowel syndrome) may influence the pharmacokinetic profile of the drug and potentially result in lower plasma concentrations.

In a multiple dose study in 26 males with type 2 diabetes mellitus, the pharmacokinetics of glipizide were linear with glipizide extended-release tablets in that the plasma drug concentrations increased proportionately with dose. In a single dose study in 24 healthy subjects, four 5-mg, two 10-mg, and one 20-mg glipizide extended-release tablets were bioequivalent. In a separate single dose study in 36 healthy subjects, four 2.5-mg glipizide extended-release tablets were bioequivalent to one 10-mg glipizide extended-release tablets.

### **Distribution**

The mean volume of distribution was approximately 10 liters after single intravenous

doses in patients with type 2 diabetes mellitus. Glipizide is 98–99% bound to serum proteins, primarily to albumin.

## **Metabolism**

The major metabolites of glipizide are products of aromatic hydroxylation and have no hypoglycemic activity. A minor metabolite, an acetamino-ethyl benzene derivative, which accounts for less than 2% of a dose, is reported to have 1/10 to 1/3 as much hypoglycemic activity as the parent compound.

## **Elimination**

Glipizide is eliminated primarily by hepatic biotransformation: less than 10% of a dose is excreted as unchanged drug in urine and feces; approximately 90% of a dose is excreted as biotransformation products in urine (80%) and feces (10%). The mean total body clearance of glipizide was approximately 3 liters per hour after single intravenous doses in patients with type 2 diabetes mellitus. The mean terminal elimination half-life of glipizide ranged from 2 to 5 hours after single or multiple doses in patients with type 2 diabetes mellitus.

## **Specific Populations**

### **Pediatric**

Studies characterizing the pharmacokinetics of glipizide in pediatric patients have not been performed.

### **Geriatric**

There were no differences in the pharmacokinetics of glipizide after single dose administration to older diabetic subjects compared to younger healthy subjects. [see *Use in Specific Populations* (8.5)]

### **Renal Impairment**

The pharmacokinetics of glipizide has not been evaluated in patients with varying degree of renal impairment. Limited data indicates that glipizide biotransformation products may remain in circulation for a longer time in subjects with renal impairment than that seen in subjects with normal renal function.

### **Hepatic Impairment**

The pharmacokinetics of glipizide has not been evaluated in patients with hepatic impairment.

## **Drug-drug Interactions**

### **Miconazole**

A potential interaction between oral miconazole and oral glipizide leading to severe hypoglycemia has been reported. Whether this interaction also occurs with the intravenous, topical, or vaginal preparations of miconazole is not known. [see *Drug Interactions* (7.2)]

### **Fluconazole**

Concomitant treatment with fluconazole increases plasma concentrations of glipizide. The effect of concomitant administration of Diflucan® (fluconazole) and Glipizide has been demonstrated in a placebo controlled crossover study in healthy volunteers. All

been demonstrated in a placebo-controlled crossover study in healthy volunteers. All subjects received Glipizide alone and following treatment with 100 mg of Diflucan® as a single daily oral dose for 7 days. The mean percentage increase in the glipizide AUC after fluconazole administration was 56.9% (range: 35 to 81%). [see *Drug Interactions* (7.3)]

### **Colesevelam**

Colesevelam can reduce the maximum plasma concentration and total exposure of glipizide when the two are coadministered. In studies assessing the effect of colesevelam on the pharmacokinetics of glipizide ER in healthy volunteers, reductions in glipizide AUC<sub>0-∞</sub> and C<sub>max</sub> of 12% and 13%, respectively were observed when colesevelam was coadministered with glipizide ER. When glipizide ER was administered 4 hours prior to colesevelam, there was no significant change in glipizide AUC<sub>0-∞</sub> or C<sub>max</sub>, -4% and 0%, respectively. [see *Drug Interactions* (7.4)]

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

A twenty month study in rats and an eighteen month study in mice at doses up to 75 times the maximum human dose revealed no evidence of drug-related carcinogenicity. Bacterial and *in vivo* mutagenicity tests were uniformly negative. Studies in rats of both sexes at doses up to 20 times the human dose based on body surface area showed no effects on fertility.

## **15 REFERENCES**

1. **1. *Diabetes*, 19, SUPP. 2: 747-830, 1970**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

Glipizide Extended-release Tablets USP, 2.5 mg are yellow colored, round, biconvex film-coated tablets imprinted with "2" on one side with black ink and plain on the other side and are supplied as follows:

NDC 16714-894-01 in bottle of 30 tablets with child-resistant closure

Glipizide Extended-release Tablets USP, 5 mg are orange colored, round, biconvex film-coated tablets imprinted with "3" on one side with black ink and plain on the other side and are supplied as follows:

NDC 16714-895-01 in bottle of 100 tablets with child-resistant closure

NDC 16714-895-02 in bottle of 500 tablets

Glipizide Extended-release Tablets USP, 10 mg are white colored, round, biconvex film-coated tablets imprinted with "4" on one side and plain on the other side and are supplied as follows:

NDC 16714-896-01 in bottle of 100 tablets with child-resistant closure

NDC 16714-896-02 in bottle of 500 tablets

### **Storage:**

The tablets should be protected from moisture and humidity.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

## **17 PATIENT COUNSELING INFORMATION**

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Inform patients of the potential adverse reactions of glipizide extended-release tablets including hypoglycemia. Explain the risks of hypoglycemia, its symptoms and treatment, and conditions that predispose to its development to patients and responsible family members. Also inform patients about the importance of adhering to dietary instructions, of a regular exercise program, and of regular testing of glycemic control.

Inform patients that glipizide extended-release tablets should be swallowed whole. Inform patients that they should not chew, divide or crush tablets.

### ***Pregnancy***

Advise females of reproductive potential to inform their prescriber of a known or suspected pregnancy ***[see Use in Specific Populations (8.1)]***.

### ***Lactation***

Advise breastfeeding women taking glipizide extended-release tablets to monitor breastfed infants for signs of hypoglycemia (e.g., jitters, cyanosis, hypothermia, excessive sleepiness, poor feeding, seizures) ***[see Use in Specific Populations (8.2)]***.

The 2.5 mg tablets contain FD&C Yellow No.5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No.5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

Trademarks are the property of their respective owners.

### **Manufactured for:**

Northstar Rx LLC

Memphis, TN 38141.

### **Manufactured by:**

Zydus Lifesciences Ltd.

Ahmedabad, India.

Rev.: 12/25

## **PATIENT INFORMATION**

### **Glipizide (GLIP-i-zide) Extended-release Tablets, USP**

The 2.5 mg tablets contain FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

## **What are glipizide extended-release tablets?**

- Glipizide extended-release tablets are a prescription medicine you take by mouth used along with diet and exercise to lower blood sugar in adults with type 2 diabetes mellitus.
- Glipizide extended-release tablets are not for people with type 1 diabetes or people with diabetic ketoacidosis.

It is not known if glipizide extended-release tablets are safe and effective in children under 18 years of age.

## **Who should not take glipizide extended-release tablets?**

### **Do not use glipizide extended-release tablets if you:**

- have a condition called diabetic ketoacidosis
- have ever had an allergic reaction to glipizide or any of the other ingredients in glipizide extended-release tablets. See the end of this Patient Information for a complete list of ingredients in glipizide extended-release tablets.

## **What should I tell my doctor before taking glipizide extended-release tablets?**

### **Before you take glipizide extended-release tablets, tell your healthcare provider if you:**

- Have ever had a condition called diabetic ketoacidosis
- Have kidney or liver problems
- Have had a blockage or narrowing of your intestines due to illness or past surgery
- Have chronic (continuing) diarrhea
- Have glucose-6-phosphate dehydrogenase (G6PD) deficiency. This condition usually runs in families. People with G6PD deficiency who take glipizide extended-release tablets may develop hemolytic anemia (fast breakdown of red blood cells).
- Are pregnant or might be pregnant. It is not known if glipizide extended-release tablets will harm your unborn baby. If you are pregnant, talk to your healthcare provider about the best way to control your blood sugar while you are pregnant. You should not take glipizide extended-release tablets during the last two weeks of pregnancy.
- Are breastfeeding or plan to breastfeed. It is not known if glipizide passes into your breast milk. You and your healthcare provider should decide the best way to feed your baby during treatment with glipizide extended-release tablets.

**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Glipizide extended-release tablets may affect the way other medicines work, and other medicines may affect how glipizide extended-release tablets works.

Some medicines can affect how well glipizide extended-release tablets works or may affect you blood sugar level. Know the medicines you take. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

## **How should I take glipizide extended-release tablets?**

- Take glipizide extended-release tablets exactly as your healthcare provider tells you to take it.
- Your healthcare provider will tell you how much glipizide extended-release tablets to

take and when to take it.

- Take glipizide extended-release tablets by mouth, 1 time each day with breakfast or your first meal of the day.
- Each glipizide extended-release tablets tablet will release the medicine slowly over 24 hours. This is why you take it only 1 time each day.
- Swallow the glipizide extended-release tablets whole. **Do not break, crush, dissolve, chew, or cut** the tablet in half. This will damage the tablet and release too much medicine into your body at one time.
- It is important to take glipizide extended-release tablets every day to help keep your blood sugar level under good control. Your healthcare provider may change your dose depending on your blood sugar test results. If your blood sugar level is not under control, call your healthcare provider. Do not change your dose unless your healthcare provider tells you to.
- If you take too much glipizide extended-release tablets, call your healthcare provider or go to the nearest emergency room right away.
- Your healthcare provider may tell you to take glipizide extended-release tablets with other diabetes medicines. Low blood sugar can happen more often when glipizide extended-release tablets is taken with other diabetes medicines. See "**What are the possible side effects of glipizide extended-release tablets?**"
- Check your blood sugar as your healthcare provider tells you to.
- Stay on your prescribed diet and exercise program while taking glipizide extended-release tablets.

### **What should I avoid while taking glipizide extended-release tablets?**

- Do not drink alcohol while taking glipizide extended-release tablets. It can increase your chances of getting serious side effects.
- Do not drive, operate machinery, or do other dangerous activities until you know how glipizide extended-release tablets affects you.

### **What are the possible side effects of glipizide extended-release tablets?**

#### **Glipizide extended-release tablets can cause serious side effects, including:**

- **Low blood sugar.** Glipizide extended-release tablets may cause low blood sugar. Signs and symptoms of low blood sugar may include:
  - a cold clammy feeling
  - hunger
  - unusual sweating
  - fast heartbeat
  - dizziness
  - headache
  - weakness
  - blurred vision
  - trembling
  - slurred speech
  - shakiness
  - tingling in the lips or hands

If you have signs or symptoms of low blood sugar, eat or drink something with sugar in it right away. If you do not feel better or your blood sugar level does not go up, call your healthcare provider or go to the nearest emergency room.

**The most common side effects of glipizide extended-release tablets include:** dizziness, diarrhea, nervousness, tremor, and gas.

These are not all the possible side effects of glipizide extended-release tablets. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How to store glipizide extended-release tablets?**

- Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].
- Store glipizide extended-release tablets in a dry place, in its original container.
- Drug glipizide extended-release tablets come in child-resistant bottles of 30s and 100s.
- **Keep glipizide extended-release tablets and all medicines out of reach of children.**

**General information about the safe and effective use of glipizide extended-release tablets** Medicines are sometimes prescribed for purposes other than those listed in a patient information leaflet. Do not use glipizide extended-release tablets for a condition for which it was not prescribed. Do not give glipizide extended-release tablets to other people, even if they have the same symptoms you have. It may harm them.

This Patient Information summarizes the most important information about glipizide extended-release tablets. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about glipizide extended-release tablets that is written for healthcare professionals.

Please address all medical inquiries to, Northstar Rx LLC at Tel.: 1-800-206-7821.

**What are the ingredients in glipizide extended-release tablets?**

**Active ingredient:** Glipizide

**Inactive ingredients:** acetyltributyl citrate, colloidal silicon dioxide, hydroxyethyl cellulose, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, methacrylic acid copolymer and polyethylene glycol. Additionally each 2.5 mg tablet contains: FD&C yellow #5 aluminum lake and titanium dioxide. Each 5 mg tablet contains: FD&C yellow #6 aluminum lake and titanium dioxide. The tablet is imprinted with opacode black S-1-17823 which contains following ingredients: ammonium hydroxide, iron oxide black, isopropyl alcohol, n-butyl alcohol, propylene glycol and shellac.

This Patient Information has been approved by the U.S. Food and Drug Administration

**Manufactured for:**

Northstar Rx LLC

Memphis, TN 38141.

**Manufactured by:**

Zydus Lifesciences Ltd.

Ahmedabad, India.

Rev.: 12/25


## PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 16714-894-01 in bottle of 30 tablets

Glipizide Extended-release Tablets USP, 2.5 mg

R<sub>x</sub> only

30 tablets




NDC 16714-895-01 in bottle of 100 tablets


Glipizide Extended-release Tablets USP, 5 mg

R<sub>x</sub> only

100 tablets



NDC 16714-896-01 in bottle of 100 tablets  
 Glipizide Extended-release Tablets USP, 10 mg  
 Rx only  
 100 tablets



## GLIPIZIDE

glipizide tablet, extended release

### Product Information

#### Product Type

HUMAN PRESCRIPTION DRUG

#### Item Code (Source)

NDC:16714-894

|                                                                           |                                          |                                                   |                      |                    |
|---------------------------------------------------------------------------|------------------------------------------|---------------------------------------------------|----------------------|--------------------|
| Route of Administration                                                   |                                          | ORAL                                              |                      |                    |
|                                                                           |                                          |                                                   |                      |                    |
| Active Ingredient/Active Moiety                                           |                                          |                                                   |                      |                    |
| Ingredient Name                                                           |                                          | Basis of Strength                                 | Strength             |                    |
| GLIPIZIDE (UNII: X7WDT95N5C) (GLIPIZIDE - UNII:X7WDT95N5C)                |                                          | GLIPIZIDE                                         | 2.5 mg               |                    |
|                                                                           |                                          |                                                   |                      |                    |
| Inactive Ingredients                                                      |                                          |                                                   |                      |                    |
| Ingredient Name                                                           |                                          |                                                   | Strength             |                    |
| ACETYLTRIBUTYL CITRATE (UNII: 0ZBX0N59RZ)                                 |                                          |                                                   |                      |                    |
| ALUMINUM OXIDE (UNII: LMI26O6933)                                         |                                          |                                                   |                      |                    |
| AMMONIA (UNII: 5138Q19F1X)                                                |                                          |                                                   |                      |                    |
| BUTYL ALCOHOL (UNII: 8PJ61P6TS3)                                          |                                          |                                                   |                      |                    |
| FD&C YELLOW NO. 5 (UNII: I753WB2F1M)                                      |                                          |                                                   |                      |                    |
| FERROSOFERRIC OXIDE (UNII: XM0M87F357)                                    |                                          |                                                   |                      |                    |
| HYDROXYETHYL CELLULOSE (2000 CPS AT 1%) (UNII: S38J6RZN16)                |                                          |                                                   |                      |                    |
| HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)                   |                                          |                                                   |                      |                    |
| HYPROMELLOSES (UNII: 3NXW29V3WO)                                          |                                          |                                                   |                      |                    |
| ISOPROPYL ALCOHOL (UNII: ND2M416302)                                      |                                          |                                                   |                      |                    |
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)                                    |                                          |                                                   |                      |                    |
| MAGNESIUM STEARATE (UNII: 70097M6I30)                                     |                                          |                                                   |                      |                    |
| METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13) |                                          |                                                   |                      |                    |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)                       |                                          |                                                   |                      |                    |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3)                                       |                                          |                                                   |                      |                    |
| SHELLAC (UNII: 46N107B71O)                                                |                                          |                                                   |                      |                    |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)                                        |                                          |                                                   |                      |                    |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)                                       |                                          |                                                   |                      |                    |
|                                                                           |                                          |                                                   |                      |                    |
| Product Characteristics                                                   |                                          |                                                   |                      |                    |
| Color                                                                     | YELLOW (Yellow)                          | Score                                             | no score             |                    |
| Shape                                                                     | ROUND (Round)                            | Size                                              | 6mm                  |                    |
| Flavor                                                                    |                                          | Imprint Code                                      | 2                    |                    |
| Contains                                                                  |                                          |                                                   |                      |                    |
|                                                                           |                                          |                                                   |                      |                    |
| Packaging                                                                 |                                          |                                                   |                      |                    |
| #                                                                         | Item Code                                | Package Description                               | Marketing Start Date | Marketing End Date |
| 1                                                                         | NDC:16714-894-01                         | 30 in 1 BOTTLE; Type 0: Not a Combination Product | 12/01/2018           |                    |
|                                                                           |                                          |                                                   |                      |                    |
| Marketing Information                                                     |                                          |                                                   |                      |                    |
| Marketing Category                                                        | Application Number or Monograph Citation |                                                   | Marketing Start Date | Marketing End Date |
| ANDA                                                                      | ANDA203499                               |                                                   | 12/01/2018           |                    |

# GLIPIZIDE

glipizide tablet, extended release

## Product Information

|                         |                         |                    |               |
|-------------------------|-------------------------|--------------------|---------------|
| Product Type            | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:16714-895 |
| Route of Administration | ORAL                    |                    |               |

## Active Ingredient/Active Moiety

| Ingredient Name                                            | Basis of Strength | Strength |
|------------------------------------------------------------|-------------------|----------|
| GLIPIZIDE (UNII: X7WDT95N5C) (GLIPIZIDE - UNII:X7WDT95N5C) | GLIPIZIDE         | 5 mg     |

## Inactive Ingredients

| Ingredient Name                                                           | Strength |
|---------------------------------------------------------------------------|----------|
| ACETYLTRIBUTYL CITRATE (UNII: 0ZBX0N59RZ)                                 |          |
| AMMONIA (UNII: 5138Q19F1X)                                                |          |
| BUTYL ALCOHOL (UNII: 8PJ61P6TS3)                                          |          |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8)                                      |          |
| FERROSOFERRIC OXIDE (UNII: XM0M87F357)                                    |          |
| HYDROXYETHYL CELLULOSE (2000 CPS AT 1%) (UNII: S38J6RZN16)                |          |
| HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)                   |          |
| HYPROMELLOSES (UNII: 3NXW29V3WO)                                          |          |
| ISOPROPYL ALCOHOL (UNII: ND2M416302)                                      |          |
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)                                    |          |
| MAGNESIUM STEARATE (UNII: 70097M6I30)                                     |          |
| METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13) |          |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)                       |          |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3)                                       |          |
| SHELLAC (UNII: 46N107B71O)                                                |          |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)                                        |          |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP)                                       |          |

## Product Characteristics

|          |                 |              |          |
|----------|-----------------|--------------|----------|
| Color    | ORANGE (Orange) | Score        | no score |
| Shape    | ROUND (Round)   | Size         | 6mm      |
| Flavor   |                 | Imprint Code | 3        |
| Contains |                 |              |          |

## Packaging

| # | Item Code        | Package Description                                | Marketing Start Date | Marketing End Date |
|---|------------------|----------------------------------------------------|----------------------|--------------------|
| 1 | NDC:16714-895-01 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 12/01/2018           |                    |

|                       |                                          |                                                    |                    |  |
|-----------------------|------------------------------------------|----------------------------------------------------|--------------------|--|
| 2                     | NDC:16714-895-02                         | 500 in 1 BOTTLE; Type 0: Not a Combination Product | 12/01/2018         |  |
|                       |                                          |                                                    |                    |  |
| Marketing Information |                                          |                                                    |                    |  |
| Marketing Category    | Application Number or Monograph Citation | Marketing Start Date                               | Marketing End Date |  |
| ANDA                  | ANDA203499                               | 12/01/2018                                         |                    |  |

|                                                                           |                         |                    |               |
|---------------------------------------------------------------------------|-------------------------|--------------------|---------------|
| GLIPIZIDE                                                                 |                         |                    |               |
| glipizide tablet, extended release                                        |                         |                    |               |
|                                                                           |                         |                    |               |
| Product Information                                                       |                         |                    |               |
| Product Type                                                              | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:16714-896 |
| Route of Administration                                                   | ORAL                    |                    |               |
|                                                                           |                         |                    |               |
| Active Ingredient/Active Moiety                                           |                         |                    |               |
| Ingredient Name                                                           |                         | Basis of Strength  | Strength      |
| GLIPIZIDE (UNII: X7WDT95N5C) (GLIPIZIDE - UNII:X7WDT95N5C)                |                         | GLIPIZIDE          | 10 mg         |
|                                                                           |                         |                    |               |
| Inactive Ingredients                                                      |                         |                    |               |
| Ingredient Name                                                           |                         |                    | Strength      |
| ACETYLTRIBUTYL CITRATE (UNII: 0ZBX0N59RZ)                                 |                         |                    |               |
| AMMONIA (UNII: 5138Q19F1X)                                                |                         |                    |               |
| BUTYL ALCOHOL (UNII: 8PJ61P6TS3)                                          |                         |                    |               |
| FERROSOFERRIC OXIDE (UNII: XM0M87F357)                                    |                         |                    |               |
| HYDROXYETHYL CELLULOSE (2000 CPS AT 1%) (UNII: S38J6RZN16)                |                         |                    |               |
| HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)                   |                         |                    |               |
| HYPROMELLOSES (UNII: 3NXW29V3WO)                                          |                         |                    |               |
| ISOPROPYL ALCOHOL (UNII: ND2M416302)                                      |                         |                    |               |
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)                                    |                         |                    |               |
| MAGNESIUM STEARATE (UNII: 70097M6I30)                                     |                         |                    |               |
| METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13) |                         |                    |               |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)                       |                         |                    |               |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3)                                       |                         |                    |               |
| SHELLAC (UNII: 46N107B71O)                                                |                         |                    |               |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)                                        |                         |                    |               |
|                                                                           |                         |                    |               |
| Product Characteristics                                                   |                         |                    |               |
| Color                                                                     | WHITE (White)           | Score              | no score      |
| Shape                                                                     | ROUND (Round)           | Size               | 6mm           |
| Flavor                                                                    |                         | Imprint Code       | 4             |
| Contains                                                                  |                         |                    |               |

**Packaging**

| # | Item Code        | Package Description                                | Marketing Start Date | Marketing End Date |
|---|------------------|----------------------------------------------------|----------------------|--------------------|
| 1 | NDC:16714-896-01 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 12/01/2018           |                    |
| 2 | NDC:16714-896-02 | 500 in 1 BOTTLE; Type 0: Not a Combination Product | 12/01/2018           |                    |

**Marketing Information**

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|------------------------------------------|----------------------|--------------------|
| ANDA               | ANDA203499                               | 12/01/2018           |                    |

**Labeler** - Northstar Rx LLC. (830546433)**Registrant** - Zydus Pharmaceuticals USA Inc. (156861945)**Establishment**

| Name                       | Address | ID/FEI    | Business Operations                                                                      |
|----------------------------|---------|-----------|------------------------------------------------------------------------------------------|
| Zydus Lifesciences Limited |         | 918596198 | ANALYSIS(16714-894, 16714-895, 16714-896) , MANUFACTURE(16714-894, 16714-895, 16714-896) |

Revised: 1/2026

Northstar Rx LLC.