ADULT LOW STRENGTH ASPIRIN- aspirin tablet, delayed release DISCOUNT DRUG MART

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1146-DDM-2021-1126

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID ¹)

1 nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

• temporarily relieves minor aches and pains

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours; do not exceed 48 tablets in 24 hours
- children under 12 years: do not use unless directed by a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

acetylated monoglycerides, anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, FD&C red #40, FD&C yellow #6, hypromellose, hypromellose phthalate, iron oxide yellow, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polyethylene glycol, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

PRINCIPAL DISPLAY PANEL

DISCOUNT drug mart FOOD FAIR

†COMPARE TO THE ACTIVE INGREDIENT IN ST. JOSEPH® ADULT LOW STRENGTH ASPIRIN

ADULT LOW STRENGTH ASPIRIN

SAFETY COATED PAIN RELIEVER (NSAID)

SEE NEW WARNINGS INFORMATION

180 TABLETS - 81 mg each

81mg

Enteric Coated



ADULT LOW STRENGTH ASPIRIN

aspirin tablet, delayed release

Product Information					
Product Type HUMAN OTC DRUG Item Code (Source)			(Source)	NDC:53943-146	
Route of Administration	ORAL				
Active Ingredient/Active Moiety					
Ingredient Name			Basis of Strength		Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPI	RIN - UNII:R16CO5Y76E)	AS	SPIRIN		81 mg
Inactive Ingredients					

Ingredient Name	Strength
DIACETYLATED MONOGLYCERIDES (UNII: 5Z17386USF)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C RED NO. 40 (UNII: WZ B9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
HYPROMELLOSE PHTHALATE (24% PHTHALATE, 55 CST) (UNII: 87Y6436BKR)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 60ZP39ZG8H)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	orange (PEACH)	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	heart
Contains			

Packaging

#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943- 146-01	1 in 1 CARTON	06/25/2010	
1		180 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:53943- 146-02	1 in 1 CARTON	06/25/2010	
2		300 in 1 PACKAGE; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part343	06/25/2010		

Revised: 11/2021

DISCOUNT DRUG MART