

CABINET COUGH RELIEF- dextromethorphan hydrobromide capsule, liquid filled

Spirit Pharmaceuticals LLC

Cabinet Cough Relief

Drug Facts

Active ingredient (in each softgel)

Dextromethorphan HBr, USP 15mg

Purpose

Cough suppressant

Use

temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold.

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if you have

- a cough that occurs with too much phlegm (mucus)
- a cough that lasts or is chronic as occurs with smoking, asthma, or emphysema

Stop use and ask doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than 8 capsules in any 24-hour period
- this adult product is not intended for use in children under 12 years of age

take 2 capsules

adults and children 12 years and over	take 2 capsules every 6 to 8 hours, as needed
children under 12 years	do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)
- protect from light

Inactive ingredients

FD&C Red No. 40, gelatin, glycerin, polyethylene glycol 400, propylene glycol, povidone, sorbitol sorbitan solution, titanium dioxide

Questions or comments?

1-888-333-9792

Distributed by:
Cabinet Health, Inc.
wearecabinet.com
Made in India

PRINCIPAL DISPLAY PANEL

Compare to the active ingredient
in Robitussin® Long Acting Cough Gels®*

CABINET:

Cough Relief

Dextromethorphan HBr 15mg

50 softgels



Distributed by:
Cabinet Health, Inc.
www.cabinet.com
Made in India

Compare to the active ingredient
in Robitussin® Long Acting Cough Gels*

CABINET® Cough Relief

Dextromethorphan HBr 15mg

50 Softgels

Tamper evident: do not use if print safety seal under cap
is broken or missing

Drug Facts

Active ingredient
(in each softgel) Dextromethorphan HBr 15 mg

Purpose
Cough suppressant

Drug Facts (continued under label)

Lot:
Exp:



Drug Facts (continued)

Do not use if you are on a prescription drug for pain, or if you are taking any other prescription drug for pain, or if you are taking any other prescription drug for cough.

Warnings
Do not use if you are on a prescription drug for pain, or if you are taking any other prescription drug for pain, or if you are taking any other prescription drug for cough. Do not use if you are on a prescription drug for depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
any of the following conditions: asthma, emphysema, or other lung disease; a history of seizures; a history of heart disease; a history of high blood pressure; a history of kidney disease; a history of liver disease; a history of stomach or intestinal disease; a history of urinary tract infection; a history of diabetes; a history of thyroid disease; a history of adrenal gland disease; a history of pheochromocytoma; a history of paraganglioma; a history of paraganglioid; a history of paraganglioma; a history of paraganglioid; a history of paraganglioma; a history of paraganglioid.

Stop use and ask a doctor if you have
any of the following symptoms: dizziness, lightheadedness, or fainting; a change in your voice; a change in your breathing; a change in your heart rate; a change in your blood pressure; a change in your blood sugar; a change in your thyroid function; a change in your adrenal gland function; a change in your paraganglioma function; a change in your paraganglioid function.

Directions • Do not take more than 8 softgels in any 24-hour period. • This adult softgel is not intended for use in children under 12 years of age.

Other information
• Store at 20-25°C (68-77°F). • Avoid excessive heat above 30°C (86°F). • Protect from light.

Inactive ingredients
DSCC, PEG 400, polyethylene glycol 400, polyethylene glycol 600, polyethylene glycol 800, polyethylene glycol 1000, polyethylene glycol 1500, polyethylene glycol 2000, polyethylene glycol 3000, polyethylene glycol 4000, polyethylene glycol 6000, polyethylene glycol 8000, polyethylene glycol 10000, polyethylene glycol 15000, polyethylene glycol 20000, polyethylene glycol 30000, polyethylene glycol 40000, polyethylene glycol 60000, polyethylene glycol 80000, polyethylene glycol 100000, polyethylene glycol 150000, polyethylene glycol 200000, polyethylene glycol 300000, polyethylene glycol 400000, polyethylene glycol 600000, polyethylene glycol 800000, polyethylene glycol 1000000.

Questions or comments?
1-888-333-9792

*This product is not manufactured or distributed by Phazax, Inc., owner of the registered trademark Robitussin® Long Acting Cough Gels®.

Phazax, Inc. is the owner of the registered trademark Robitussin® Long Acting Cough Gels®.

Lot:
Exp:

Compare to the active ingredient in
Robitussin® Long Acting Cough Gels®

CABINET® Cough Relief

Dextromethorphan HBr 15mg

25 Softgels

Tamper evident: do not use if print safety seal under cap is broken or missing

Uses temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold

Drug Facts (continued under label)



Warnings

Do not use if you are on a new taking prescription drug for pain, or if you are taking any other prescription drug for pain, or if you are taking any other prescription drug for cough. Do not use if you are on a prescription drug for depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have a cough that occurs with too much phlegm (mucus) or a cough that lasts or is chronic as occurs with smoking, asthma, or emphysema

Stop use and ask a doctor if you have any of the following symptoms: dizziness, lightheadedness, or fainting; a change in your voice; a change in your breathing; a change in your heart rate; a change in your blood pressure; a change in your blood sugar; a change in your thyroid function; a change in your adrenal gland function; a change in your paraganglioma function; a change in your paraganglioid function.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Do not use if you are on a prescription drug for pain, or if you are taking any other prescription drug for pain, or if you are taking any other prescription drug for cough. Do not use if you are on a prescription drug for depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Directions • Do not take more than 8 softgels in any 24-hour period. • This adult product is not intended for use in children under 12 years of age.

adults & children 12 years & over	take 2 softgels every 6 to 8 hours
children under 12 years of age	do not use


Other information
• Store at 20-25°C (68-77°F). • Avoid excessive heat above 30°C (86°F). • Protect from light.

Questions or comments?
1-888-333-9792

*This product is not manufactured or distributed by Phazax, Inc., owner of the registered trademark Robitussin® Long Acting Cough Gels®.

Phazax, Inc. is the owner of the registered trademark Robitussin® Long Acting Cough Gels®.

Distributed by: Cabinet Health, Inc. www.cabinet.com Made in India



CABINET COUGH RELIEF

dextromethorphan hydrobromide capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68210-6010
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII: 7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	

WATER (UNII: 059QF0KO0R)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
SORBITOL (UNII: 506T60A25R)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	13mm
Flavor		Imprint Code	404
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68210-6010-3	1 in 1 CARTON	10/28/2019	
1		25 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:68210-6010-2	50 in 1 PACKAGE; Type 0: Not a Combination Product	10/28/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/28/2019	

Labeler - Spirit Pharmaceuticals LLC (179621011)