

**DG HEALTH DAYTIME SEVERE COLD AND FLU- acetaminophen,
dextromethorphan hydrobromide, guaifenesin, phenylephrine
hydrochloride capsule
Dolgenercorp Inc**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DayTime SEVERE Cold & Flu Relief

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purposes

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • minor aches & pains • headache • fever • sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 8 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may

include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin; taking sedatives or tranquilizers.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- do not exceed 8 softgels per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hours
children 4 to under	ask a doctor

12 yrs	
children under 4 yrs	do not use

Other information

- store at 20-25°C (68-77°F) • protect from light, heat and moisture

Inactive ingredients

edible printing ink, FD&C blue no. 1, FD&C red no. 40, gelatin, glycerin, polyethylene glycol 400, povidone K30, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments?

Call **1-888-577-8033** Monday - Friday 8am - 4pm EST

COMPARE TO VICKS® DAYQUIL® SEVERE COLD & FLU ACTIVE INGREDIENTS* MULTI-SYMP TOM RELIEF

- Chest Congestion, Thins and Loosens Mucus

Antihistamine Free

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.


*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademarks Vicks® and DayQuil™.

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

Distributed by:
Strive Pharmaceuticals Inc.
East Brunswick, NJ 08816

Product of United Arab Emirates

Packaging



Day Time

SEVERE Cold & Flu Relief

PAIN RELIEVER-FEVER REDUCER - ACETAMINOPHEN
COUGH SUPPRESSANT - DEXTROMETHORPHAN HBr
EXPECTORANT - GUAIFENESIN
NASAL DECONGESTANT - PHENYLEPHRINE HCl

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Chest Congestion, Thins and Loosens Mucus
- Nasal Congestion, Sinus Pressure
- Cough

Antihistamine Free



actual size

24 SOFTGELS

KEEP OUTER CARTON FOR COMPLETE WARNING

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 8 softgels in 24 hours, which is the maximum daily amount for this product
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

taking the blood thinning drug warfarin; taking sedatives or tranquilizers.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- do not exceed 8 softgels per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hours
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Drug Facts (continued)

Other information ■ store at 20-25°C (68-77°F) ■ protect from light, heat and moisture

Inactive ingredients ■ edible printing ink, FD&C blue no. 1, FD&C red no. 40, gelatin, glycerin, polyethylene glycol, purified water, sorbitol sorbitan solution

Questions or comments? Call 1-800-577-8003 Monday - Friday 8am - 4pm EST

*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademarks Vicksi[®] and Day/Cal

Distributed by:
Shive Pharmaceuticals Inc.
East Brunswick, NJ 08816

Product of
United Arab Emirates



LOT: COATING FREE AREA
EXP:

DRUG FACTS TABLE

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Purposes

Drug Facts (continued)

Uses

- temporarily relieves common cold/flu symptoms:
 - nasal congestion
 - sinus congestion & pressure
 - cough due to minor throat & bronchial irritation
 - minor aches & pains
 - headache
 - fever
 - sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 8 softgels in 24 hours, which is the maximum daily amount for this product
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if

Drug Facts (continued)

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin; taking sedatives or tranquilizers.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- do not exceed 8 softgels per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hours
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

Other information ■ store at 20-25°C (68-77°F) ■ protect from light, heat and moisture

Inactive ingredients edible printing ink, FD&C blue no. 1, FD&C red no. 40, gelatin, glycerin, polyethylene glycol 400, povidone K30, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments? Call 1-888-577-8033 Monday - Friday 8am - 4pm EST

DG HEALTH DAYTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride capsule

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-285
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE K30 (UNII: U725QWY32X)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL (oblong)	Size	20mm
Flavor		Imprint Code	811
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-285-24	2 in 1 CARTON	08/08/2023	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/08/2023	

Labeler - Dolgencorp Inc (068331990)

Revised: 8/2023

Dolgencorp Inc