

CHLORPHENIRAMINE MALEATE 4MG- chlorpheniramine maleate 4mg tablet
Strive Pharmaceuticals Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Allergy - Alergia (Chlorpheniramine Maleate 4mg)

Chlorpheniramine Maleate 4mg

D&C yellow #10, lactose anhydrous, magnesium stearate, microcrystalline cellulose, pregelatinized starch

- take with a full glass of water
- do not take more than directed

adults and children 12 yrs of age and over	1 tablet every 4 to 6 hours, do not take more than 6 tablets in 24 hours
children 6 to under 12 years of age	1/2 tablet (break tablet in half) every 4 to 6 hours, not to exceed 3 whole tablets in 24 hours
children under 6 years of age	ask a doctor

temporarily relieves the following symptoms due to hay fever or other upper respiratory allergies

- sneezing
- runny nose
- itchy, watery eyes
- itching of the nose or throat

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or a pharmacist before use if

you are taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- drowsiness may occur
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating a machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

If pregnant or breast-feeding,

ask a health professional before use

Keep out of reach of children


Overdose warning: In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children

Overdose Warning: In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Antihistamine

- store at room temperature between 20 °C - 25 °C (68 °F - 77 °F)
- protect from excessive heat and moisture



		LOT: EXP:
	<p>DRUG FACTS</p> <p>Active ingredient (in each tablet) Chlorpheniramine Maleate 4 mg</p> <p>Purpose Antihistamine</p> <p>Uses:</p> <ul style="list-style-type: none"> temporarily relieves the following symptoms due to hay fever or other upper respiratory allergies: sneezing runny nose itchy, watery eyes itching of the nose or throat <p>Warnings:</p> <p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> a breathing problem such as emphysema or chronic bronchitis glaucoma trouble urinating due to an enlarged prostate gland <p>Ask a doctor or a pharmacist before use if you are taking sedatives or tranquilizers.</p> <p>When using this product:</p> <ul style="list-style-type: none"> excitability may occur, especially in children drowsiness may occur avoid alcoholic beverages use caution when driving a motor vehicle or operating a machinery alcohol, sedatives, and tranquilizers may increase drowsiness <p style="text-align: right;">LIFT PANEL FOR CONTINUED DRUG FACTS</p>	

<p>DRUG FACTS (Continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p>Overdose Warning: In case of accidental overdose, get medical help or contact a Poison Control Center, right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions:</p> <ul style="list-style-type: none"> Take with a full glass of water. do not take more than directed. <table border="1" style="width: 100%;"> <tr> <td>adults and children 12 years of age and over:</td> <td>1 tablet every 4 to 6 hours, do not take more than 6 tablets in 24 hours</td> </tr> <tr> <td>children 6 to under 12 years of age:</td> <td>½ tablet (break tablet in half) every 4 to 6 hours, not to exceed 3 whole tablets in 24 hours</td> </tr> <tr> <td>children under 6 years of age:</td> <td>ask a doctor</td> </tr> </table> <p>Other information:</p> <ul style="list-style-type: none"> Store at room temperature between 20-25°C (68-77°F). protect from excessive heat and moisture <p>Inactive ingredients: D1AC, yellow 10, lactose anhydrous, magnesium stearate, microcrystalline cellulose, pregelatinized starch</p>	adults and children 12 years of age and over:	1 tablet every 4 to 6 hours, do not take more than 6 tablets in 24 hours	children 6 to under 12 years of age:	½ tablet (break tablet in half) every 4 to 6 hours, not to exceed 3 whole tablets in 24 hours	children under 6 years of age:	ask a doctor	<p>DRUG FACTS (Continued)</p> <p>QUESTIONS OR COMMENTS? 1-833-685-9444</p> <p>READ AND KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION</p> <p>Distributed By: O4 GLOBAL TRADING USA LLC, 2241 S Watson Rd, Suite 121 Arlington, TX 76010 www.onlyfor.com Packaged By: Sinite Pharmaceuticals Inc., East Brunswick, NJ 08818</p> <p>TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF A BUBBLER UNIT IS TORN, BROKEN OR SHOWS ANY SIGN OF TAMPERING</p> <div style="text-align: center;"> <p>Tear at arrows push through</p> </div> <p>www.onlyfor.com CT2ALR4-REV.102018</p>
adults and children 12 years of age and over:	1 tablet every 4 to 6 hours, do not take more than 6 tablets in 24 hours						
children 6 to under 12 years of age:	½ tablet (break tablet in half) every 4 to 6 hours, not to exceed 3 whole tablets in 24 hours						
children under 6 years of age:	ask a doctor						

chlorpheniramine maleate 4 mg tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-113
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	S72
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-113-41	4 in 1 PACKAGE; Type 0: Not a Combination Product	12/20/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/20/2018	

Labeler - Strive Pharmaceuticals Inc (080028013)