

CHILDRENS IBUPROFEN- ibuprofen suspension
Kroger Company

Children's Ibuprofen

Drug Facts

Active ingredient (in each 5 mL)

Ibuprofen 100 mg (NSAID) ¹

¹ nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin,

ibuprofen, naproxen, or others)

- takes more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning

Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

- the child does not get any relief within first day (24 hours) of treatment
- fever or pain get worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6 to 8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL) *
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- **Store at 20° to 25°C (68° to 77°F)**[see USP Controlled Room Temperature].
- **do not use if carton is opened or if bottle wrap imprinted with "SEALED FOR SAFETY" is broken or missing**
- see bottom panel of carton for lot number and expiration date

Inactive ingredients

acesulfame potassium, berry flavor natural & artificial, citric acid anhydrous, glycerin, polysorbate 80, pregelatinized modified starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions?

1-800-632-6900

DIST. BY THE KROGER CO., CINCINNATI, OHIO 45202

PRINCIPAL DISPLAY PANEL - 120 mL Bottle Carton

NDC 30142-965-08

Kroger®

Pharmacist Read Direction
→

NO VARNISH
ON THIS FLAP

NO VARNISH
ON THIS FLAP

T181C
B98.2
ENG19.59

NO VARNISH
ON THIS FLAP

5254617
1124
43

NO VARNISH/ NO AQ
NO COPY / NO COLOR
THIS FLAP FOR LOT #
AND EXP DATE PRINT



0 41260 01288 4

Kroger health

CHILDREN'S
Dye-Free
Ibuprofen

Oral Suspension, USP
100 mg per 5 mL
Pain Reliever / Fever Reducer (NSAID)
ALCOHOL FREE

LASTS UP TO 8 HOURS

AGES
2 to 11
YEARS

Kroger health

CHILDREN'S
Dye-Free
Ibuprofen

Oral Suspension, USP
100 mg per 5 mL
Pain Reliever /
Fever Reducer
(NSAID)

AGES
2 to 11
YEARS

LASTS UP TO
8 HOURS

4 FL. OZ.
(120 mL)

BERRY FLAVOR

Drug Facts (continued)

Ask a doctor before use if:

- stomach bleeding warning appears to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is:

- under a doctor's care for any serious condition
- taking any other drug

When taking this product:

- take with food or milk. If stomach upset occurs

Stop use and ask a doctor if:

- child experiences any of the following signs of stomach bleeding:
 - black stool
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- this product does not contain salicylates or aspirin-like warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.

Drug Facts (continued)

- use only enclosed dosing cup. Do not use any other dosing devices.
- if needed, repeat dose every 6 to 8 hours
- do not use more than 4 doses a day
- replace original bottle cap to maintain child resistance

Weight (lb)	Age (yr)	Dose (mL)*
Under 24	Under 2 years	ask a doctor
24-35 lb	2-3 years	5 mL
36-47 lb	4-5 years	7.5 mL
48-59 lb	6-8 years	10 mL
60-71 lb	9-10 years	12.5 mL
72-85 lb	11 years	15 mL

* or as directed by a doctor

Other Information

- Store at 20° to 25° (68° to 77°F) (see USP Controlled Room Temperature).
- do not use if cap is opened or if bottle wrap impervious with "SEALED FOR YOUR PROTECTION" is broken or missing
- see bottom panel of carton for lot number and expiration date

Inactive Ingredients

saccharin potassium, berry flavor natural & artificial, citric acid anhydrous, glycerin, polyorbital 80, propylchlorid modified starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
Call 1-888-632-6900

Motrin® is a registered trademark of Kenova Inc. PROUDLY DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202

MADE IN CANADA

For More Product Information, Scan UPC Using Your Kroger App or Call 800-632-6900

*OUR PHARMACIST RECOMMENDED

Our Brands. Our Guarantee.
Love It or Your Money Back.
www.kroger.com

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active Ingredient (for each 5 mL)
Ibuprofen 100 mg (NSAID) ... Pain reliever/fever reducer (nonsteroidal anti-inflammatory drug)

Purpose

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin rash/itching
- nausea
- blister

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinner (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warnings: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Severe throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, rash, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or as directed in children under 3 years of age unless directed by doctor.

Do not use:

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

CHILDRENS IBUPROFEN

ibuprofen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-965
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
WATER (UNII: 059QF0KO0R)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	white (WHITE to OFF-WHITE)	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-965-08	1 in 1 CARTON	05/14/2021	
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209207	06/27/2017	

Labeler - Kroger Company (006999528)**Establishment**

Name	Address	ID/FEI	Business Operations
Sun Pharma Canada Inc.		243339023	manufacture(30142-965)

Revised: 7/2025

Kroger Company