

SHISEIDO UV PROTECTIVE FOUNDATION LIGHT BEIGE- octinoxate and titanium dioxide cream

SHISEIDO UV PROTECTIVE FOUNDATION LIGHT IVORY- octinoxate and titanium dioxide cream

SHISEIDO UV PROTECTIVE FOUNDATION LIGHT OCHRE- octinoxate and titanium dioxide cream

SHISEIDO UV PROTECTIVE FOUNDATION MEDIUM BEIGE- octinoxate and titanium dioxide cream

SHISEIDO UV PROTECTIVE FOUNDATION MEDIUM IVORY- octinoxate and titanium dioxide cream

SHISEIDO UV PROTECTIVE FOUNDATION DARK BEIGE- octinoxate and titanium dioxide cream

SHISEIDO UV PROTECTIVE FOUNDATION DARK IVORY- octinoxate and titanium dioxide cream

SHISEIDO AMERICAS CORPORATION

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

SHISEIDO UV PROTECTIVE LIQUID FOUNDATION

Drug Facts

<i>Active ingredients</i>	<i>Purpose</i>
OCTINOXATE 1.9%	Sunscreen
TITANIUM DIOXIDE 8.3%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours

- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor

Inactive Ingredients

WATER, DIMETHICONE, TRIMETHYLSILOXYSILICATE, ISODODECANE, POLYMETHYL METHACRYLATE, CYCLOMETHICONE, SD ALCOHOL 40-B, PEG/PPG-19/19 DIMETHICONE, HYDROGENATED POLYISOBUTENE, BIS-BUTYLDIMETHICONE POLYGLYCERYL-3, PEG/PPG-14/7 DIMETHYL ETHER, SAXIFRAGA SARMENTOSA EXTRACT, SODIUM HYALURONATE, SOPHORA ANGUSTIFOLIA ROOT EXTRACT, ALUMINUM HYDROXIDE, ALUMINUM DISTEARATE, ALUMINA, ISOSTEARIC ACID, TRIETHOXYCAPRYLYLSILANE, TRISODIUM EDTA, SILICA, POLYSILICONE-2, ALCOHOL, SODIUM MAGNESIUM SILICATE, HYDRATED SILICA, BUTYLENE GLYCOL, GLYCERIN, TOCOPHEROL, DISTEARYLDIMONIUM CHLORIDE, BHT, SYZYGIUM JAMBOS LEAF EXTRACT, HYDROGEN DIMETHICONE, TIN OXIDE, PHENOXYETHANOL, FRAGRANCE, TITANIUM DIOXIDE, IRON OXIDES, MICA,

Other information

- protect this product in this container from excessive heat and direct sun.

Questions or comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Light Beige

SHI SEIDO

Anti-Aging Foundation

42

UV Protective
Liquid Foundation

BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT

SUNSCREEN

30mL 1 FL. OZ.

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

Light Beige
(SP20)

SHISEIDO

Anti-Aging Foundation



42

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT
SUNSCREEN

30mL 1 FL. OZ.

Drug Facts (continued)

Directions

■ children under 6 months: Ask a doctor

Inactive ingredients

WATER • DIMETHICONE • TRIMETHYLSILOXYSILICATE •
ISODODECANE • POLYMETHYL METHACRYLATE • CYCLOMETHICONE •
SD ALCOHOL 40-8 • PEG/PPG-19/19 DIMETHICONE •
HYDROGENATED POLYISOBUTENE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
PEG/PPG-14/7 DIMETHYL ETHER •
SAXIFRAGA SARMENTOSA EXTRACT • SODIUM HYALURONATE •
SOPHORA ANGUSTIFOLIA ROOT EXTRACT • ALUMINUM HYDROXIDE •
ALUMINUM DISTEARATE • ALUMINA • ISOSTEARIC ACID •
TRIETHOXYCAPRYLYLSILANE • TRISODIUM EDTA • SILICA •
POLYSILICONE-2 • ALCOHOL • SODIUM MAGNESIUM SULFATE •
HYDRATED SILICA • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL •
DISTEARYLDIMONIUM CHLORIDE • BHT •
SYZYGIUM JAMBOS LEAF EXTRACT • HYDROGEN DIMETHICONE •
TIN OXIDE • PHENOXYETHANOL • FRAGRANCE • TITANIUM DIOXIDE •
IRON OXIDES • MICA •

Other Information

■ protect this product in this container
from excessive heat and direct sun. ▼

SHISEIDO
 UV PROTECTIVE LIQUID FOUNDATION
 BROAD SPECTRUM
 SPF 42

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil-UV 360™ technology.
- Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,* cell damage* and DNA damage* linked to the formation of wrinkles and dark spots.
- Provides a fresh touch and a dewy finish.
- NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- * in vitro test

Drug Facts

Active ingredients	Purpose
OCTINOXATE 1.9%	Sunscreen
TITANIUM DIOXIDE 8.3%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin ▶

SHISEIDO AMERICAS CORPORATION DIST.
 NEW YORK, NY 10022
 MADE IN U.S.A.
 GLO. 11199
www.shiseido.com

Drug Facts (continued)

Warnings

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions


For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses ▶

Drug Facts (continued)

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Light Ivory

SHI SEIDO

Anti-Aging Foundation

42

UV Protective
Liquid Foundation

BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT

SUNSCREEN

30mL 1 FL. OZ.

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

Light Ivory

Anti-Aging Foundation



42

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT
SUNSCREEN

30mL 1 FL. OZ.

SHISEIDO

Drug Facts (continued)

Directions

■ children under 6 months: Ask a doctor

Inactive ingredients

WATER • DIMETHICONE • TRIMETHYLSILOXYSILICATE •
DODODECANE • POLYMETHYL METHACRYLATE • CYCLOMETHICONE •
SD ALCOHOL 40-8 • PEG/PPG-19/19 DIMETHICONE •
HYDROGENATED POLYISOBUTENE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
PEG/PPG-14/7 DIMETHYL ETHER •
SAXIFRAGA SARMENTOSA EXTRACT • SODIUM HYALURONATE •
SOPHORA ANGUSTIFOLIA ROOT EXTRACT • ALUMINUM HYDROXIDE •
ALUMINUM DISTEARATE • ALUMINA • ISOSTEARIC ACID •
TRIETHOXYCAPRYLYLSILANE • TRISODIUM EDTA • SILICA •
POLYSILOXANE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE •
HYDRATED SILICA • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL •
DISTEARYLDIMONIUM CHLORIDE • BHT •
SYZYGIUM JAMBOS LEAF EXTRACT • HYDROGEN DIMETHICONE •
TIN OXIDE • PHENOXYETHANOL • FRAGRANCE • TITANIUM DIOXIDE •
IRON OXIDES • MICA •

Other Information

■ protect this product in this container
from excessive heat and direct sun. ▼

SHISEIDO
 UV PROTECTIVE LIQUID FOUNDATION
 BROAD SPECTRUM
 SPF 42

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil-UV 360™ technology.
 - Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,* cell damage* and DNA damage* linked to the formation of wrinkles and dark spots.
 - Provides a fresh touch and a dewy finish.
 - NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- * in vitro test

Drug Facts

Active ingredients	Purpose
OCTINOXATE 1.9%	Sunscreen
TITANIUM DIOXIDE 8.3%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin ▶

SHISEIDO AMERICAS CORPORATION DIST.
 NEW YORK, NY 10022
 MADE IN U.S.A.
 GLO. 11200
www.shiseido.com

Drug Facts (continued)

Warnings

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions


For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses ▶

Drug Facts (continued)

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Light Ochre

SHI SEIDO

Anti-Aging Foundation

42

UV Protective
Liquid Foundation

BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT

SUNSCREEN

30mL 1 FL. OZ.

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

Light Ochre
(SP30)

SHISEIDO

Anti-Aging Foundation



42

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT
SUNSCREEN

30mL 1 FL. OZ.

Drug Facts (continued)

Directions

■ children under 6 months: Ask a doctor

Inactive ingredients

WATER • DIMETHICONE • TRIMETHYLSILOXYSILICATE •
ISODODECANE • POLYMETHYL METHACRYLATE • CYCLOMETHICONE •
SD ALCOHOL 40-B • PEG/PPG-19/19 DIMETHICONE •
HYDROGENATED POLYISOBUTENE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
PEG/PPG-14/7 DIMETHYL ETHER •
SAXIFRAGA SARMENTOSA EXTRACT • SODIUM HYALURONATE •
SOPHORA ANGUSTIFOLIA ROOT EXTRACT • ALUMINUM HYDROXIDE •
ALUMINUM DISTEARATE • ALUMINA • ISOSTEARIC ACID •
TRIETHOXYCAPRYLYLSILANE • TRISODIUM EDTA • SILICA •
POLYSILICONE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE •
HYDRATED SILICA • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL •
DISTEARYLDIMONIUM CHLORIDE • BHT •
SYZYGIUM JAMBOS LEAF EXTRACT • HYDROGEN DIMETHICONE •
TIN OXIDE • PHENOXYETHANOL • FRAGRANCE • TITANIUM DIOXIDE •
IRON OXIDES • MICA •

Other Information

■ protect this product in this container
from excessive heat and direct sun. ▼

SHISEIDO
UV PROTECTIVE LIQUID FOUNDATION
BROAD SPECTRUM
SPF 42

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil-UV 360™ technology.
 - Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,* cell damage* and DNA damage* linked to the formation of wrinkles and dark spots.
 - Provides a fresh touch and a dewy finish.
 - NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- * in vitro test

Drug Facts

Active ingredients	Purpose
OCTINOXATE 1.9%	Sunscreen
TITANIUM DIOXIDE 8.3%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10022
MADE IN U.S.A.
G.L.O. 11201
www.shiseido.com

Drug Facts (continued)

Warnings

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions


For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses ▶

Drug Facts (continued)

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Medium Beige

SHI SEIDO

Anti-Aging Foundation

42

UV Protective

Liquid Foundation

BROAD SPECTRUM

SPF 42

WATER RESISTANT

(80 MINUTES)

TRANSFER RESISTANT

SUNSCREEN

30mL 1 FL. OZ.

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

Medium Beige
(SP60)

Anti-Aging Foundation



42

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT
SUNSCREEN

30mL 1 FL. OZ.

SHISEIDO

Drug Facts (continued)

Directions

■ children under 6 months: Ask a doctor

Inactive ingredients

WATER • DIMETHICONE • TRIMETHYLSILOXYSILICATE •
ISODODECANE • POLYMETHYL METHACRYLATE • CYCLOMETHICONE •
SD ALCOHOL 40-B • PEG/PPG-19/19 DIMETHICONE •
HYDROGENATED POLYISOBUTENE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
PEG/PPG-14/7 DIMETHYL ETHER •
SAXIFRAGA SARMENTOSA EXTRACT • SODIUM HYALURONATE •
SOPHORA ANGUSTIFOLIA ROOT EXTRACT • ALUMINUM HYDROXIDE •
ALUMINUM DISTEARATE • ALUMINA • ISOSTEARIC ACID •
TRIETHOXYCAPRYLYLSILANE • TRISODIUM EDTA • SILICA •
POLYSILOXANE-2 • ALCOHOL • SODIUM MAGNESIUM SULFATE •
HYDRATED SILICA • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL •
DISTEARYLDIMONIUM CHLORIDE • BHT •
SYZYGIUM JAMBOS LEAF EXTRACT • HYDROGEN DIMETHICONE •
TIN OXIDE • PHENOXYETHANOL • FRAGRANCE • TITANIUM DIOXIDE •
IRON OXIDES • MICA •

Other Information

■ protect this product in this container
from excessive heat and direct sun. ▼

SHISEIDO
 UV PROTECTIVE LIQUID FOUNDATION
 BROAD SPECTRUM
 SPF 42

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil UV 360™ technology.
- Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,* cell damage* and DNA damage* linked to the formation of wrinkles and dark spots.
- Provides a fresh touch and a dewy finish.
- NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- * in vitro test

Drug Facts

Active ingredients	Purpose
OCTINOXATE 1.9%	Sunscreen
TITANIUM DIOXIDE 8.3%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin ▶

SHISEIDO AMERICAS CORPORATION DIST.
 NEW YORK, NY 10022
 MADE IN U.S.A.
 GLO. 11202
www.shiseido.com

Drug Facts (continued)

Warnings

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions


For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses ▶

Drug Facts (continued)

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Medium Ivory

SHI SEIDO

Anti-Aging Foundation

42

UV Protective

Liquid Foundation

BROAD SPECTRUM

SPF 42

WATER RESISTANT

(80 MINUTES)

TRANSFER RESISTANT

SUNSCREEN

30mL 1 FL. OZ.

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

Medium Ivory
(SP40)

Anti-Aging Foundation



42

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT
SUNSCREEN

30mL 1 FL. OZ.

Drug Facts (continued)

Directions

■ children under 6 months: Ask a doctor

Inactive ingredients

WATER • DIMETHICONE • TRIMETHYLSILOXYSILICATE •
ISODODECANE • POLYMETHYL METHACRYLATE • CYCLOMETHICONE •
SD ALCOHOL 40-8 • PEG/PPG-19/19 DIMETHICONE •
HYDROGENATED POLYISOBUTENE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
PEG/PPG-14/7 DIMETHYL ETHER •
SAXIFRAGA SARMENTOSA EXTRACT • SODIUM HYALURONATE •
SOPHORA ANGUSTIFOLIA ROOT EXTRACT • ALUMINUM HYDROXIDE •
ALUMINUM DISTEARATE • ALUMINA • ISOSTEARIC ACID •
TRIETHOXYCAPRYLYLSILANE • TRISODIUM EDTA • SILICA •
POLYSILICONE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE •
HYDRATED SILICA • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL •
DISTEARYLDIMONIUM CHLORIDE • BHT •
SYZYGIUM JAMBOS LEAF EXTRACT • HYDROGEN DIMETHICONE •
TIN OXIDE • PHENOXYETHANOL • FRAGRANCE • TITANIUM DIOXIDE •
IRON OXIDES • MICA •

Other Information

■ protect this product in this container
from excessive heat and direct sun. ▼

SHISEIDO

SHISEIDO
 UV PROTECTIVE LIQUID FOUNDATION
 BROAD SPECTRUM
 SPF 42

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil^{UV} 360TM technology.
- Formulated with Profense CELTM, a complex of ingredients that reduces enzyme activity,* cell damage* and DNA damage* linked to the formation of wrinkles and dark spots.
- Provides a fresh touch and a dewy finish.
- NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- * in vitro test

Drug Facts

Active ingredients	Purpose
OCTINOXATE 1.9%	Sunscreen
TITANIUM DIOXIDE 8.3%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin ▶

SHISEIDO AMERICAS CORPORATION DIST.
 NEW YORK, NY 10022
 MADE IN U.S.A.
 GLO. 11203
www.shiseido.com

Drug Facts (continued)

Warnings

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions


For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses ▶

Drug Facts (continued)

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Dark Beige

SHI SEIDO

Anti-Aging Foundation

42

UV Protective
Liquid Foundation

BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT

SUNSCREEN

30mL 1 FL. OZ.

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

Dark Beige

Anti-Aging Foundation



42

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT
SUNSCREEN

30mL 1 FL. OZ.

SHISEIDO

Drug Facts (continued)

Directions

■ children under 6 months: Ask a doctor

Inactive ingredients

WATER • DIMETHICONE • TRIMETHYLSILOXYLICATE •
ISODODECANE • POLYMETHYL METHACRYLATE • CYCLOMETHICONE •
SD ALCOHOL 40-8 • PEG/PPG-19/19 DIMETHICONE •
HYDROGENATED POLYISOBUTENE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
PEG/PPG-14/7 DIMETHYL ETHER •
SAXIFRAGA SARMENTOSA EXTRACT • SODIUM HYALURONATE •
SOPHORA ANGUSTIFOLIA ROOT EXTRACT • ALUMINUM HYDROXIDE •
ALUMINUM DISTEARATE • ALUMINA • ISOSTEARIC ACID •
TRIETHOXYCAPRYLYLSILANE • TRISODIUM EDTA • SILICA •
POLYSILOXANE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE •
HYDRATED SILICA • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL •
DISTEARYLDIMONIUM CHLORIDE • BHT •
SYZYGIUM JAMBOS LEAF EXTRACT • HYDROGEN DIMETHICONE •
TIN OXIDE • PHENOXYETHANOL • FRAGRANCE • TITANIUM DIOXIDE •
IRON OXIDES • MICA •

Other Information

■ protect this product in this container
from excessive heat and direct sun. ▼

SHISEIDO
 UV PROTECTIVE LIQUID FOUNDATION
 BROAD SPECTRUM
 SPF 42

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil^{UV} 360TM technology.
- Formulated with Profense CELTM, a complex of ingredients that reduces enzyme activity,* cell damage* and DNA damage* linked to the formation of wrinkles and dark spots.
- Provides a fresh touch and a dewy finish.
- NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- * in vitro test

Drug Facts

Active ingredients	Purpose
OCTINOXATE 1.9%	Sunscreen
TITANIUM DIOXIDE 8.3%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin ▶

SHISEIDO AMERICAS CORPORATION DIST.
 NEW YORK, NY 10022
 MADE IN U.S.A.
 GLO. 11204 |
 www.shiseido.com

Drug Facts (continued)

Warnings

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions


For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses ▶

Drug Facts (continued)

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - Dark Ivory

SHI SEIDO

Anti-Aging Foundation

42

UV Protective
Liquid Foundation

BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT

SUNSCREEN

30mL 1 FL. OZ.

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

Dark Ivory
(SP70)

Anti-Aging Foundation



42

UV Protective
Liquid Foundation
BROAD SPECTRUM
SPF 42

WATER RESISTANT
(80 MINUTES)
TRANSFER RESISTANT
SUNSCREEN

30mL 1 FL. OZ.

SHISEIDO

Drug Facts (continued)

Directions

■ children under 6 months: Ask a doctor

Inactive ingredients

WATER • DIMETHICONE • TRIMETHYLSILOXYSILICATE •
DODODECANE • POLYMETHYL METHACRYLATE • CYCLOMETHICONE •
SD ALCOHOL 40-8 • PEG/PPG-19/19 DIMETHICONE •
HYDROGENATED POLYISOBUTENE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
PEG/PPG-14/7 DIMETHYL ETHER •
SAXIFRAGA SARMENTOSA EXTRACT • SODIUM HYALURONATE •
SOPHORA ANGUSTIFOLIA ROOT EXTRACT • ALUMINUM HYDROXIDE •
ALUMINUM DISTEARATE • ALUMINA • ISOSTEARIC ACID •
TRIETHOXYCAPRYLYLSILANE • TRISODIUM EDTA • SILICA •
POLYSILOXANE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE •
HYDRATED SILICA • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL •
DISTEARYLDIMONIUM CHLORIDE • BHT •
SYZYGIUM JAMBOS LEAF EXTRACT • HYDROGEN DIMETHICONE •
TIN OXIDE • PHENOXYETHANOL • FRAGRANCE • TITANIUM DIOXIDE •
IRON OXIDES • MICA •

Other Information

■ protect this product in this container
from excessive heat and direct sun. ▼

SHISEIDO
 UV PROTECTIVE LIQUID FOUNDATION
 BROAD SPECTRUM
 SPF 42

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil UV 360™ technology.
 - Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,* cell damage* and DNA damage* linked to the formation of wrinkles and dark spots.
 - Provides a fresh touch and a dewy finish.
 - NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- * in vitro test

Drug Facts

Active ingredients	Purpose
OCTINOXATE 1.9%	Sunscreen
TITANIUM DIOXIDE 8.3%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin ▶

SHISEIDO AMERICAS CORPORATION DIST.
 NEW YORK, NY 10022
 MADE IN U.S.A.
 GLO. 11205
www.shiseido.com

Drug Facts (continued)

Warnings

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions


For sunscreen use:

- apply liberally 15 minutes before sun exposure
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses ▶

Drug Facts (continued)

Questions or Comments?

Call toll free 1-800-906-7503



SHISEIDO UV PROTECTIVE FOUNDATION LIGHT BEIGE

octinoxate and titanium dioxide cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-130
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	644 mg in 33.9 g
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2814 mg in 33.9 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
TRIMETHYLSILOXYSILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)	
ISODODECANE (UNII: A8289P68Y2)	
CYCLOMETHICONE (UNII: NMQ347994Z)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
HYDROGENATED POLYBUTENE (1300 MW) (UNII: 7D1YQ9Y5EZ)	
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)	
SAXIFRAGA STOLONIFERA LEAF (UNII: O3TMV4903H)	
HYALURONATE SODIUM (UNII: YSE9PPT4TH)	
SOPHORA FLAVESCENS ROOT (UNII: IYR6K8KQ5K)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)	
ALUMINUM OXIDE (UNII: LM26O6933)	
ISO STEARIC ACID (UNII: X33R8U0062)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
EDETATE TRISODIUM (UNII: 420IP921MB)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ALCOHOL (UNII: 3K9958V90M)	
HYDRATED SILICA (UNII: Y6O7T4G8P9)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
GLYCERIN (UNII: PDC6A3C0OX)	

.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DISTEARYLDIMONIUM CHLORIDE (UNII: OM9573ZX3X)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
SYZYGIUM JAMBOS LEAF (UNII: 407Z4W5LFF)
STANNIC OXIDE (UNII: KM7N50LOS6)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-130-10	1 in 1 CARTON	02/01/2014	
1		33.9 g in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	02/01/2014	

SHISEIDO UV PROTECTIVE FOUNDATION LIGHT IVORY

octinoxate and titanium dioxide cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-131
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	644 mg in 33.9 g
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2814 mg in 33.9 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
TRIMETHYLSILOXYSILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)	
ISODODECANE (UNII: A8289P68Y2)	
CYCLOMETHICONE (UNII: NMQ347994Z)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
HYDROGENATED POLYBUTENE (1300 MW) (UNII: 7D1YQ9Y5EZ)	
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)	

SAXIFRAGA STOLONIFERA LEAF (UNII: O3TMV4903H)
HYALURONATE SODIUM (UNII: YSE9PPT4TH)
SOPHORA FLAVESCENS ROOT (UNII: IYR6K8KQ5K)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
ALUMINUM DISTEARATE (UNII: 7P1HP1B9U)
ALUMINUM OXIDE (UNII: LM26O6933)
ISOSTEARIC ACID (UNII: X33R8U0062)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
EDETATE TRISODIUM (UNII: 420IP921MB)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
ALCOHOL (UNII: 3K9958V90M)
HYDRATED SILICA (UNII: Y6O7T4G8P9)
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)
GLYCERIN (UNII: PDC6A3C0OX)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DISTEARYLDIMONIUM CHLORIDE (UNII: OM9573ZX3X)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
SYZYGIUM JAMBOS LEAF (UNII: 407Z4W5LFF)
STANNIC OXIDE (UNII: KM7N50LOS6)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-131-10	1 in 1 CARTON	02/01/2014	
1		33.9 g in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	02/01/2014	

SHISEIDO UV PROTECTIVE FOUNDATION LIGHT OCHRE

octinoxate and titanium dioxide cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-132
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	644 mg in 33.9 g
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2814 mg in 33.9 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y10)	
TRIMETHYLSILOXYLSILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)	
ISODODECANE (UNII: A8289P68Y2)	
CYCLOMETHICONE (UNII: NMQ347994Z)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
HYDROGENATED POLYBUTENE (1300 MW) (UNII: 7D1YQ9Y5EZ)	
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)	
SAXIFRAGA STOLONIFERA LEAF (UNII: O3TMV4903H)	
HYALURONATE SODIUM (UNII: YSE9PPT4TH)	
SOPHORA FLAVESCENS ROOT (UNII: IYR6K8KQ5K)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)	
ALUMINUM OXIDE (UNII: LM26O6933)	
ISO STEARIC ACID (UNII: X33R8U0062)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
EDETATE TRISODIUM (UNII: 420IP921MB)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ALCOHOL (UNII: 3K9958V90M)	
HYDRATED SILICA (UNII: Y6O7T4G8P9)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
GLYCERIN (UNII: PDC6A3C0OX)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DISTEARYLDIMONIUM CHLORIDE (UNII: OM9573ZX3X)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SYZYGIUM JAMBOS LEAF (UNII: 407Z4W5LFF)	
STANNIC OXIDE (UNII: KM7N50LOS6)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-132-10	1 in 1 CARTON	02/01/2014	
1		33.9 g in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	02/01/2014	

SHISEIDO UV PROTECTIVE FOUNDATION MEDIUM BEIGE

octinoxate and titanium dioxide cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-133
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	644 mg in 33.9 g
TITANIUM DIOXIDE (UNII: 15FIX9 V2JP) (TITANIUM DIOXIDE - UNII:15FIX9 V2JP)	TITANIUM DIOXIDE	2814 mg in 33.9 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
TRIMETHYLSILOXYLSILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)	
ISODODECANE (UNII: A8289P68Y2)	
CYCLOMETHICONE (UNII: NMQ347994Z)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
HYDROGENATED POLYBUTENE (1300 MW) (UNII: 7D1YQ9Y5EZ)	
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)	
SAXIFRAGA STOLONIFERA LEAF (UNII: O3TMV4903H)	
HYALURONATE SODIUM (UNII: YSE9PPT4TH)	
SOPHORA FLAVESCENS ROOT (UNII: IYR6K8KQ5K)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALUMINUM DISTEARATE (UNII: 7P1HP1B9U)	
ALUMINUM OXIDE (UNII: LM26O6933)	
ISOSTEARIC ACID (UNII: X33R8U0062)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
EDETATE TRISODIUM (UNII: 420IP921MB)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ALCOHOL (UNII: 3K9958V90M)	
HYDRATED SILICA (UNII: Y6O7T4G8P9)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
GLYCERIN (UNII: PDC6A3C0OX)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DISTEARYLDIMONIUM CHLORIDE (UNII: OM9573ZX3X)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SYZYGIUM JAMBOS LEAF (UNII: 407Z4W5LFF)	
STANNIC OXIDE (UNII: KM7N50LOS6)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

FERROSO FERRIC OXIDE (UNII: XM0M87F357)

MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-133-10	1 in 1 CARTON	02/01/2014	
1		33.9 g in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	02/01/2014	

SHISEIDO UV PROTECTIVE FOUNDATION MEDIUM IVORY

octinoxate and titanium dioxide cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-134
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	644 mg in 33.9 g
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2814 mg in 33.9 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
TRIMETHYLSILOXY SILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)	
ISODODECANE (UNII: A8289P68Y2)	
CYCLOMETHICONE (UNII: NMQ347994Z)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
HYDROGENATED POLYBUTENE (1300 MW) (UNII: 7D1YQ9Y5EZ)	
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)	
SAXIFRAGA STOLONIFERA LEAF (UNII: O3TMV4903H)	
HYALURONATE SODIUM (UNII: YSE9PPT4TH)	
SOPHORA FLAVESCENS ROOT (UNII: IYR6K8KQ5K)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)	
ALUMINUM OXIDE (UNII: LM26O6933)	
ISO STEARIC ACID (UNII: X33R8U0062)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	

EDETATE TRISODIUM (UNII: 420IP921MB)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
ALCOHOL (UNII: 3K9958V90M)
HYDRATED SILICA (UNII: Y6O7T4G8P9)
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)
GLYCERIN (UNII: PDC6A3C0OX)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DISTEARYLDIMONIUM CHLORIDE (UNII: OM9573ZX3X)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
SYZYGIUM JAMBOS LEAF (UNII: 407Z4W5LFF)
STANNIC OXIDE (UNII: KM7N50LOS6)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-134-10	1 in 1 CARTON	02/01/2014	
1		33.9 g in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	02/01/2014	

SHISEIDO UV PROTECTIVE FOUNDATION DARK BEIGE

octinoxate and titanium dioxide cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-135
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	644 mg in 33.9 g
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2814 mg in 33.9 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	

TRIMETHYLSILOXYSILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)
ISODODECANE (UNII: A8289P68Y2)
CYCLOMETHICONE (UNII: NMQ347994Z)
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)
HYDROGENATED POLYBUTENE (1300 MW) (UNII: 7D1YQ9Y5EZ)
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)
SAXIFRAGA STOLONIFERA LEAF (UNII: O3TMV4903H)
HYALURONATE SODIUM (UNII: YSE9PPT4TH)
SOPHORA FLAVESCENS ROOT (UNII: IYR6K8KQ5K)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
ALUMINUM DISTEARATE (UNII: 7P1HP1B9UI)
ALUMINUM OXIDE (UNII: LMI26O6933)
ISOSTEARIC ACID (UNII: X33R8U0062)
TRIETHOXYCAPRYLSILANE (UNII: LDC331P08E)
EDETATE TRISODIUM (UNII: 420IP921MB)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
ALCOHOL (UNII: 3K9958V90M)
HYDRATED SILICA (UNII: Y6O7T4G8P9)
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)
GLYCERIN (UNII: PDC6A3C0OX)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DISTEARYLDIMONIUM CHLORIDE (UNII: OM9573ZX3X)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
SYZYGIUM JAMBOS LEAF (UNII: 407Z4W5LFF)
STANNIC OXIDE (UNII: KM7N50LOS6)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-135-10	1 in 1 CARTON	02/01/2014	
1		33.9 g in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	02/01/2014	

SHISEIDO UV PROTECTIVE FOUNDATION DARK IVORY
 octinoxate and titanium dioxide cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-136
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	644 mg in 33.9 g
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2814 mg in 33.9 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
TRIMETHYLSILOXYSILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)	
ISODODECANE (UNII: A8289P68Y2)	
CYCLOMETHICONE (UNII: NMQ347994Z)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
HYDROGENATED POLYBUTENE (1300 MW) (UNII: 7D1YQ9Y5EZ)	
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)	
SAXIFRAGA STOLONIFERA LEAF (UNII: O3TMV4903H)	
HYALURONATE SODIUM (UNII: YSE9PPT4TH)	
SOPHORA FLAVESCENS ROOT (UNII: IYR6K8KQ5K)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
ALUMINUM DISTEARATE (UNII: 7P1HP1B9U)	
ALUMINUM OXIDE (UNII: LM26O6933)	
ISOSTEARIC ACID (UNII: X33R8U0062)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
EDETATE TRISODIUM (UNII: 420IP921MB)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ALCOHOL (UNII: 3K9958V90M)	
HYDRATED SILICA (UNII: Y6O7T4G8P9)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
GLYCERIN (UNII: PDC6A3C0OX)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DISTEARYLDIMONIUM CHLORIDE (UNII: OM9573ZX3X)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SYZYGIUM JAMBOS LEAF (UNII: 407Z4W5LFF)	
STANNIC OXIDE (UNII: KM7N50LOS6)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-136-10	1 in 1 CARTON	02/01/2014	

1	33.9 g in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	02/01/2014	

Labeler - SHISEIDO AMERICAS CORPORATION (193691821)

Establishment			
Name	Address	ID/FEI	Business Operations
SHISEIDO AMERICA INC.		782677132	manufacture(58411-130, 58411-131, 58411-132, 58411-133, 58411-134, 58411-135, 58411-136), analysis(58411-130, 58411-131, 58411-132, 58411-133, 58411-134, 58411-135, 58411-136)

Revised: 12/2018

SHISEIDO AMERICAS CORPORATION