DIPHENHYDRAMINE HCL- diphenhydramine hcl liquid Akron Pharma Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Akron Pharma, Inc.

Active ingredient (in each teaspoonful (5 mL))

Diphenhydramine HCl 12.5 mg

Purpose

Antihistamine

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat

Warnings

Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- marked drowsiness may occur
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in 24 hours
- mL = milliliter; FL OZ = fluid ounce
- find right dose on chart below
- take every 4 to 6 hours, or as directed by a doctor

| Age | Dose |
|---------------------------|---|
| | 2 - 4 teaspoonsful (25 mg to 50 mg) |
| children 6 to 11 years | 1 - 2 teaspoonsful (12.5 mg to 25 mg) |
| children 2 to 5 Vears | do not use unless directed by a doctor |
| children under 2 years | do not use |

Other information

- each teaspoonful (5 mL) contains: sodium 5 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

Bubble Gum Flavor, Citric Acid, Purified Water, Sodium Benzoate, Sodium Citrate, Sodium Saccharin, Sorbitol Solution 70% & Sucrose.

Questions or comments?

Please Call 1(877) 225-6999

Manufactured for:

Akron Pharma, Inc, Fairfield, NJ 07004 Manufactured in U.S.A

Drug Facts (continued)

| find | the righ | t dose on | the chart | belo |
|------|----------|-----------|-------------|------|
| | | 00000011 | 1110 011011 | 0010 |

| Age (yr) | Dose (mL) |
|------------------------|--|
| children 6 to 11 years | 5 mL to 10 mL |
| children 2 to 5 years | Do not use unless directed by a doctor |
| children under 2 years | Do not use |

Other information

each 5 mL contains: sodium 8 mg
 store between 20'- 25'C (68'-77'F). Do not refrigerate
 protect from light.

Inactive ingredients

Bubble Gum Flavor, Citric Acid, Purified Water, Sodium Benzoate, Sodium Citrate, Sodium Saccharin, Sorbitol Solution 70% & Sucrose.

Ouestions or comments? Call (877) 225-6999 Monday - Friday 9AM - 5PM EST

This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Benadryl Allergy.

Lot

Exp

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING

Akrán Pharma Manufactured for Akron Pharma, Inc. Fairfield, NJ - 07004 Rev 09-19 Manufactured In USA



For Ages 6 to 11 Years Alcohol-Free

Compare to the active Ingredient in Children's

Benadryl Allergy

Dye-free Bubble Gum Flavored

4 FL OZ (118 mL)

Compare to the active

Drug Facts

Active ingredient (in each 5 mL) Purpose Diphenhydramine HCI 12.5 mg Antihistamine Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies. sneezing =runny nose = tichy,watery eyes = tichy of the nose or throat

Warnings

NDC 71399-0125-4

Akrän Pharma

NDC 71399-0125-8

Do not use with any other product containing diphenhydramine, even one used on the skin to make a child sleepy

Ask a doctor before use if the child has a breathing problem such as chronic bronchitis

glaucoma
 a sodium-restricted diet

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers.

When using this product marked drowsiness may occur sedatives and tranquilizers may increase drowsiness excitability may occur, especially in children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

Directions -do not take more than 6 doses in any 24 hours -take every 4 to 6 hours, or as directed by a doctor -measure only with dosing cup provided. Do not use any other dosing device. -mL=milliliter - keep dosing cup with product

Drug Facts (continued)

| find the right dose on the chart below | | | | | |
|--|------|------|----|---|---|
| Ade | (vr) | Dose | mL |) | _ |

| | , |
|------------------------|--|
| children 6 to 11 years | 5 mL to 10 mL |
| children 2 to 5 years | Do not use unless directed by a doctor |
| children under 2 years | Do not use |

Other information

each 5 mL contains: sodium 8 mg
 store between 20'- 25'C (68'-77'F). Do not refrigerate
 protect from light.

1

Inactive ingredients

Bubble Gum Flavor, Citric Acid, Purified Water, Sodium Benzoate, Sodium Citrate, Sodium Saccharin, Sorbitol Solution 70% & Sucrose.

Questions or comments?

Call (877) 225-6999 Monday - Friday 9AM - 5PM EST * This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Benadry Allergy.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING

Akráin Pharma Manufactured for Akron Pharma, Inc., Fairfield, NJ - 07004

Manufactured In USA

Rev 09-19



Ingredient in Children's Benadryl Allergy **Dye-Free**



Antihistamine

Diphenhydramine Hydrochloride 12.5 mg/5 mL Oral Solution

Relieves

Runny Nose • Sneezing • Itchy, Watery Eyes • Itchy Throat For Ages 6 to 11 Years Alcohol-Free

Dye-free Bubble Gum Flavored



Drug Facts Active ingredient (ineach 5 mL)

Purpose

•

Diphenhydramine HCI 12.5 mg Antihistamine Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies.
 sneezing = runny nose = ltchy, watery eyes = itchy of the nose or throat

Warnings

Do not use
 when any other product containing diphenhydramine,
 even one used on the skin
 to make a child sleepy

- Ask a doctor before use if the child has = a breathing problem such as chronic bronchitis glaucoma = a socium-restricted diet

Ask a doctor or pharmacist before use if the child is taking sedatives or tranquilizers.

When using this product marked drowsiness may occur sedatives and tranquilizers may increase drowsiness excitability may occur, especially in children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

do not take more than 6 doses in any 24 hours

- take every 4 to 6 hours, or as directed by a doctor
 measure only with dosing cup provided. Do not use any other dosing device.
 mL=milliliter keep dosing cup with product

8 FL OZ (236 mL)

Akrän Pharma

| Drug Facts | NDC 71399-0125-6 *Compare to the active Ingredient in Children's | | Drug Facts (Co | ontinued) | |
|--|---|-------------------------------|--|---|--|
| Active ingredient Purpose | | Benadryl [®] Allergy | | | |
| (in each 5 mL) | | | Age (yr) | Dose (mL) | |
| Diphenhydramine HCI 12.5 mgAntihistamine | Dye-free | | children 6 to 11 years | 5 mL to 10 mL | |
| Uses temporarily relieves these symptoms due to | | | children 2 to 5 years | Do not use unless directed by a doctor | |
| hay fever or other upper respiratory allergies. ■ sneezing ■ runny nose ■ ltchy, watery eyes ■ itchy | | | children under 2 years | | |
| nose or throat | Childre | 'N S | Other Informati | 0.0 | |
| Warnings | | | each 5 mL contai | | |
| ■ with any other product containing diphenhydramine, | | | store between 20° | | |
| even one used on the skin | Alle | | Do not refrigerate protect from light | | |
| to make a child sleepy Ask a doctor before use if the child has | | | Inactive Ingred | ents | |
| a breathing problem such as chronic bronchitis | Antihistam | ho | Bubble Gum Flavor, (| Sitric Acid, Purified Water, | |
| glaucoma a sodium-restricted diet | Entrement | ШЭ | Sodium Benzoate, So Sorbitol Solution 70% | odium Citrate, Sodium Saccharin | |
| Ask a doctor or pharmacist before use if the child is | Diphenhydramine | Hydrochloride | | <i>mments?</i> 1-877-225-6999 | |
| taking sedatives or tranquilizers. | 12.5 mg/5 mL Ora | | | NOT USE IF PRINTED SAFETY SEAL | |
| When using this product | Relieves: | | | NDER CAP IS BROKEN OR MISSING. | |
| marked drowsiness may occur sedatives and tranquilizers may increase drowsiness | Runny Nose Sneezin | na a ltchy | *This product is not man | nufactured or distributed by McNeil listributor of Benadryl Allergy. | |
| excitability may occur, especially in children | Watery Eyes Itchy T | | | istributor of bonduryt natogy. | |
| Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center | | | | | |
| (1-800-222-1222) right away. | For Ages 6 to 11 year | | | | |
| Directions | Alcohol-Free | | Rev.:07/22 | | |
| do not take more than 6 doses in any 24 hours | Bubble Gum Flavo | red | Akrájn Pharma | | |
| take every 4 to 6 hours, or as directed by a doctor measure only with dosing cup provided. Do not use | | | Manufactured for: Akron Pharma, Inc. | | |
| any other dosing device. | | Made in USA | Fairfield, NJ 07004 | | |
| ■ mL=milliliter. ■ keep dosing cup with product | Akr 🗐 n Pharma 🛛 1 | 6 FL.OZ (473 mL) | www.akronpharma.com | n 3 ∎ 71399 01256 2 | |

| DIPHENHYDRAMINE | HCL | | | | | |
|---------------------------------|----------------|----------------------------------|---------------|--------------------|----------|--|
| diphenhydramine hcl liquid | | | | | | |
| | | | | | | |
| Product Information | | | | | | |
| Product Type | HUMAN OTC DRUG | Item Code (| Source) | NDC:713 | 99-0125 | |
| Route of Administration | ORAL | | | | | |
| | | | | | | |
| | | | | | | |
| Active Ingredient/Active | Moiety | | | | | |
| Ingredient Name Basis of Stre | | | | | Strength | |
| DIPHENHYDRAMINE HYDROCHL | | DIPHENHYDRAMINE HYDROCHLORIDE | | 12.5 mg in 5 ml | | |
| (DIPHENHYDRAMINE - UNII:8GTS825 | 583™) | | HIDROCHLORIDE | | IN 5 ML | |
| | | | | | | |
| Inactive Ingredients | | | | | | |
| Ingredient Name | | | | | Strength | |
| ANHYDROUS CITRIC ACID (UNII: | XF417D3PSL) | | | | | |
| WATER (UNII: 059QF0K00R) | | | | | | |
| SODIUM BENZOATE (UNII: 0J245) | FE5EU) | | | | | |
| SODIUM CITRATE (UNII: 1Q73Q2J | ULR) | | | | | |
| SACCHARIN SODIUM (UNII: SB8Z | UX40TY) | | | | | |
| SORBITOL (UNII: 506T60A25R) | | | | | | |

SUCROSE (UNII: C151H8M554)

| P | roduct Cha | racterist | ics | | | | |
|---------------------------|----------------------|--|--|--------------|-------------------------|------------------|---|
| С | olor | | | Scor | e | | |
| SI | hape | | | Size | | | |
| | avor | | BUBBLE GUM | Imprint Code | | | |
| С | ontains | | | | | | |
| | | | | | | | |
| P | ackaging | | | | | | |
| # | ltem Code | Package Description | | | Marketing Start Date | Marketin Dat | - |
| 1 | NDC:71399- 0125-4 | 118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | | 10/25/2019 | | |
| 2 | NDC:71399- 0125-8 | 236 mL in Combinatio | 1 BOTTLE, PLASTIC; Type 0: Not a on Product | | 10/25/2019 | | |
| 3 | NDC:71399- 0125-6 | 473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | | 02/01/2023 | | |
| | | | | | | | |
| M | larketing | Inforn | nation | | | | |
| Marketing App Category | | Арр | blication Number or Monograp Citation | h | Marketing Start Date | Marketin Date | - |
| | TC MONOGRAPH | part34 | 1 |] | 0/25/2019 | | |

Labeler - Akron Pharma Inc. (067878881)

Revised: 2/2023

Akron Pharma Inc.