

## **MEIJER CLEAR ZINC SPF 50 - octocrylene, zinc oxide lotion**

**Meijer Distribution Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Drug Facts**

#### **Active ingredients**

Octocrylene 4.0%

Zinc Oxide 5.0%

#### **Purpose**

Sunscreen

#### **Uses**

- Helps prevent sunburn
- If used as directed with other sun protection measures (see Directions), decrease the risk of skin cancer and early skin aging caused by the sun.

#### **Warnings**

For external use only. Do not use on damaged or broken skin.

#### **When using this product**

Keep out of eyes. Rinse with water to remove.

#### **Stop use and ask a doctor**

if rash occurs

#### **Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away. May stain some fabrics.

#### **Directions**

- apply liberally 15 minutes before sun exposure
- reapply
- after 80 minutes of swimming or sweating
- immediately after towel drying
- at least every 2 hours
- **Sun Protection Measures:** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats and sunglasses
  - children under 6 months of age: Ask a doctor

**Inactive ingredients**

Water, Ethylhexyl Stearate, Diisopropyl Adipate, Cetyl Dimethicone, Polyglyceryl-4 Isostearate, Cetyl PEG/PPG-10/1 Dimethicone, Hexyl Laurate, Beeswax, Isohexadecane, Sodium Chloride, Hydrogenated Castor Oil, Silica, Triethoxycaprylylsilane, Dimethicone, Tocopherol, Disodium EDTA, Aloe Barbadensis Leaf Juice Powder, Sodium Ascorbyl Phosphate, Phenoxyethanol, Methylparaben, Propylparaben.

**Other Information**

- protect this product from excessive heat and direct sun

**Principal Display Panel**

meijer

Clear Zinc

Sunscreen lotion

50

Broad Spectrum SPF 50

Water Resistant (80 Minutes)

Goes on Clear Non-Greasy

Dermatologist Tested UVA/UVB Protection

FPO

BLUE CIRCLE REPRESENTS JAR LID

RED DOTTED LINE REPRESENTS OUTSIDE EDGE OF CLEAR BLISTER

(DO NOT PRINT)

NET WT. 1 OZ. (28 g)

**meijer**

**Clear Zinc Sunscreen**  
lotion

**50**  
Broad Spectrum  
SPF 50

Water Resistant (80 Minutes)    Goes on Clear Non-Greasy    Dermatologist Tested UVA/UVB Protection

**FPO**  
**BLUE CIRCLE**  
**REPRESENTS JAR LID**  
**RED DOTTED LINE**  
**REPRESENTS OUTSIDE EDGE**  
**OF CLEAR BLISTER**  
**(DO NOT PRINT)**

**NET WT.**  
**1 OZ. (28 g)**

**Drug Facts**

<b>Active ingredients</b> Octocrylene 4.0%, Zinc Oxide 5.0%	<b>Purpose</b> Sunscreen
<b>Uses</b> • helps prevent sunburn • If used as directed with other sun protection measures (see <i>Directions</i> ), decreases the risk of skin cancer and early skin aging caused by the sun	
<b>Warnings</b> For external use only Do not use on damaged or broken skin When using this product keep out of eyes. Rinse with water to remove. Stop use and ask a doctor if rash occurs Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away. • May stain some fabrics.	
<b>Directions</b> • apply liberally 15 minutes before sun exposure • reapply after 80 minutes of swimming or sweating • immediately after towel drying • at least every 2 hours • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. - 2 p.m. • wear long-sleeve shirts, pants, hats and sunglasses • children under 6 months of age. Ask a doctor	
<b>Inactive ingredients</b> Water, Ethylhexyl Stearate, Diisopropyl Adipate, Cetyl Dimethicone, Polyglyceryl-4 Isocostearate, Cetyl PEG/PPG-10/1 Dimethicone, Hexyl Laurate, Beeswax, Isohexadecane, Sodium Chloride, Hydrogenated Castor Oil, Silica, Triethoxycaprylylsilane, Dimethicone, Tocopherol, Disodium EDTA, Aloe Barbadosis Leaf Juice Powder, Sodium Ascorbyl Phosphate, Phenoxyethanol, Methylparaben, Propylparaben.	
<b>Other information</b> • protect this product from excessive heat and direct sun	

**Directions**  
• apply liberally 15 minutes before sun exposure  
• reapply after 80 minutes of swimming or sweating  
• immediately after towel drying • at least every 2 hours • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. - 2 p.m. • wear long-sleeve shirts, pants, hats and sunglasses • children under 6 months of age. Ask a doctor

**Inactive ingredients** Water, Ethylhexyl Stearate, Diisopropyl Adipate, Cetyl Dimethicone, Polyglyceryl-4 Isocostearate, Cetyl PEG/PPG-10/1 Dimethicone, Hexyl Laurate, Beeswax, Isohexadecane, Sodium Chloride, Hydrogenated Castor Oil, Silica, Triethoxycaprylylsilane, Dimethicone, Tocopherol, Disodium EDTA, Aloe Barbadosis Leaf Juice Powder, Sodium Ascorbyl Phosphate, Phenoxyethanol, Methylparaben, Propylparaben.

**Other information** • protect this product from excessive heat and direct sun

7 1333 8933 109

Made in USA  
**DIST. BY MEIJER DISTRIBUTION, INC.**  
2929 WALKER AVE NW  
GRAND RAPIDS, MI 49544  
www.meijer.com

93007814

## MEIJER CLEAR ZINC SPF 50

octocrylene, zinc oxide lotion

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-038
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	4 g in 100 g
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	5 g in 100 g

## Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
ETHYLHEXYL STEARATE (UNII: EG3PA2K3K5)	
DIISOPROPYL ADIPATE (UNII: P7E6YFV72X)	
POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7)	
HEXYL LAURATE (UNII: 4CG9F9W01Q)	
YELLOW WAX (UNII: 2ZA36H0S2V)	
ISOHEXADECANE (UNII: 918X1OUF1E)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
HYDROGENATED CASTOR OIL (UNII: ZF94AP8MEY)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
TOCOPHEROL (UNII: R0ZB2556P8)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
SODIUM ASCORBYL PHOSPHATE (UNII: 836SJG51DR)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-038-81	28 g in 1 BOTTLE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	07/12/2013	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 7/2013

Meijer Distribution Inc