SAFCO SENSICAINE ULTRA TOPICAL ANESTHETIC GEL- benzocaine gel Safco Dental Supply Co.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredients

Benzocaine 200mg (in each g)

Purpose

Oral Anesthetic

Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use. Keep out of reach of children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other Information


Store at 59°-86°F (15°-30°C). Protect from freezing.

Inactive ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?

safcodental.com * 1-800-621-2178



benzocaine gel

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:67239-0220

Route of Administration DENTAL

Active Ingredient/Active Moiety

	Ingredient Name	Basis of Strength	Strength
l	BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients		
Ingredient Name	Strength	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)		
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
FD&C RED NO. 3 (UNII: PN2Z H5LOQY)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)		
D&C RED NO. 28 (UNII: 767IP0Y5NH)		
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)		

Product Characteristics			
Color	green	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:67239- 0220-1	30 g in 1 JAR; Type 0: Not a Combination Product	03/05/2019		

Distributed by: Safco Dental Supply Co.

Buffalo Grove, IL 60089 800-621-2178 Made in USA www.safcodental.com

For professional use only

OPEN



SensiCaine Ultra Topical Anesthetic Gel

1 oz. (29.6 ml)

99091779 Rev 03/2019

Questions or comments? safcodental.com • 800-621-2178

Contains 20% Benzocaine

(tartrazine), yellow #6, as a color

additive.

blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5,

sodium saccharin. May contain

flavoring, PEG 3350, PEG 400,

nactive ingredients

Mint REF 102-0450 NDC 67239-0220-1

Gluten Free

mouth symptoms do not improve f swelling, rash or fever develops redness persists or worsens; or directed by a physician. If sore in 7 days; irritation, pain, or see your physician promptly for teething

For oral mucosal use only, as

Oral Anesthetic

emporary relief of pain due to minor dental procedures.

directed by dentist. For the

Benzocaine 200 mg (in each g)

Active ingredients

Drug Facts

in children under 2 years of age When using this product

Do not exceed recommender contact occurs, flush with water dosage. If more than used for pain is accidentally swallowed Avoid contact with eyes. If

warning: Use of this product

Methemoglobinemia

Warnings

Poison Control Center right away get medical help or contact a

if pregnant or breast feeding Keep out of reach of children ask a physician before use.

may cause methemoglobinemia a serious condition that must be attention if you or a child in your occur even if you have used this reduces the amount of oxygen product before. Stop use and carried in the blood. This can treated promptly because it seek immediate medical

pale, gray or blue colored skin care develops: cyanosis

Apply only amount needed to

Directions

the oral mucosa to prevent or

children under 2 years of age

relieve pain.

Store at 59" - 86" F (15 - 30"C).

Protect from freezing.

Other information

do not use

dizziness or lightheadedness shortness of breath rapid heart rate headache

Allergy Alert: Do not use on allergies to local anesthetics such as procaine, butacaine penzocaine or other "caine" fatigue or lack of energy patients with a history of

for more than 7 days unless Do not use anesthetics.

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	03/05/2019	

Labeler - Safco Dental Supply Co. (020417380)

Revised: 1/2022 Safco Dental Supply Co.