#### SAFCO SENSICAINE ULTRA TOPICAL ANESTHETIC GEL- benzocaine gel Safco Dental Supply Co.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

# **Active Ingredients**

Benzocaine 200mg (in each g)

### Purpose

Oral Anesthetic

#### Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

# Warnings

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy Alert:** Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

### Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

**Do not exceed recommended dosage.** If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

# Keep out of reach of children.

#### Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

# **Other Information**


Store at 59°-86°F (15°-30°C). Protect from freezing.

#### Inactive ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

# **Questions or comments?**

safcodental.com \* 1-800-621-2178



Product Inform	ation						
					NDC		
Product Type		HUMAN OTC DRUG	Item Code (Source)		NDC:0	IDC:67239-0222	
Route of Administ	ration	DENTAL					
Active Ingredier	nt/Active	Moiety					
Ingredient Name				Basis of Stre	ngth	Strength	
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY4			48JW5)	BENZOCAINE		200 mg in 1 g	
Inactive Ingredi	ents						
		Ingredient Name				Strength	
POLYETHYLENE GLY	COL 3350	(UNII: G2M7P15E5P)					
POLYETHYLENE GLY							
SACCHARIN SODIUM							
FD&C YELLOW NO. 5							
FD&C YELLOW NO. 6							
FD&C RED NO. 3 (UN							
FD&C RED NO. 40 (U							
FD&C BLUE NO. 1 (U							
FD&C GREEN NO. 3 (							
D&C RED NO. 28 (UN							
D&C GREEN NO. 5 (U	NII: 8JORDU	8697)					
Product Charac	teristics						
Color	re	ed	Score				
Shape			Size				
Flavor		ASPBERRY	Imprint	Imprint Code			
Contains							
Packaging							
# Item Code	Pa	ckage Description	Ma	Marketing Start Ma Date		rketing End Date	
NDC:67239- 3	0	; Type 0: Not a Combination		/2019			

Distributed by: Safco Dental Supply Co. Buffalo Grave, IL 60089 = 800-621-2178 www.safcodental.com Made in USA	s	afco	
For professional use only OPEN HERE	Raspberry REF 102-0468 NDC 67239-0222-1 Gluten Free	SensiCaine Topical Anest	
directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly. • for teething • in children under 2 years of age When using this product Avoid contact with eyes. If contact occurs, flush with water. Do not excond recommonded	dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away. If pregnant or breast feeding, ask a physician before use. Xeep out of reach of children. Directions • Apply only amount needed to the oral mucosa to prevent or	relieve pain. • children under 2 years of age: do not use Other information Store at 59° - 86°F (15 - 30°C). Protect from freezing. Inactive ingredients flavoring, PEG 3350, PEG 400, sodium sacharin, Max contain	blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive. <b>Questions or comments?</b> safcodental.com + 800-621-2178
Drug Facts Active ingredients Benzocaine 200 mg (in each g) Purpose Oral Anesthetic Use For oral mucosal use only, as directed by dentist. For the temporary relief of pain due temporary relief of pain due to minor dental procedures. Warnings Use of this product	a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: pale, gray or blue colored skin	<ul> <li>(cyanosis)</li> <li>headache</li> <li>rapid heart rate</li> <li>shortness of breath</li> <li>dizziness or lightheadedness</li> <li>fatigue or lack of energy</li> <li>Allergy Alert: Do not use on patients with a history of allergies to local anesthetics</li> </ul>	benzocaline or other "caline" anesthetics. <b>Do not use</b> • for more than 7 days unless
Iarketing Information Marketing Application N Category	lumber or Monograph Citation	Marketing Start Date	Marketing Enc Date
TC monograph not part356		03/05/2019	Pate

# Labeler - Safco Dental Supply Co. (020417380)

Revised: 1/2022

Safco Dental Supply Co.