

SAFCO SENSICAINA ULTRA TOPICAL ANESTHETIC GEL- benzocaine gel
Safco Dental Supply Co.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredients

Benzocaine 200mg (in each g)

Purpose

Oral Anesthetic

Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Keep out of reach of children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other Information

Store at 59°-86°F (15°-30°C). Protect from freezing.

Inactive ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?

safcodental.com * 1-800-621-2178

Distributed by:
Safco Dental Supply Co.
 Buffalo Grove, IL 60089 ☎ 800-621-2178
 www.safcodental.com Made in USA

safco

For professional use only

Strawberry
REF 102-0439
 NDC 67239-0223-1
 Gluten Free

SensiCaine™ Ultra
 Topical Anesthetic Gel

1 oz. (29.6 ml)
Contains 20% Benzocaine

OPEN HERE

| |
|--|
| <p>directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.</p> <ul style="list-style-type: none"> • for teething • in children under 2 years of age <p>When using this product Avoid contact with eyes. If contact occurs, flush with water.</p> <p>Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.</p> <p>If pregnant or breast feeding, ask a physician before use.</p> <p>Keep out of reach of children.</p> <p>Directions</p> <ul style="list-style-type: none"> • Apply only amount needed to the oral mucosa to prevent or relieve pain. • children under 2 years of age: do not use <p>Other information Store at 59° - 86° F (15 - 30° C). Protect from freezing.</p> <p>Inactive ingredients flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.</p> <p>Questions or comments? safcodental.com • 800-621-2178</p> <p style="text-align: right; font-size: small;">99091776 Rev 03/2019</p> |
|--|

| |
|---|
| <p>Drug Facts</p> <p>Active ingredients Benzocaine 200 mg (in each g)</p> <p>Purpose Oral Anesthetic</p> <p>Use For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.</p> <p>Warnings</p> <p>Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:</p> <ul style="list-style-type: none"> • pale, gray or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy <p>Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.</p> <p>Do not use</p> <ul style="list-style-type: none"> • for more than 7 days unless |
|---|

SAFCO SENSICAINE ULTRA TOPICAL ANESTHETIC GEL

benzocaine gel

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:67239-0223 |
| Route of Administration | DENTAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|--------------------------|-----------------|
| BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) | BENZOCAINE | 200 mg in 1 g |

Inactive Ingredients

| Ingredient Name | Strength |
|--|-----------------|
| POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) | |
| POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| FD&C YELLOW NO. 5 (UNII: I753WB2F1M) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| FD&C RED NO. 3 (UNII: PN2ZH5LOQY) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S) | |
| D&C RED NO. 28 (UNII: 767IP0Y5NH) | |
| D&C GREEN NO. 5 (UNII: 8J6RDU8L9X) | |

Product Characteristics

| | | | |
|-----------------|------------|---------------------|--|
| Color | red | Score | |
| Shape | | Size | |
| Flavor | STRAWBERRY | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|--|-----------------------------|---------------------------|
| 1 | NDC:67239-0223-1 | 30 g in 1 JAR; Type 0: Not a Combination Product | 03/05/2019 | |

Distributed by:
Safco Dental Supply Co.
 Buffalo Grove, IL 60089 ☎ 800-621-2178
 www.safcodental.com Made in USA



For professional use only

Strawberry

REF 102-0439

NDC 67239-0223-1

Gluten Free

SensiCaine™ Ultra
Topical Anesthetic Gel

1 oz. (29.6 ml)

Contains 20% Benzocaine



directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.

- for teething
- in children under 2 years of age

When using this product
 Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Keep out of reach of children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other information
 Store at 59° - 86° F (15 - 30° C). Protect from freezing.

Inactive ingredients
 flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?
 safcodental.com • 800-621-2178

99091776 Rev 03/2019

| |
|--|
| Drug Facts |
| Active ingredients Benzocaine 200 mg (in each g) |
| Purpose Oral Anesthetic |
| Use For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures. |
| Warnings Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: <ul style="list-style-type: none"> • pale, gray or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics. Do not use <ul style="list-style-type: none"> • for more than 7 days unless |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part356 | 03/05/2019 | |

Labeler - Safco Dental Supply Co. (020417380)

Revised: 1/2022

Safco Dental Supply Co.