IGNATIA AMARA- strychnos ignatii seed pellet Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Ignatia amara 6X

Ignatia amara 6X

(**contains 0.443 mg of the active ingredient per pellet)

Apprehension, Hypersensitivity To Stress*

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen

If pregnant or breast-feeding ask a health professional before use

Keep out of reach of children

Do not use if pellet dispenser seal is broken.

Contains approx 80 pellets.

How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue. *CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

lactose, sucrose

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

1-800-BOIRON-1 (1-800-264-7661), BoironUSA.com Info@boiron.com Distributed by Boiron, Inc. Newtown Square, PA 19073





Drug Facts

Active ingredient**: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).


Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Directions: Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Other information: Do not use if pellet dispenser seal is broken.

Drug Facts (continued) Inactive ingredients: lactose, sucrose



| IGNATIA AMARA | | | | | | | |
|---|---------------------------------------|------|-----------------------|-------------------|----------------------|--------------|--|
| strychnos ignatii seed pellet | | | | | | | |
| | | | | | | | |
| Product Information | | | | | | | |
| Product Type | HUMAN OTC DRUG | | Item Code (Source) ND | | | DC:0220-2613 | |
| Route of Administration | ORAL | | | | | | |
| | | | | | | | |
| | | | | | | | |
| Active Ingredient/Active | Molety | | | | | | |
| Ingredient Name | | | | | Basis of Strength | | |
| STRYCHNOS IGNATII SEED (UNI UNII:1NM3M2487K) | 1NM3M2487K) (STRYCHNOS IGNATII SEED - | | | STRYCHNOS SEED | 5 IGNATII | 6 [hp_X] | |
| | | | | | | | |
| Inactive Ingredients | | | | | | | |
| Ingredient Name | | | | | Strength | | |
| SUCROSE (UNII: C151H8M554) | | | | | | | |
| LACTOSE (UNII: J2B2A4N98G) | | | | | | | |
| | | | | | | | |
| Product Characteristics | | | | | | | |
| Color | white | Scor | e | | | | |
| Shape | ROUND | Size | | | 4mr | m | |

| Flavor | | | Imprint Code | | | | | |
|-----------------------|-----------------------|------------------------|---|--|---------------------------|-------------------------|-----------------------|--|
| Contains | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| Packaging | | | | | | | | |
| # | ltem Code | Pa | Package Description | | M | larketing Start Date | Marketing End Date | |
| 1 | NDC:0220-2613- 41 | 80 in 1 TUB Product | 1 TUBE; Type 0: Not a Combination | | 03/03/1983 | | | |
| | | | | | | | | |
| | | | | | | | | |
| Marketing Information | | | | | | | | |
| | Marketing Category | Applic | pplication Number or Monograp Citation | | h Marketing Start Date | | Marketing End Date | |
| | approved meopathic | | | | | 03/03/1983 | | |
| | | | | | | | | |

Labeler - Boiron (282560473)

Registrant - Boiron Inc. (014892269)

| Establishment | | | | | |
|---------------|---------|-----------|----------------------------|--|--|
| Name | Address | ID/FEI | Business Operations | | |
| Boiron | | 282560473 | manufacture(0220-2613) | | |

Revised: 2/2023

Boiron